

Foundations of Electro-Magnetism Slide Series

- [Major Points of Presentation](#)
- [Chung's Carbon Filament Negative Resistor](#)
- [Aharanov-Bohm Effect](#)
- [Absorption and emission reactions \(1\)](#)
- [Absorption and emission reactions \(2\)](#)
- [Anti-engine for cell's deviation reverses cell back to normal](#)
- [Asymmetric regauging produces excess force, which can be used to do work on the system](#)
- [Asymmetric regauging produces excess force, which can be used to do work on the regauging system](#)
- [Becker's bone fracture healing experiments](#)
- [Becker's theoretical control system governing regeneration](#)
- [Becker's theoretical DC control system involved with response to injury](#)
- [Geometrical Distortion of Poynting Energy Flow](#)
- [Cosmic Bootstrap: Before the Big Bang](#)
- [Bridging Enables COP > 1](#)

- [Aspects of Strong Local Symmetry](#)
- [Other Aspects of Strong Local Symmetry](#)
- [Bunge on the Status of Electrodynamics](#)
- [Bunge on the Status of Electrodynamics and Physics](#)
- [Bunge on the Status of Classical and Quantum Physics](#)
- [Fractions of BW Casualties Surviving](#)
- [First Step in Self-Promotion of Cancer Due to Hypoxia](#)
- [Differentiation and Dedifferentiation Timelines](#)
- [Evolution of Cells on Earth](#)
- [A Charged Particle \$q\$ is a Coupled System of \$m\$ and \$\emptyset\$](#)
- [The Ubiquitous Assumption: Two Asymmetrical Regaugings for Net Symmetry \(1\)](#)
- [The Ubiquitous Assumption: Two Asymmetrical Regaugings for Net Symmetry \(2\)](#)
- [The Ubiquitous Assumption: Two Asymmetrical Regaugings for Net Symmetry \(3\)](#)
- [A Charged Particle is a Coupled System](#)
- [The Anomalies in Navy Electrolyte Experiments at China Lake](#)
- [Chung's Carbon Filament Negative Resistor](#)

- [Typical circuit has about \$10^{-13}\$ energy collection efficiency](#)
- [Classical View of EM](#)
- [Curved Spacetime acts as source or sink](#)
- [Chronic Disease Puzzling](#)
- [Cancer Characteristics](#)
- [Solving a Cancer Enigma](#)
- [Cancer: Characteristics](#)
- [Nonlinear Optics Distortion Correction Theorem](#)
- [The Electronuclear Reaction: Nuclear Reactions in TR-Zones \(1\)](#)
- [The Electronuclear Reaction: Nuclear Reactions in TR-Zones \(2\)](#)
- [The Electronuclear Reaction: Nuclear Reactions in TR-Zones \(3\)](#)
- [In Time-Reversed Zone: Nuclear reactions are biased](#)
- [In Time-Reversed Zone: Dynamics may be reversed](#)
- [Time-Reversed Zone: Significant only after time-density charging](#)
- [A New Conservation of Energy Law](#)
- [Questions on Carcinogens and EM](#)

Radiation

- [AS DEFINED, fields and potentials only exist in and on charged matter](#)
- [Serious flaws and errors in classical EM theory](#)
- [Electrodynamicists' Reaction to Removal of the Material Ether](#)
- [Reaction to Loss of Material Ether \(1\)](#)
- [Reaction to Loss of Material Ether \(2\)](#)
- [Some EM waves in Spacetime](#)
- [Air Medium Disturbances Generated When Air is Perturbed by a Plucked Taut String](#)
- [Launching a Spacetime Perturbation \("EM Wave"\) from a Wire Antenna](#)
- [Phase Conjugate Wavepairs Produce New Waves](#)
- [Scalar Electromagnetics \(Energetics\) View of EM \(1\)](#)
- [Scalar Electromagnetics \(Energetics\) View of EM \(2\)](#)

- [Scalar Electromagnetics \(Energetics\) View of EM \(3\)](#)
- [Extended General Relativity Principle](#)
- [Extension of Work-Energy Theorem](#)
- [34 Flaws in Classical EM Theory](#)
- [MTW's General Relativity Principle](#)
- [Use of General Relativity \(GR\) in Particle Physics](#)
- [The Missing Infolded Electrodynamics](#)
- [Infolded Engines: Excluded by Present Physics](#)
- [Mass curves and structures spacetime; ST curvatures change and structure mass \(1\)](#)
- [Mass curves and structures spacetime; ST curvatures change and structure mass \(2\)](#)
- [At Infinite Velocity: Each Is Every Other](#)
- [Lisitsyn's Report: Brain Code Broken](#)
- [Lorentz Closed Surface Integration](#)
- [Lorentz's physically insignificant energy flow can be collected and utilized](#)
- [Lorentz surface integration of the Poynting vector around a closed surface](#)
- [Longitudinal EM Photon Interaction with Charge](#)
- [Mass is transparent to longitudinal EM waves, which move through the infolded interiors of internal waves, potentials, and fields in the mass.](#)


- [Pumping with Longitudinal EM waves](#)
[Time-reverses the Mass](#)
- [Maxwell's Quaternion Theory](#)
- [Maxwellian Systems before and after](#)
[Regauging](#)
- [Maxwell's four equations reduce to: Page 1](#)
- [Maxwell's four equations reduce to: Page 2](#)
- More to come

Major Points of Presentation

- **Electrodynamics is seriously flawed**
 - Some major flaws
 - Corrections Indicated
- **Why past corrective attempts failed**
 - Force field concept is material
 - Missing two infolded transforms (M->ST; ST->M)
- **Missing Infolded general relativity inside electrostatics**
- **Several Kinds of EM Waves in Space**
 - How to make
 - Transduction
- **A French medical example as a deep experimental demonstration**
 - Spectacular regenerative cures funded by French Govt
 - Not comprehended, therefore suppressed
- **Cold Fusion examples of time-density wave effects**
 - China Lake Instrumental anomalies in electrolysis
 - Explanation of anomalous behavior of instruments
 - Explanation of new nuclear reactions
 - Wave energy transduction provides excess heat
- **Urge theorists to consider dramatically new reactions**

Chung's Carbon Filament Negative Resistor

UNIVERSITY AT BUFFALO, NY



© 1997 T. E. Bearden



Aharonov-Bohm Effect



- **In field-free regions**

- **E-field is zero**
- **B-field is zero**

© T.E. BEARDEN 1990



- **Potentials still exist, just no gradients**

- **Cause real effects**
- **Interference is the key**
- **Contain sum-zeroed substructures**



- **Does not follow from**


- **Mechanics**
- **Classical electromagnetics**




- **Required by quantum mechanics**

Absorption and emission reactions





ELECTRON ABSORBS PHOTON ENERGY, MOVES TO HIGHER ENERGY LEVEL. DECAYS BY EMITTING PHOTON



ELECTRON ABSORBS TIME DENSITY, MOVES TO DIFFERENT TIME DENSITY LEVEL. DECAYS BY EMITTING LONGITUDINAL OR PSEUDOLONGITUDINAL PHOTON



© T.E. BEARDEN 1998



Absorption and emission reactions

TRANSVERSE PHOTON

ELECTRON ABSORBS PHOTON ENERGY, MOVES TO HIGHER ENERGY LEVEL. DECAYS BY EMITTING PHOTON, REMOVING Δt




Figure A. Excited energy density level in one jump.

STREAM OF TIME-POLARIZED PHOTONS

ELECTRON ABSORBS TIME DENSITY, MOVES TO $1/\sqrt{2}$ DIFFERENT TIME DENSITY LEVEL. DECAYS BY EMITTING LONGITUDINAL OR PSEUDOLONGITUDINAL PHOTONS





Figure B. Excited time density level significant only after enormous number of increases.

© T.E. BEARDEN 1998


ANTI-ENGINE FOR CELL'S DEVIATION REVERSES CELL BACK TO NORMAL



© 1994, 1995 T.E. BEARDEN


By pumping the internal EM channel, process is dramatically accelerated.
This is the body's normal recovery mechanism, within its limits.

Asymmetric Regauging Produces Excess Force, Which Can be Used to Do Work on the System



Net Symmetric Regauging Does No Excess Work on the System


This little bird put one foot on wire A, and then the second foot on wire A also. He net symmetrically regauged, so he had no net excess force across him, even though his potential energy was increased. He asymmetrically regauged each foot, but both simultaneously so that one countered the other. Wise little bird!



Net Asymmetric Regauging Can Do Excess Work on the System


This little bird put one foot on wire A, and then the second foot on wire B. He net asymmetrically regauged, so he had lots of net excess force across him. That force then violently translated his little body parts every which way, doing lots of work in and on him. Bummer!

© T.E. BEARDEN 1999



Net Symmetric Regauging Does No Excess Work on the System

This little bird put one foot on wire A, and then the second foot on wire A also. He net symmetrically regauged, so he had no net excess force across him, even though his potential energy was increased. He asymmetrically regauged each foot, but both simultaneously so that one countered the other. Wise little bird!



Net Asymmetric Regauging Can Do Excess Work on the System

This little bird put one foot on wire A, and then the second foot on wire B. He net asymmetrically regauged, so he had lots of net excess force across him. That force then violently translated his little body parts every which way, doing lots of work in and on him. Bummer!


© T.E. BEARDEN 1996

Figure 32. Asymmetric regauging produces excess force, which can be used to do work on the regauging system.

Selected clipart © by Lotus SmartPics™

Becker's bone fracture healing experiments

- Tiny DC currents (picoamperes)
- Pulsed DC current can be utilized
- Pulsed magnetic fields may be utilized



© 1993, 1995 T.E. BEARDEN

RED BLOOD CELL
DEDIFFERENTIATES
Shucks hemoglobin coat
Grows nucleus

NEW CELL
REDIFFERENTIATES
Turns into type of cell that
makes cartilage

NEW CELL
REDIFFERENTIATES
Turns into type of cell that
makes bone
Deposits in fracture site,
healing the fracture

Becker's theoretical control system governing regeneration*

Trauma Producing Tissue Loss

Local cellular effect

Local nerve effect

CNS effect

Alterations in hormone pattern (prolactin)

Alterations in local DC field pattern

Local electrical effects

Local and systemic pools of target cells

Dedifferentiation into primitive mesenchymal cells

Phase I

Blastema formation

Phase II


growth

Redifferentiation into required cell types

Restoration of body part or tissue

*Becker & Spadaro, "Electrical stimulation of partial limb regeneration in mammals," Bull. N. Y. Acad. Med., 48(4), May 1972, p. 629.


Becker's theoretical DC control system involved with response to injury*



*Robert O. Becker, "The significance of bioelectrical potentials," *Bioelectrochemistry and Bioenergetics*, Vol. 1, 1974, p. 191.

Geometrical Distortion of Poynting Energy Flow

At or near resonance frequency, in the case shown the energy collection fraction (reaction cross section) increases dramatically.



a. Around an aluminum sphere at light energy 8.8 eV. Absorption efficiency = 18.00.

b. Around an aluminum sphere at light energy 5eV. Absorption efficiency = 0.1.

Figures per Craig F. Bohren, "How can a particle absorb more than the light incident upon it?", American Journal of Physics, 51(4), Apr. 1983, p. 326.



Cosmic Bootstrap: Before the Big Bang:




- *Cosmic repulsion behaved like a fluid with negative pressure*
- *The "universe" inflated (expanded) as a "false vacuum"*
- *As the negative-pressure fluid expanded, its energy went up rather than down*
- *When inflation stopped, the false vacuum decayed from its excited state*
- *Its excitation energy was released in a single great burst*

Paul Davies, Superforce, 1984, p. 194

BRIDGING ENABLES COP > 1.0

S-flow, $d\phi/dt$ flow, and EMF flow



Patent Pending

Current loop B is asymmetrically regauged by S, $d\phi/dt$, and EMF flow from current loop A.

© 1995 T.E. BEARDEN

Aspects of Strong Local Asymmetry

■ If local asymmetry is strong, conservation laws may be appreciably violated

- Energy
- Charge
- Spin
- Momentum
- Angular momentum

© T. E. Bearden 1995, 1996

■ Properties of an object may differ appreciably for

- Different observers
- Different detecting means
- One time to another
- One position to another

■ STRUCTURING THE INFOLDED EM INSIDE POTENTIALS, FIELDS, AND WAVES

- Strongly breaks local symmetry
- Provides spacetime engines giving the above effects

Other Aspects of Strong Local Asymmetry

- Local spacetime is curved
- Lorentz invariance of vacuum is violated
- May be a local "sink" or "source"
- Gravitational-inertial effects from EM
- Translation between virtual and observable
- Electrogravitational solitons
- Action at a distance
- Transmutation effects may exist
- Scalar/pseudoscalar field translation
- **COMMENT:** Thus the use of infolded longitudinal EM fields and waves to strongly break local symmetry allows interested clustering of spacetime curvatures. These spacetime structures are vacuum engines, or *spacetime engines*.

© T. E. Bearden 1995, 1998

Bunge on the Status of Electrodynamics

"... it is not usually acknowledged that electro-dynamics, both classical and quantal, are in a sad state..."

"... the best modern physicist is the one who acknowledges that neither classical nor quantum physics are cut and dried, both being full of holes and in need of a vigorous overhauling..."

*Mario Bunge, Foundations of Physics, Springer-Verlag, New York, NY, 1967, p. 176..

© 1996 T.E. BEARDEN

Bunge on the Status of Electrodynamics and Physics

"... it is not usually acknowledged that electro-dynamics, both classical and quantal, are in a sad state..."

"... neither classical nor quantum physics are cut and dried, both being full of holes and in need of a vigorous overhauling..."

© 1996 T. E. BEARDEN

*Mario Bunge, Foundations of Physics, Springer-Verlag, New York, NY, 1967, p. 176..

Bunge on the Status of Classical and Quantum Physics

"... it is not usually acknowledged that electrodynamics, both classical and quantal, are in a sad state... the best modern physicist is the one who acknowledges that neither classical nor quantum physics are cut and dried, both being full of holes and in need of a vigorous overhauling not only to better cover their own domains but also to join smoothly so as to produce a coherent picture of the various levels of physical reality."

***Mario Bunge, Foundations of Physics, Springer-Verlag, New York, NY, 1967, p. 176..**

Fraction of Casualties Surviving

N = 1 - 40 million

FRACTION SURVIVING



CASUALTIES

TYPE TREATMENT RECEIVED

EXPRESSED IN FRACTION SURVIVING

- With conventional treatment
- With 1st generation P-treatment
- With 2nd generation P-treatment
- With 3rd generation P-treatment

© 1998 T. E. BEARDEN

FIRST STEP IN SELF-PROMOTION OF CANCER DUE TO HYPOXIA



SINGLE CELL ANAEROBIC



SINGLE CELL AEROBIC*



MULTICELL AEROBIC* (INCLUDES MAN)



OXYGEN-RICH

CENTRAL CONTROL
TIME-FORWARD PATH AS IT DEVELOPED

CANCER CELL



PARTIALLY ANAEROBIC




SINGLE CELL AEROBIC

TIME-REVERSED PATH FOR SUSTAINED OXYGEN DEFICIENCY


FIRST STEP IN OXYGEN-DEFICIENCY DE DIFFERENTIATION

© 1993 T. E. Bearden

Differentiation and Dedifferentiation Timelines



Evolution of Cells on Earth



© 1999 T. E. BEARDEN


A Charged Particle q Is a Coupled System of m and ϕ

(A proposed formal definition of a charged mass)

Note: X is a coupling operator

$$q \equiv f_q \times m_q$$

Also note that the ϕ portion of the charged particle can be internally structured, per Stoney-Whittaker-Ziolkowski, with hidden bidirectional EM wavepairs (their sums and products). Note that this constitutes an EM application of Bohm-type hidden variable theory, with many ramifications.



"...curiously enough, we do not know exactly what charge is, only what it does. Or, equally significantly, what it does not do."

M.P. Silverman, *And Yet It Moves: Strange Systems and Subtle Questions in Physics*, Cambridge University Press, 1993, p. 127.

© T.E. BEARDEN 1997

An electric charge Q consists of a massless flux component and a mass component, coupled together (interacting). Q is a broken symmetry. The mass of the charge continually and violently exchanges virtual photons with the surrounding vacuum.

Thus Q is energetically driven by the surrounding vacuum potential, which itself is to first order just a violent virtual photon flux.

The Ubiquitous Assumption: Two Asymmetrical Regaugings for Net Symmetry



**You really need
a Maxwell's demon,
to do some free work.**



**Your electrodynamicist
says he can make one,
by regauging.**

© T.E. BEARDEN 1999

The Ubiquitous Assumption: Two Asymmetrical Regaugings for Net Symmetry



He says he will do the regauging for you. Hey, what a swell guy!



He regauges twice, and makes two Maxwell's demons equal and opposite. They won't do any net work for you!

© T.E. BEARDEN 1990

The Ubiquitous Assumption: Two Asymmetrical Regaugings for Net Symmetry



He disdainfully says,
"We always do it that way!
We must enforce local
equilibrium and the second
law of thermodynamics!"



You decide to have a
little chat with your
electrodynamicist!
With friends like that,
who needs enemies!


© T.E. BEARDEN 1999

A Charged Particle Is a Coupled System

(The first formal definition of charge)

$$q \equiv \phi_q \times m_q$$

VACUUM
CHARGE



Note: X is a coupling operator

virtual
photon
flux
exchange

ϕ_q

© 1994 T. E. BEARDEN

"...curiously enough, we do not know exactly what charge is, only what it does. Or, equally significantly, what it does not do."


MP. Silverman, *And Yet It Moves: Strange Systems and Subtle Questions in Physics*, Cambridge University Press, 1993, p. 127.

An electric charge q is a broken symmetry in its fierce energy exchange with the active vacuum. It coherently integrates some of the virtual energy absorbs and outputs it as Poynting energy flow.


Chung's Carbon Filament Negative Resistor

UNIVERSITY AT BUFFALO, NY


© 1990 T.E. Bearden



Typical circuit has about 10-13 energy collection efficiency



Distance traveled by S in one hour is 1.08×10^{12} meters.



Example: 1.3 amps flowing in DC circuit, 1.8 mm diameter copper wire. $J = 51 \text{ A/cm}^2$

$d_i = 0.1368$ meters

S violently transports ϕ provided from the source.

$j\phi$ sluggishly transports the energy collected and dissipated in the circuit.

For the case discussed, the electron drift velocity in the circuit is about 3.8×10^{-5} meters/sec. So $J\phi$ moves about 0.1368 meters in one hour. Thus $J\phi$ has collected about 0.1368 meters of the ϕ -filled S-tube. During that same hour, the S-flow evoked by the power source will have traveled 1.08×10^{12} meters. The ϕ of both currents is the same. Both are involved in the same energy-filled tube. Thus S has provided and transported about 7.89×10^{12} times as much energy along the circuit in one hour as the $j\phi$ has been able to collect, transport, and dissipate as work in the circuit. Thus this circuit provides about 10^{13} collection efficiency.




Classical View of EM

- Potentials just mathematical conveniences
 - Not real
 - No internal structure, just magnitude
- Force fields primary, causative, can exist in space in absence of observable mass
 - All EM actions due to force fields, none when zero
 - No action at a distance
- Vacuum empty; without energy or structure
 - No locally curved spacetime
 - No gravity effects
 - No quantum effects
- Maxwell's equations complete
 - Symmetrically regauged *arbitrarily*
 - Topology dramatically reduced



Curved Spacetime acts as source or sink

- **May emit excess radiation and energy (e.g., heat)**
- **May absorb radiation and energy (e.g., cool)**
- **May transduce waves
TW \Leftrightarrow LW \Leftrightarrow TDW**
- **New energy conservation law applies**



© 1998 T.E. BEARDEN



CHRONIC DISEASE PUZZLING

- **SCIENCE REALLY DOES NOT UNDERSTAND THE MECHANISMS OF A SINGLE CHRONIC DISEASE, INCLUDING CANCER**
- **MANY TROUBLING FACTORS EXIST**
 - **MIND**
 - **STRESS**
 - **SPONTANEOUS REMISSION**
 - **PLACEBO EFFECTS**
 - **COMBINATIONS**
- **GENERATION OF THE PLACEBO EFFECT BY THE CELLULAR REGENERATION SYSTEM IS UNRECOGNIZED**



© 1994, 1995 T. E. Beaman

Cancer Characteristics

- Not one disease, but a whole range
- Starts in damaged aerobic cell
- Cell shakes off body's central control
 - Starts uncontrolled division
 - Becomes a lump
- Can send forth cancer cells
 - Through blood
 - Through lymph
- Form metastases (secondary tumors)
- Often become anaerobic

© 1993 T.E. Beaufee

Solving a Cancer Enigma

- **An enigma has been failure of immune system to attack some tumor cells**
 - **Suppressor cells may curb immune system?**
 - **Tumor may lack antigens normally identified by the immune system?**
- ***Promotion of damaged cell to cancer is usually ordered by cellular control system***
 - ***Regenerative and recovery system forces cell back toward anaerobe***
 - ***First step breaks free from central growth***
 - ***Control continues to recognize cell as self***


© 1993 T.E. Huxley

CANCER: CHARACTERISTICS

- **NOT ONE DISEASE BUT A WHOLE RANGE**
 - **STARTS IN ORDINARY (AEROBIC) CELL**
 - **CELL SHAKES OFF BODY'S DICTATORIAL CONTROL***
 - **STARTS UNCONTROLLED DIVISION**
 - **BECOMES A LUMP**
 - **CAN SEND FORTH CANCER CELLS**
 - **THROUGH BLOOD**
 - **THROUGH LYMPH**
 - **THESE FORM METASTASES (SECONDARY TUMORS)**
 - **OFTEN BECOME ANAEROBIC (NON-OXYGEN USING)**
 - **ENIGMA IS FAILURE OF IMMUNE SYSTEM TO ATTACK SOME TUMOR CELLS**
 - **SUPPRESSOR CELLS MAY CURB IMMUNE SYSTEM**
 - **TUMOR MAY LACK ANTIGENS NORMALLY IDENTIFIED BY THE IMMUNE SYSTEM**
- *R&R system forces cell back toward anaerobe*

© 1993 T.E. Bearden

Nonlinear Optics Distortion Correction Theorem



"If a scalar wave $E_1(r)$ propagates from left to right through an arbitrary but lossless dielectric medium, and if we generate in some region of space [say near $z = 0$] its phase conjugate replica $E_2(r)$, then E_2 will propagate backward from right to left through the dielectric medium, remaining everywhere the phase conjugate of E_1 ."

The Electronuclear Reaction: Nuclear Reactions in TR-Zones (1)

- **Nucleon interchange**
 - Changes neutron into proton and vice versa
 - At high rate (flat spacetime assumed)
- **Statistics become skewed due to curved ST**
- **Can bias statistics toward either neutron or proton end**
 - Biases toward transmutation of elements along isomer chains
 - Concept of "isomer" is vastly expanded

© 1997 E. Scuderi

The Electronuclear Reaction: Nuclear Reactions in TR-Zones (2)

- **Normal dynamics (impulses) may reverse**
 - *Like charges attract, unlike charges repel*
 - *Positive charges cluster (are drawn together)*
 - *H⁺ ions (protons) may form quasi-nuclei*
 - *Nucleus coulomb barrier becomes assistant*
- **Many "impossible" reactions now possible**
 - *Quark access and quark flipping by EM*
 - *Ion clustering*
 - *Positive charges as phase conjugate mirrors*
 - *Time-density waves and time-energy charging*

© 2011 E. Beaudin

The Electronuclear Reaction: Nuclear Reactions in TR-Zones (3)


- ***Not observed until time-energy charging has become substantial***
 - *Time delay (hours, days)*
 - *Transduction of LW --> TW involved*
 - *Specific time-charging history of detectors is involved in whether they detect or not*
 - *Longitudinal EM waves become significant*
- ***Curved spacetime can act as source or sink***
 - *May emit excess radiation*
 - *May absorb excess radiation*

© 1997 T. E. Darden

In Time-Reversed Zone: Nuclear reactions are biased


- **In normal flat ST, $n \Leftrightarrow p$ at high rate**
- **Curved ST skews statistics**
- **Biases to either n or p**
- **Biases to transmutation along isomer chain**
- **"Isomer" concept greatly expanded**

© 1998 T.E. BEARDEN




In Time-Reversed Zone: Dynamics may be reversed

- Like charges may attract and unlike charges repel
- Coulomb barrier becomes Coulomb assistant
- Clusters like charges and forms artificial nuclei
- "Impossible" nuclear reactions at low energy
- Time-density interactions



Time-Reversed Zone: ***Significant only after time-density charging***

- **Hours, for "fast" onset**
- **Days, for "slower" onset**
- **Transduction involved for
TW \Leftrightarrow LW \Leftrightarrow TDW**
- **Time-history of each
detector is significant**
- **Longitudinal EM waves
become significant**



A New Conservation of Energy Law

- Physics conserves total "energy and mass-energy"
- Now also must consider "time-energy"
- Conserve total mass-time-energy (spatiotemporal energy)
- Let E_T = total energy, E_M = mass energy, E_E = ordinary energy, and E_t = time energy. Then

$$E_T = E_E + E_M + E_t$$

© 1990 T.E. Bearden

$$(kE_t \Rightarrow E_E > 0) \Rightarrow E_T > (E_M + E_E)$$

- If some E_t is transduced into E_E , the experiment will violate the old spatial mass-energy conservation law



QUESTIONS ON CARCINOGENS AND EM RADIATION



■ UNRESOLVED QUESTIONS ON EM RADIATION

- DOSAGES
- ACTIVE MECHANISMS
- REPLICATION DIFFICULTIES
- MEASUREMENT DIFFICULTIES
- WHERE, WHEN, HOW ILL EFFECTS OCCUR

■ SIMILAR QUESTIONS EXIST ABOUT EVERY CARCINOGEN


■ SCIENCE REALLY DOES NOT UNDERSTAND THE MECHANISMS OF A SINGLE CHRONIC DISEASE, INCLUDING CANCER

- FACTORS SUCH AS MIND, STRESS, SPONTANEOUS REMISSION, PLACEBO EFFECTS, AND COMBINATIONS OF FACTORS ARE PARTICULARLY TROUBLESOME
- GENERATION OF THE PLACEBO EFFECT BY THE REGENERATION & RECOVERY SYSTEM IS UNRECOGNIZED




© 1994, 1995 T.E. Bearden

As defined, fields and potentials only exist in and on charged matter



Serious flaws and errors in classical EM theory.


1. Eliminates the Internal EM Inside the Scalar Potential.
 2. No Definition of Electrical Charge or of Scalar Potential.
 3. Equations Still Assume Material Ether Per Maxwell (Unchanged).
 4. Use of Force Fields in Vacuum is False (and Known to be So).
 5. Treats Charge q as Unitary Instead of Coupled System $q = \phi[q]m(q)$.
 6. Confuses Massless Potential Gradients as Forces (See #3, #4).
 7. Does Not Utilize Mass as a Component of Force (See #23).
-
8. Erroneously Assumes EM Force Fields as Primary Causes.
 9. Topology of EM Model Has Been Substantially Reduced.
 10. Does Not Include Quantum Potential or Action at a Distance.
 11. Does Not Include Superluminal Velocity of Inner EM Components.
 12. Does Not Utilize Extended Near-Field Coulomb Gauge Effects.
 13. Does Not Include EM Generatrix Mechanism for Time Flow.
 14. Does Not Unify Photon and Wave Aspects (Requires 7-D Model).
-
15. Does Not Include Electron Spin and Precession (See #19, #24).
 16. Treats EM Energy As Existing In "Chunks," Instead of as Flow.
 17. Confuses Energy and Energy Collection (See #16).
 18. Discards Half of Every EM Wave in Vacuum (See #22).
 19. Erroneously Uses Transverse Vacuum Wave; It's Longitudinal.
 20. Arbitrarily Regauges Maxwell's Equations to Eliminate Overunity.
 21. Omits Phase Conjugate Optics Effects (The Rule in Internal EM).
 22. Does Not Include EM Cause of Newtonian Reaction Force.
 23. Erroneously Assumes Separate Force Acting on Separate Mass.
 24. Confuses Detected Electron Precession Waves as Proving Transverse EM Waves in Vacuum (Remnant of Old "EM Fluid" Concept).
 25. Due to Error in String Wave, Omits the Ubiquitous Antiwave.
 26. Assumes Equilibrium; Not True Unless Include Vacuum Interactions.
 27. Higher Topology Required, to Model Electromagnetic Reality.
 28. Lorentz surface integration discards Poynting energy transport.
 29. Has nothing at all to say about form of EM entities in massless space.
 30. Eliminates the infolded general relativity using EM-force as curve agent.
 31. Does not include longitudinal EM waves as time domain oscillations.
 32. Does not include EM mechanism that generates time flow and flow rate.



REACTION TO LOSS OF MATERIAL ETHER

**Material Ether
Assumed in
Equations**


**Potentials,
Fields, and
Waves
Assumed
Material**




**Not a single
equation was
changed!**

© T.E. EBARDEN 1999


ELECTRODYNAMICISTS



Some EM Waves in Spacetime



Air Medium Disturbances Generated When Air is Perturbed by a Plucked Taut String




The string wave and the holder wave never leave the string and holder respectively.

Wave is alternatively compressive and rarefactive

String slaps air medium

WAVE IN STRING (TRANSVERSE)
Lower degree of freedom for string.
Can only vibrate mostly laterally.


Drude
Electron
Gas waves



ANTI WAVE IN
HOLDER
(Highly damped; not shown to scale)

Nuclear
proton
waves

Launching a Spacetime Perturbation ("EM Wave") from a Wire Antenna




© T.E. BEARDEN 1994, 1995

Phase Conjugate Wavepairs Produce New Waves



**TRANSVERSE EM WAVE PLUS
PHASE CONJUGATE REPLICAWAVE**

LONGITUDINAL EM SURGE WAVE



TIME DENSITY WAVE

*Oscillates rate of flow of time
about some average value*

**LONGITUDINAL EM WAVE PLUS
PHASE CONJUGATE REPLICAWAVE**

© T.E.BEARDEN 1994 1995



Scalar Electromagnetics (Energetics) View of EM

- Potentials real; primary causes of EM phenomena
 - Force fields made by differentiations of potentials
 - Force fields are effects, not primary causes
 - Force fields exist only in, on mass particles
- Actions due to potentials and their interference
 - Action at a distance or locally
 - Quantum potential and hidden variables
- Locally curved spacetime, engineerable by EM
 - Gravity effects not necessarily negligible
 - Strong EM force or time used as agent of curvature

© T. B. BEARDEN 1996, 1998



Scalar Electromagnetics (Energetics) View of EM (Cont'd)

- Potentials have internal structure
 - Stoney/Whittaker structure
 - Longitudinal EM phase conjugate wavepairs
 - Each wavepair is a time-polarized EM wave
- Vacuum EM is a potential and has microstructure
 - Stoney-Whittaker biwaves/t-polarized waves
 - Fluctuations exhibit chaos
- Spacetime \equiv Vacuum \equiv Potential \equiv Flux

© T. E. BARDEN 1994, 1999




Scalar Electromagnetics (Energetics) View of EM (Cont'd)

- Statistics may have hidden order (already chaotic)
- Uses spacetime curvature engines to alter matter (inside-out or outside-in)
- Engineerable EM mechanism generates rate of flow of a mass through time
- Quantum potential with specific QP can be used for instant action at a distance

Extended General Relativity Principle

- "All levels of energy structures and all levels of time structures mold spacetime geometry.
- This is called a vacuum engine (spacetime engine).
- Spatial and temporal structuring act upon mass at all levels.
- This produces templates of forces, for translations and stresses."

© T.E. BEARDEN 1999



Extension of Work-Energy Theorem

© T. E. BEARDEN 1997

- Theorem: $W = k_f - k_i = \Delta k$ [1]

- k must be interpreted not as energy per se, but as *collected/collecting* energy.

- The reaction cross section λ for the collecting process must be included.

- Extension: $W = \lambda (k_f - k_i) = \lambda \Delta k$ [2]

- Normally $\lambda \leq 1$, as for elastic collision or Stokes emission.


- However, $\lambda > 1$ is now possible, for processes which asymmetrically self-regauge. An example is Letokhov's negative absorption of the medium.

- Working models are the Patterson Power Cell[®] and Lawandy lasing without population inversion.

MTW's General Relativity Principle

- "Space acts on matter, telling it how to move."
- In turn, matter reacts back on space, telling it how to curve."

© T.E. BEARDEN 1998



Use of General Relativity (GR) in Particle Physics

- GR seldom used in particle physics
- In cold fusion, Matsumoto* has applied general relativity
- Consistent with important cold fusion results
- Used spacetime (ST) curvature by energy density
- Matsumoto did not utilize:
 - Longitudinal EM waves
 - Time density waves
 - Time density curvatures of ST (gain = 9×10^{16})

© 1998 T. E. Bearden

* T. Matsumoto, "Mechanisms of Electro-Nuclear Collapse,"
Proc. ICCF-7, Vancouver, BC, Canada, Apr 1998, p. 98



THE MISSING INFOLDED ELECTRODYNAMICS

Maxwellian electrodynamics has nothing at all to say about EM entities in space or their form

**Material
Electric
Fluid
Disturbed**

**Material
Electric
Fluid
Disturbed**



MATTER-TO-MATTER TRANSFORM

**INFOLDED INSIDE
MAXWELLIAN EM
AS HIDDEN EM
VARIABLES**

**MATTER-TO-SPACETIME
TRANSFORMATION**

**SPACETIME-TO-MATTER
TRANSFORMATION**

**INTERESTED SPACETIME CURVATURES
(VACUUM ENGINES)
INFOLDED GENERAL RELATIVITY**

**EINSTEIN'S
SPACETIME**


© T.E. BEARDEN 1998

Infolded Engines: Excluded by Present Physics

- ***The infolded dynamics of EM systems are not considered in general relativity, electromagnetics, or quantum mechanics***
- ***None of these disciplines include such spacetime curvature engines (vacuum engines)***
- ***Thus GR, EM, and QM cannot be unified, since ST curvature engines are where the unification mainly occurs***


© 1999 T. E. BEARDEN

ST curvatures propagate



The *mass-to-mass transform* contains two hidden infolded transforms: (i) the *mass-to-spacetime transform* and (ii) the *spacetime-to-mass transform*. What is transmitted and propagates in space is a *spacetime perturbation*.

Mass curves and structures spacetime; ST curvatures change and structure mass




At Infinite Velocity:

Each Is Every Other



SPACETIME



**VIRTUAL
PARTICLE
FLUX**



POTENTIAL



**MIND AND
MIND LEVELS**



MASS


PLANET



LIVING ORGANISMS

© 1997 T.E. Bearden

Lisitsyn's Report: Brain Code Broken



- ★ **THEORY DEVELOPED AND FITTED**
- ★ **HYSTERESIS MEMORY LOOP**
- ★ **CONTROLLED EM INDUCTION**
 - **IMAGES**
 - **SENSATIONS**
 - **PREDETERMINED EMERGENCE**
- ★ **23 EEG BANDS**
 - **UP TO 8.1×10^{20} HZ**
 - **11 INDEPENDENT CHANNELS**


© 1970, 1985, 1991 T.E. BEARDEN

S.K. Lisitsyn, "New Approach to the Analysis of Electroencephalograms,"
DDC Report AD73005, p. 16-25.

Lorentz Closed Surface Integration

Selects only the locally divergent portion of the Poynting energy flow that powers the component.

POYNTING GENERATOR




S.S. LORENTZ

**Only the wind on my ship
is of physical significance!**

© 1997 T.E. Bearden

Lorentz Closed Surface Integration


Selects only the locally divergent portion of the Poynting energy flow that powers the component.



© T. E. BEARDEN 1998

Lorentz's physically insignificant energy flow can be collected and utilized.

Lorentz surface integration of the Poynting vector around a closed surface




1a. Lorentz surface integration.

1b. Actual S in and S out.

Note: If the S -vector is integrated over the closed surface, then all energy *transport* passage is zeroed, leaving only the very small component of the Input S -flow that is powering the joule heating of the resistor. In short, only the small component of the S -flow that is equal in magnitude to the Slepian vector $j\phi$ remains. This measures only the tiny portion of the S -flow that is "collecting" on electrons passing through the resistor, and therefrom being dissipated out of the resistor as joule heating. It discards everything else (all S_{TR} 's and S_{OUT}). S_{IN} becomes S_R .

Longitudinal EM Photon Interaction with Charge


- Creates time-density wave pumping
- Slowly creates time-density charging




- Fundamental units are arbitrary in physics
- Can express all physics in one unit -- e.g., time
- Time is highly compressed energy
- 1 sec = 9×10^{16} joules of transverse EM wave energy

Figure 1-8. Infolded biwave composition of a scalar potential.

Mass is transparent to longitudinal EM waves, which move through the infolded interiors of internal waves, potentials, and fields in the mass



Pumping with Longitudinal EM Waves Time-reverses the Mass





Maxwell's Quaternion Theory



- **Quaternions:**

- Discovered by Hamilton in 1843
- A quaternion has a vector part
- A quaternion has a scalar part
- First significant non-arithmetic number system
- Higher topology than vector or tensor algebra



- **Maxwell's theory was in quaternion equations**

- 20 equations in 20 unknowns
- Reduced to a small 4-equation subset by Heaviside and Gibbs
- EM topology dramatically reduced




© T.E. BEARDEN 1995, 1996



Maxwellian Systems

Before regauging,
after symmetrical regauging,
and after asymmetrical regauging

Note: Any system must first break equilibrium in its vacuum exchange, before it can energize Drude electrons and "power" the load. Asymmetric regauging is always utilized. Powering loads and losses is a symmetry-restoring operation.



© 1996, 1997 T.E. BEARDEN

Selected clipart © by Lotus SmartPics™

Maxwell's four equations reduce to:

$$\nabla^2 \Phi + \frac{1}{c} \frac{\partial}{\partial t} (\nabla \cdot A) = -4 \pi \rho \quad [1]$$

$$\nabla^2 A - \frac{1}{c^2} \frac{\partial^2 A}{\partial t^2} - \nabla (\nabla \cdot A + \frac{1}{c} \frac{\partial \Phi}{\partial t}) = -\frac{4 \pi}{c} J \quad [2]$$

A is replaced by A', where

$$A' = A + \nabla \Lambda \quad [3]$$

The new B' field then becomes

$$B' = \nabla \times (A + \nabla \Lambda) = \nabla \times A + 0 = \nabla \times A = B \quad [4]$$

A new E-field will also be formed. So let

$$\Phi' = \Phi - \frac{1}{c} \frac{\partial \Lambda}{\partial t} \quad [5]$$

Per Jackson, (A, Φ) are habitually chosen so that

$$\nabla \bullet A + \frac{1}{c} \frac{\partial \Phi}{\partial t} = 0 \quad [6]$$

The *net symmetrical* regauging separates variables. Two inhomogeneous wave equations result:

$$\nabla^2 \Phi - \frac{1}{c^2} \frac{\partial^2 \Phi}{\partial t^2} = -4\pi\rho \quad [7]$$

$$\nabla^2 A - \frac{1}{c^2} \frac{\partial^2 A}{\partial t^2} = -\frac{4\pi}{c} J \quad [8]$$

© T.E. BEARDEN 1997

Equations [1] and [2] arbitrarily changed to [7] and [8]