

## NATURAL PEST & DISEASE CONTROL

Henry Elwell & Anita Maas

Our quest for 'instant' higher yields to cater for a growing, hungry population has led us to adopt short-term crisis management in the control of pests and diseases on our farms. Nature became the enemy and we used destructive 'chemical warfare'.

This book aims to return us to a new co-operation with nature by offering a number of tried and tested methods to establish a long-term healthy balance within the farm environment.

It is essential reading for all farmers and gardeners in the southern African region - small-scale, large-scale, commercial and subsistence farmers alike. Not only will you find new ways of dealing with recurring problems but you will find a whole new philosophy to guide you in coping with any challenges that may arise in the future.



NATURAL FARMING NETWORK



Cover design: Rosemary Pickard

# NATURAL PEST & DISEASE CONTROL

Henry Elwell  
&  
Anita Maas



## NOTE FROM THE AUTHORS

Over the decades much valuable information on this important subject has been lost simply for lack of systematic compilation of the experiences of practising farmers and gardeners. We hope that research and publications like this one will prevent this waste in future. This is an ongoing task and we would welcome any information or suggestions that readers may have which will help to make future editions more comprehensive.

Users are also invited to record their own experiences in line with the format adopted in this book and to forward the information to the Natural Farming Network Zimbabwe, P.O. Box 8515, Causeway, Harare, Zimbabwe.

Please note that every effort has been made to make the translations of plant names into the vernaculars as accurate and useful as possible. Names have been omitted when there appeared to be any risk of confusion or any doubt. However, despite this, we realise that ambiguities due to local variations may still occur.

## ACKNOWLEDGEMENTS

The authors wish to acknowledge the following people and organisations:  
J. Bisset for checking the entomological data; R.B. Drummond for identifying common Zimbabwean plants; J. Galloway for checking the pathological data; R. Gunner for useful suggestions; S. Mavi for editing the botanical information; K. Murwira for locally appropriate contributions; S. Page (Dr) for information on nematodes and other data; S. Z. Sithole (Dr) for identifying major pests in Zimbabwe; J. Turner & J. Nzira for generous data on pest control; J. Wilson for general technical advice and his comments on the review draft; S. van den Akker for computer expertise; H. Vukasin for editorial and general comments and for securing funding for the book, without which we could not have gone to print; E. Zitsanza for entomological contributions; P. Zvoutete for pathological contributions; I. and O. Hedberg (Drs) for compiling the list of plant name translations; W. R. Mziray (Dr) from the Tropical Pesticides Research Institute in Tanzania; P. S.M. Phiri (Dr) from the University of Zambia and B. Hargreaves from the National Museums in Botswana for providing local translations of plant names; M. Beddingfield for editorial, design and layout work.

Experts from the following provided assistance, advice and expertise:  
Agricura (PVT) LTD; Department of AGRITEX; Fambidzanai Training Centre; Intermediate Technology Development Group (ITDG); Permaculture Association of Zimbabwe (PAZ); Plant Protection Research Institute (DRSS); National Herbarium (DRSS); Natural Farming Network; Silveira House; University of Zimbabwe; Zimbabwe Organic Producers Association (ZOPA).

Grateful thanks are also due to the many Zimbabweans who contributed their knowledge and experience.

Finally, the production of this book would not have been possible without the generous financial assistance given by the Interchurch Fund For International Development (ICFID), Toronto, Canada.

## THE PLANT PROTECTION IMPROVEMENT PROGRAMME

The Plant Protection Improvement Programme (PPIP) supports the development and use of appropriate and sustainable crop protection methods for small-scale farmers in Southern and East Africa. Activities include research, training and extension. PPIP is funded by the Swedish International Development Authority (SIDA).

## THE NATURAL FARMING NETWORK

The Natural Farming Network is a network of twelve development agencies co-operating to promote sustainable agriculture in Zimbabwe. The network provides training and technical assistance to staff of member groups and to farmers and groups of farmers identified by members. The network core programme is funded by W.K. Kellogg Foundation, USA and NOVIB, The Netherlands.

## Contents

Preface .....	5	Aphids .....	37
Foreword .....	6	Armyworm .....	39
<b>PART I : THE FOUNDATIONS OF PEST AND DISEASE PREVENTION</b>			
Introduction .....	9	Bean Stem Maggot or Bean Fly .....	42
The Natural Order .....	10	Beetles — Blister .....	43
Regaining control .....	11	Beetles — Leaf .....	44
Beetles — Snout .....	46	Cutworms .....	48
Four steps to re-establishing the Natural Order .....	14	Diamond Back Moth .....	50
1 Reinststate Soil Fertility .....	15	Fruit Fly .....	52
i Dig less .....	15	Grasshoppers .....	53
ii Mulch more .....	15	Maize Stalk Borer .....	54
iii Compost always .....	16	Nematodes .....	57
2 Provide Natural Habitats For Predators .....	17	Red Spider Mite .....	59
3 Re-introduce Plant Diversity .....	18	Scale Insects .....	61
i Rotations .....	19	Slugs & Snails .....	62
ii Mixed cropping .....	20	Stink or Shield Bugs .....	63
iii Intercropping .....	21	Termites .....	65
iv Strip cropping .....	22	Thrips .....	67
v Permaculture designs .....	23	Tip or Twig Wilts .....	69
4 Use Natural Remedies to Replace .....	24	Whitefly .....	70
Agro-chemicals .....	24	White Grub .....	71
Supportive Strategies .....	26	<b>Diseases</b>	
Hygiene .....	26	Bacteria .....	73
Select resistant varieties .....	26	Fungi .....	74
Timing of planting .....	27	Viruses .....	76
Plant spacing .....	27	Other Pests and Diseases .....	79
Insect barriers and decoys .....	28	<b>PART III : REMEDIES FOR PROTECTING FIELD CROPS</b>	
Encourage predators .....	28	<b>A Miscellaneous substances and Methods</b>	
Summary .....	30	..... 83	
<b>PART II: PESTS, DISEASES AND CONTROL MEASURES</b>			
<b>The Advantages of Natural Pest Control</b>			
Common pests .....	33	A1 Aromatic Plants .....	83
The correct approach .....	33	A2 Ash .....	83
How to use this section .....	34	A3 Baking or Washing Soda .....	84
American Bollworm .....	35	A4 Biological Controls .....	84
Ants .....	36	A5 Bordeaux mixture .....	85
		A6 Brushing .....	85
		A7 Builders' lime .....	85
		A8 Burgundy mixture .....	86
		A9 Burning .....	86

A10	Clay, lime & other powdery materials	86
A11	Coffee grounds & tea leaves	87
A12	Compost	87
A13	Glues	88
A14	Grease	88
A15	Hand picking	88
A16	Hens, bantams & ducks	89
A17	Manure, dung & urine	89
A18	Milk	90
A19	Mulch	90
A20	Noise & vibration	91
A21	Oil	91
A22	Potassium permanganate	92
A23	Salt preparations	92
A24	Smoke & soot	92
A25	Soap solutions	92
A26	Soil	93
A27	Sugar	93
A28	Sulphur	94
A29	Sun	94
A30	Vinegar	94
A31	Water	95
B	Plant Materials	96
B1	Useful Common Cultivated Plants	96
	Guidelines for the use of Botanical Sprays	96
B1.1	<i>Allium cepa</i> ; Onion	96
B1.2	<i>Allium sativum</i> ; Garlic	96
B1.3	<i>Camelia sinensis</i> ; Tea	97
B1.4	<i>Capiscum frutescense</i> & <i>annuum</i> ; Chilli, Sweet peppers	97
B1.5	<i>Carica papaya</i> ; Pawpaw	97
B1.6	<i>Chrysanthemum cinerariifolium</i> ; Pyrethrum	98
B1.7	<i>Crotalaria juncea</i> ; Sunnhemp	98
B1.8	<i>Eleusine coracane</i> ; Finger millet, Rapoko	98
B1.9	<i>Eucalyptus</i> spp; Gum tree	98
B1.10	<i>Glycine max</i> ; Soyabean	99
B1.11	<i>Impomoea batatas</i> ; Sweet potato	99
B1.12	<i>Lycopersicon esculentum</i> ; Tomato	99
B1.13	<i>Manihot esculenta</i> ; Cassava	100
B1.14	<i>Matricaria inximia</i> ; Feverfew	100
B1.15	<i>Melia azedarach</i> ; Syringa	100
B1.16	<i>Nerium indicum</i> ; Oleander	100
B1.17	<i>Nicotiana tabacum</i> ; Tobacco	101
B1.18	<i>Ocimum basilicum</i> ; Sweet basil	101
B1.19	<i>Rheum</i> spp; Rhubarb	101
B1.20	<i>Tagetes erecta</i> ; African marigold	102
B 2	Useful Wild Plants and Trees	102
B2.1	<i>Agave americana</i> ; American aloe & other spp.	102
B2.2	<i>Ageratum conyzoides</i> ; Goat weed	102
B2.3	<i>Annona muricata</i> , <i>reticulata</i> & <i>squamosa</i> ; Sugar apple	103
B2.4	<i>Annona senegalensis</i> ; Custard apple	103
B2.5	<i>Argemone mexicana</i> ; Mexican poppy	103
B2.6	<i>Balanites aegyptiaca</i> ; Desert date	104
B2.7	<i>Bersama abyssinica</i> ; Bitter bark	104
B2.8	<i>Bidens pilosa</i> ; Blackjack	104
B2.9	<i>Catunaregam spinosa</i>	104
B2.10	<i>Chenopodium ambrasioides</i> ; Wormwood	105
B2.11	<i>Datura stramonium</i> ; Thorn apple, Jimson weed	105
B2.12	<i>Euphorbia hirta</i> ; Asthma weed	105
B2.13	<i>Euphorbia tirucalli</i> ; Rubber hedge including spp called <i>Euphorbia</i> or <i>Cadelabra</i>	105
B2.14	<i>Jatropha curcas</i> ; Barbados nut	106
B2.15	<i>Lantana camara</i> ; Lantana	106
B2.16	<i>Leonotis nepetifolia</i> ; Lion's ear	106
B2.17	<i>Lippia javanica</i> ; Lippia	106
B2.18	<i>Lobelia colummaris</i> ; God's tobacco	107
B2.19	<i>Lanchocarpus capassa</i> ; Rain tree including <i>L. bussei</i> , <i>ericolalyx</i> & <i>nelsii</i> known as lance-pods	107
B2.20	<i>Mundulea sericea</i> ; Cork bush	107
B2.21	<i>Nicandra physalodes</i> ; Apple of Peru	107
B2.22	<i>Nicotiana rustica</i> ; Wild tobacco	108
B2.23	<i>Ocimum canum</i> & <i>urticifolium</i> ; Wild basil	108
B2.24	<i>Ricinis communis</i> ; Castor oil plant	108
B2.25	<i>Tagetes minuta</i> ; Mexican marigold	108
B2.26	<i>Tephrosia vogelii</i> ; Tephrosia	109
B 3	Some Easy Remedies from Plant Mixtures	
B3.1	Chilli & garlic	109
B3.2	Chilli, garlic & onions	109
B3.3	Chilli, Mexican marigold & onions	109
B3.4	Chilli, garlic, Mexican marigold & onions	109
B3.5	Garlic & Tephrosia vogelii	110
B3.6	Mexican marigold & syringa	110
B 4	More Plants with Potential	110
B4.1	<i>Albizia anthelmintica</i>	110
B4.2	<i>Albizia lebbek</i> ; Woman's tongue	110

B4.3	<i>Aleurites fordii</i> ; Tung tree	110	
B4.4	<i>Amaranthus spinosus</i> ; Pigweed	110	
B4.5	<i>Andropogon</i> spp; Blue, Snowflake & Stab grass	110	
B4.6	<i>Calotropis procera</i> ; Swallow wort	110	
B4.7	<i>Chenopodium album</i> ; Fat hen	110	
B4.8	<i>Clausena anisata</i>	110	
B4.9	<i>Combretum imberbe</i> ; Leadwood	110	
B4.10	<i>Dichrostachys cinerea</i> ; Chinese lantern	110	
B4.11	<i>Digitaria ternata</i> ; Finger grass & other spp	110	
B4.12	<i>Heteropogon contortus</i> ; Spear grass	110	
B4.13	<i>Khaya nyasica</i> ; Red mahogany	110	
B4.14	<i>Myrothamnus flabellifolius</i> ; Resurrection bush	111	
B4.15	<i>Securinega virasa</i> ; Snow berry	111	
B4.16	<i>Solanum panduriforme</i> ; Sodom apple	111	
B4.17	<i>Strychnos</i> spp; Monkey or Wild orange	111	
B4.18	<i>Tithonia rotundiflora</i> ; Red sunflower	111	
B 5	Summary of plants with insecticidal properties	111	
B 6	Summary of plants with repellent properties	111	
7	Late Entries	112	
7.1	<i>Cucumis anguria</i> ; Wild gherkin	112	
7.2	<i>Spirostachys africana</i> ; African sandalwood	112	
C	Baits, barriers, scarecrows and traps	113	
C1	Armyworm & other marching caterpillars	113	
C2	Banana weevil	113	
C3	Beetles, butterflies, grasshoppers & moths	113	
C4	Beetles	113	
C5	Birds, baboons & buck	113	
C6	Blister beetles	113	
C7	Caterpillars, grubs & slugs	113	
C8	Cutworm & cabbage root fly	114	
C9	Flies & fruit flies	114	
C10	Leaf miner	115	
C11	Locusts	115	
C12	Moths	115	
C13	Rats, mice, moles, rabbits & hares	115	
C14	Slugs, snails & squash bugs	115	
C15	Whitefly	115	
C16	Wild pig	116	
C17	Various insects	116	
<b>PART IV: REMEDIES FOR PROTECTING STORED SEED AND GRAIN</b>			
A	Controlling pests in stored seed (Small Amounts for replanting)	119	
1	Ashes, other fine powdery materials & sand	119	
2	Butter milk	119	
3	Cold	119	
4	Heat	119	
5	Hot water & air	119	
6	Naphthalene balls	119	
7	Oil	119	
8	Salt	119	
9	Smoke	119	
B	Controlling pests in stored grain (Large quantities)	120	
1	Non-plant materials	120	
1.1	Animal fat	120	
1.2	Fine powdery materials	120	
1.3	Manure	120	
1.4	Peels	120	
1.5	Sand	120	
1.6	Salt	120	
1.7	Solar energy	120	
1.8	Stirring	121	
2	Plant materials	121	
2.1	General	121	
2.2	Specific plants	121	
Recommended reading			124
Reference material			125
List of plants and their local names			127

Illustrated by Rose Elwell

© Henry Elwell

Published by the Natural Farming Network.

P.O. Box 301, Causeway, Harare, Zimbabwe with assistance from the Plant Protection Improvement Programme.

First published 1995 ISBN : 0-7974-1429-0

Printed by Mambo Press, Gweru



## The Natural Order

The soil and the air above it are teeming with life, both visible and microscopic. The existence of every single one of these life forms is essential to the natural order.

These forms of life are dependent upon each other for food, for mutual support and to develop competitive strength. They work together to produce an environment beneficial to their own existence and that of all other life forms, including human life. It can be said that without insects there would be no plants and we would not exist.

So in the natural order, all forms of life (insects, plants, animals and people) co-exist to mutual benefit, in mutual harmony and in perfect balance.

In such a situation insect numbers and types are controlled naturally, soil fertility is at its maximum potential, plant growth is vigorous and we reap the benefit of living in a healthy and supportive environment.

We have grossly under-estimated the importance of maintaining this natural order. Many current agricultural methods, particularly annual ploughing, burning, monocropping and the use of agro-chemicals, destroy this harmony. These methods reduce the numbers of some insects while encouraging others to increase and spread.

Pesticides were developed to combat the increased incidence of problem insects. Agro-chemicals appeared to work well but it soon became apparent that the chemicals were often more effective against the predators of the pest than against the pests themselves. Pest numbers increased. Worse was to follow when scientists discovered that the target pests themselves were becoming resistant to the very chemicals being used to eradicate them.

Attempts have been made to overcome these difficulties by applying stronger doses and more lethal products but pest populations have continued to increase, costs of control have escalated, the natural balance has been upset further, and greater numbers of people are now suffering from poisoning.

INSECTS ARE GOD-  
MADE. PESTS ARE  
MAN-MADE.

## Regaining Control

Regaining control does not mean eliminating all members of a particular pest from our fields and gardens. A certain level of pest presence is necessary to feed the predator population. Our aim is to minimise crop damage by regulating pest numbers.

The first step in regaining control is therefore to regard nature as an ally, to learn from it and to work with it. Nature has many lessons to teach us to our benefit. Our journey will be much easier — like floating downstream instead of forever battling upstream against the current.

Having learnt some of the more basic lessons nature has to teach us, the next move should be to re-establish a natural order of mutual benefit to insects, plants, birds, animals and ourselves.

### Befriending nature

Probably the most important lesson we have learned this century is that trying to fight nature is foolish, to co-operate is common sense. We have tried fighting with nature and found that not only do the current problems increase but it becomes more difficult to see what should be done next. When we co-operate with nature the reverse happens. Nature helps us to solve those very same problems and the way forward becomes clear.

The traditional form of agriculture once widely practised, called shifting cultivation, is a very good example of agriculture in harmony with nature. In those days farmers did not clear the land for the timber. Certain trees beneficial to society, providing fruit, medicine, wood, shade, spiritual needs and soil fertility were left untouched, while others were cut off at one metre above the ground. None of the tree roots were removed and this allowed them to re-grow quickly, returning the soil to full fertility in as short a time as possible after the farmer had moved to another piece of land. When the cut branches had dried, they were burned to clear the grass and so the ash could add fertility to the soil. The land was not ploughed but a form of zero tillage was practised, with seeds being planted into holes made with a sharp stick (later a hoe).

Several crops were planted onto the land in the same season, thus ensuring plant diversity, soil protection and soil fertility. In addition, the people had a sound knowledge of the basis of plant breeding for maintaining yield and for resistance to pests and diseases.

Prayers were offered at all important stages and a crop loss of

NATURE IS A FRIEND  
NOT AN ENEMY.  
TRY WORKING  
TOGETHER!

some 5-10% (to animals and insects) was accepted as a necessary contribution to maintaining the natural order or ecosystem. Immediately the land began to show signs of exhaustion it was allowed to rest for a lengthy period (about 20 years).

The people practising this system were clearly in harmony with nature in thought, word and deed. In those days, environmental degradation (soil erosion, soil exhaustion, pollution, loss of plant and animal species and deforestation) was unknown.

The success of this system can be judged from the fact that before writing this book a survey was carried out to record any surviving, truly indigenous, traditional knowledge on methods of pest and disease control other than the cultural methods inherent in the shifting agricultural system just described.

Although there was a little knowledge of protection of field crops against animals and birds, and some well established methods of protecting stored grain, there was a marked absence of knowledge on technologies for eradicating pests on the standing crop.

This absence of any technology for protecting field crops (although technologies had been developed for protecting grain) suggests that there was little need to protect field crops from pests and diseases under shifting agriculture. No such problems existed perhaps because the farming system did not disturb prey/predator balances.

Although shifting agriculture can no longer be practised because of a shortage of land, harmony can and must be reinstated. The principles governing harmony need to be researched by scientists and practised by the land user. In the meantime we must put to good use what we already know and use our powers of observation to learn as quickly as possible.

### Learning from nature

Nature has much to teach us. Nature is expert in zero tillage, in providing plant diversity, in recycling energy and nutrients through sunlight, animal wastes and vegetation, and in balancing prey/predator numbers. Our intelligence means we can learn from nature and then forge ahead to enhance natural processes for the benefit of all life forms and to make planet earth itself more fruitful.

On the general and detailed scales there is much to learn. Modern crop breeding methods to increase plant yields and for resistance to pests and diseases, biological methods of pest and disease control, and some genetic engineering initiatives, are examples of such beneficial developments. Although these technologies may be too specialised for the farmer to develop himself, there are no limits to the ways in which on-farm input-

outputs can be favourably altered through detailed observation and application of basic natural principles.

If, for instance, a plant is infested with insects, we can be sure that nature is teaching us an important lesson. We must examine our methods and determine whether the prey/predator balance has been upset or if the plant itself is less healthy than it should be.

If the plant is less healthy than it should be, examine soil fertility, the watering regime, plant hygiene (diseases), plant suitability or timeliness of planting. Look for clues from the colour and growth pattern of the plant above and below ground.

Examine the type of insect infesting the plants as this will indicate which predators are absent and should be encouraged back into the farming environment. The presence of large numbers of aphids, for example, is a sure sign that numbers of ladybirds, lacewings and hoverflies are too low.

In addition, observe the development of the insect through all its stages and note the duration and timing of each stage. This will assist in combating the insect as it can best be controlled at vulnerable stages in its life cycle. The most vulnerable stage of the maize stalk borer, for instance, is in the pupal form when it is hibernating in the bottom of the maize stalk. At this stage in its life cycle it can be eradicated very effectively by sun-drying the stalks in the field, by simply feeding the stover to livestock or by composting it.

Observe which crops are being attacked and at which times of the year as this knowledge can be used to avoid growing susceptible crops at peak pest periods.

Experiment with different crop mixtures in the same field to discover which protect each other and which do not.

All this information will lead to better selection of crop type or variety, of crop mixtures and rotations, and of time of planting.

Observe which plants, both wild and cultivated, are and are not attacked by certain insects. The ones that are not attacked may be useful for repelling those insects, whereas the ones which are attacked can be used as decoys.

Look carefully at the soil. Many and varied insects indicate a soil in good condition.

Learn what each insect, bird and animal eats, for that way we begin to understand the extent to which all life is linked and balanced. This knowledge will create within us a respect for nature which is evidently lacking at the moment.

Gradually through such observations made on our own land we develop a detailed knowledge which assists us to re-establish the natural order.

OBSERVE NATURE  
CLOSELY.  
THEREIN LIE THE  
ANSWERS.

FIGHTING WITH  
NATURE IS FOOLISH.  
CO-OPERATING IS  
PLAIN COMMON  
SENSE.



## Four Steps to Re-establishing the Natural Order

Ploughing or digging disturbs the balance among soil insects, fungi, viruses, bacteria and other soil life. Soil disturbance breaks up root channels and soil structural units so that the soil cannot perform its intended function. Disturbance of the soil also causes a rapid loss of organic matter upon which the crop and many soil organisms feed.

Soil is the foundation of agriculture yet on most of our lands it is at its lowest possible fertility level. In such circumstances the balance between the many and varied life forms in the soil is bound to be dangerously disturbed.

### 1 Re-instate soil fertility

The first step in re-establishing the natural order is therefore to reinstate the fertility of the soil by minimising soil disturbance and maximising soil organic matter.

### 2 Provide habitats for predators

The second step is to provide suitable natural habitats for predators and to preserve existing ones. Here agro-forestry can play an important role; and a diversity of wild plant species should be encouraged on gravelly or rocky outcrops and on field edges. There is great scope for using contour ridges for a variety of purposes: for growing fruits, firewood, fodder and trees as windbreaks, all of which will also help to re-establish the pest/predator balance.

### 3 Re-introduce plant diversity

The third step is to re-introduce plant diversity into the cropping system as diversity is nature's most effective way of maintaining the pest/predator balance and of ensuring continued soil fertility. In place of monocropping, farmers should consider ways of increasing plant diversity through the use of rotations, intercropping, mixed cropping, strip cropping and permaculture designs.

### 4 Gradually reduce agro-chemicals

The fourth step is to gradually reduce the amounts of agro-chemicals used as fertilizers, herbicides, insecticides and fungicides. Substitute organic matter and manure in place of mineral fertilizer; use rotations, underplanting and mechanical cultivations in place of herbicides; and adopt the methods recommended in the rest of this book to replace the harmful chemicals currently used for pest and disease control.

1  
REINSTATE THE  
FERTILITY OF THE  
SOIL

2  
PROVIDE AND  
PRESERVE NATURAL  
HABITATS.

3  
REINTRODUCE PLANT  
DIVERSITY.

4  
USE NATURAL  
REMEDIES TO  
REPLACE AGRO-  
CHEMICALS.

## 1 Reinstating soil fertility

A healthy soil produces healthy plants and healthy plants resist attack from pests and diseases.

To improve the soil *dig less, mulch more and compost always.*

### i DIG LESS

Start by digging deeply to break up surface compaction, to remove hard layers in the soil profile and to incorporate initial compost applications. In subsequent years, disturb the soil as little as possible. Rely on crop roots, insect life and surface mulch to maintain the soil in good condition. Occasional digging, ripping, fork lifting or ploughing may be necessary in the initial stages until the soil structure has improved sufficiently; or if the soil has become compact due to foot, wheel or hoof movement when the surface is moist.

Soil fertility and organic matter content can be improved by growing more leguminous crops. Certain legumes can serve several useful purposes. Cowpea, for example, will not only improve soil fertility, structure and permeability but also produces edible leaves and beans.

The initial application of compost can be incorporated into the soil but subsequent applications should be applied to the soil surface around the base of the plants.

### ii MULCH MORE

Mulching is the process of covering the soil with a layer of plant residues of varying thickness (thickness depending on the purpose of the mulch and the availability of suitable material) to conserve soil moisture, keep the surface cool, protect it from erosion, prevent the development of surface crusts, enhance infiltration, improve soil structure, minimise compaction, suppress weeds and to add nutrients to the soil through decomposition (in-situ composting).

Choose seedless materials as mulches and obtain the material from as many diverse plants and sites as possible to provide a wide range of nutrients. Mulch as heavily as possible wherever possible but first make sure the soil is in a fit condition to be mulched. If it is a wet soil, drain it first or make raised ridges or beds.

The combined effect of disturbing the soil less, of applying compost and of mulching causes soil life to flourish. Soil structure quickly improves, but needs more nitrogen during the initial stage. The yellowing leaves of the plants and reduced growth warns us that nitrogen is limited.

This condition can be rapidly corrected by watering the plants

LOOK AFTER THE  
SOIL AND IT WILL  
LOOK AFTER YOU.

MULCH MORE TO  
SAVE MOISTURE,  
WORK AND WORRY.

every few weeks with liquid manures, made preferably from fresh chicken manure mixed in water. Mix liberal amounts of fresh manure in water (2 to 4 heaped handfuls to a 10 litre bucket), and water into the soil around the base of the plants. Avoid pouring water over the plant leaves as the nitrogen may burn them. The manure can be added directly to the water, stirred and applied immediately when a quick nitrogen boost is required, or put into a cloth (an old sock is ideal) and soaked in water in a closed tub for 3-7 days before use, to allow nutrients to be dissolved. During the rains, a little dry manure can be placed near the plant stem but not too close.

It is true that mulches encourage certain pests (like armoured crickets and slugs). Some people avoid mulches for that reason. But residues also discourage other pests (such as cutworm and nematodes) and do far more good than harm. Termites are thought to be a major problem but this is not the case. Certainly they are attracted by the mulch but do not attack healthy living plants when there is an abundance of residue for them to feed off. In fact the termites do considerable good to the soil by rapidly breaking the mulch down and carrying it into the soil to form organic compounds. Their tunnels improve soil aeration and thus increase the water infiltration rate. When termites are prevalent, however, mulch should be cleared from around the immediate area of the plant stems, particularly of woody plants.

Residues which are infested with pests should not be used. Such material should be put on the compost heap where the heat of composting will kill the pest, or the residue should be fed to livestock and the resultant manure used to make compost.

### iii COMPOST ALWAYS

Compost is nature's fertilizer formed out of decomposition of plant, insect and animal residues and wastes. Applications of it are essential to maintain the soil in good structural condition so that it is fertile, easy to work, resistant to erosion and also for pest and disease control.

The key to good compost making is to collect as wide a range of suitable materials as possible: manures, crop residues, grass, leaves, weeds, and so on, to ensure a varied supply of nutrients. Sufficient manure and/or green matter must be added to create hot conditions in the heap.

The watering must also be just right. Apply only sufficient water to moisten the materials so that a drop or two of moisture (no more, no less) can be squeezed from a handful. Mix water in very thoroughly by watering and turning a small amount of the heap at a time as if mixing cement. When thoroughly moist, place

COMPOST ALWAYS  
FOR SOIL FERTILITY.

the material in a thin layer on the heap. Don't take the short-cut of placing the materials on the heap dry and then trying to water them. It never works.

Build the compost heap in the shade and near to a water supply if possible; and turn it every two weeks. Cover it with a thin layer of soil in the dry season to reduce evaporation and to minimise fly breeding.

Good quality compost can be made during the rains. Keep turning the compost after every storm until the heap is moist throughout. Cover it with a plastic sheet, thatching grass, or a dry-mud skin so that too much rainwater does not destroy the heat of composting and prevent the correct breakdown of plant materials.

The amount of compost required to obtain top yields varies from about 10-20 tonnes per hectare; 20 tonnes per hectare when the soil is unmulched and in poor structural condition and 10 tonnes per hectare when the soil is mulched and in good condition. The compost should not be broadcast but banded or spread next to the plant.

The addition of compost from field-edge heaps may be reduced considerably if plant diversity is maintained and if sufficient densely planted legumes are included in the rotation. The legumes should be selected to supply soil nitrogen, to maintain organic matter levels and to provide mulch when the plants have been slashed down. This is often called 'sheet composting'.

## 2 Provide natural habitats for predators

Removal of the natural vegetation on farms and around gardens has greatly disturbed the pest/predator balance. Nationwide the essential food supplies of birds, animals and insects have been reduced to the extent that they have no option but to feed off the crops. Very often the natural vegetation has been so depleted by deforestation and overgrazing that the cropland offers the only source of food.

The opportunity for encouraging the natural habitat to re-establish on unused areas such as field edges, rocky or gravelly outcrops and elsewhere on farms is very great indeed. The natural vegetation will normally re-establish itself on these areas if allowed to do so; but by careful selection of trees, shrubs and grasses in these locations this unused land can be made even more productive.

Shelter belts of mixtures of indigenous trees around farm fields or on stony areas in the fields will do much to protect the lands from drying winds, provide wood for fuel and building purposes and habitats for insect predators.

ENCOURAGE A  
NATURAL HABITAT  
ON UNUSED AREAS.

ESTABLISH SHELTER  
BELTS OF TREES  
AROUND FARM  
FIELDS.



Similarly, there is great scope for increasing production by growing fruit and fodder trees, shrubs and grasses along the contour banks. The trees will act as wind breaks, provide firewood and encourage the predators into the lands, particularly bird life. Certain indigenous trees should also be encouraged to grow on the contour banks. Choose trees or shrubs (whether indigenous, or exotic) carefully, avoiding those with shallow, spreading roots likely to compete with the crop. In many of the dry parts of the world, shelter belts have been found to dramatically increase crop yields by reducing evaporation rates.

### FULLY USE CONTOUR BANKS.

### 3 Re-introduce plant diversity

One major cause of soil exhaustion is that too many grains are being grown and too few legumes. In addition, the inclusion of too much white flour in our diet (like maize and wheat) causes constipation, sluggishness and stomach complaints. The area under grain crops could be reduced with benefits to both soil conservation and human health!

When the same crop is planted on the same piece of land year after year, the soil suffers from constant removal of specific nutrients and pests and diseases build up. For example, stalk borer and red spider mite numbers build up rapidly in maize and tomato (or cotton) fields respectively when these crops are grown continuously.

This happens simply because the law of plant diversity has been ignored. Each specific insect flourishes because the host crop provides an abundant food supply and continuous shelter for it. The situation is worsened by chemical spraying programmes which eradicate the natural predators of the pests.

The law of plant diversity says that there should be a wide variety of different plants on the land at any one time. Violation of this law brings swift repercussions particularly in the tropics and subtropics.

We have only to look at nature to see how this works, how each plant supports and protects its neighbour. In the undisturbed bush on the edges of the cropped area we find plants of every kind and height living harmoniously together. Here and there a plant may have some insects on it but the predators are also evident in sufficient numbers to prevent an outbreak.

Each plant of one kind is separated from another of the same kind so that insects on one plant cannot easily get to the other. The space between them is filled with other plants, all of which protect each other in some way either by acting as physical barriers or by repelling the insects through the aroma released by their leaves, flowers or roots. This is happening both above and

below ground with the roots acting as repellants or barriers to keep soil-borne pests and diseases in check.

In addition, each plant helps to create an entire habitat (moisture, organic matter, nutrients, insect life, wind protection, and so on) suited to the production of the maximum amount of plant material at that place.

Scientists say that there is also an element of competition between the plants for moisture, sunlight and nutrients. But in virgin bush such competition is healthy and favours the growth and reproduction of strong plants of each type and represses the weaker specimens. The competition becomes 'unhealthy' only when plants of the same species are grown together as a field crop or when two or more poorly matched plants are put close together. This does not happen in nature.

Research has found that plant diversity decreases the number of harmful insect species while at the same time attracting a greater number of enemies of the existing pests; and that one plant protects another different type of plant by obscuring that plant's shape and odour so that the pests cannot easily find it. Also, on the whole, plant diversity has been found to reduce the incidence of plant diseases.

Let us now look at some of the ways nature's plant diversity can be imitated.

#### i ROTATIONS

Rotation should be considered as the first step towards providing the necessary plant diversity. A rotation is a sequence of different crops grown on the same field one after the other.

For example, a three-year, rain-fed rotation may consist of maize-soyabean-finger millet or rape-green maize-tomatoes in a vegetable garden.

Rotational sequences were once chosen with the purpose not only of minimising pests and diseases but also of improving soil fertility and preventing erosion. However, in recent years, the needs for fertility and erosion prevention have been pushed aside and when rotations are practised, they are selected for pest and disease control only. The need to emphasise pest control is yet another indication of the extent to which modern agricultural practices have upset the natural pest/predator balance. This over-emphasis on pests and diseases can be corrected once some positive steps towards restoring the balance have been taken.

Pay more attention to including legumes, fodder crops and weed fallows in rotations to provide the combined benefits of soil fertility, erosion control and sustained food production. The value of weed fallows for all these purposes, including pest and

GROW MORE  
LEGUMES AND LESS  
GRAINS.



ROTATIONS ARE THE FIRST STEP TOWARDS PLANT DIVERSITY.

disease control, has been grossly under-estimated in the past. This is largely because soil fertility is currently so poor (because of the annual ploughing/chemical farming approach) that growth on weed fallows has been impoverished.

ii MIXED CROPPING


In mixed cropping, planting diversity in any one year is increased by growing several plants on the same land at the same time. The types of plants which grow well together are carefully chosen, for example in the maize-soyabean-finger millet rotation mentioned in the previous section, the maize rows (with normal spacing between rows unaltered) may be underplanted with pumpkin and cowpeas.

Apart from pest and disease control benefits, mixed cropping minimises the risk of total crop failure, suppresses weeds, protects the soil and improves the amount of organic matter in the system.

One old variation was to mix the seed of the different plants together in the correct proportions and broadcast them. This method avoids continuous rows of the same kind of plant down which pests and diseases can readily travel but is not suited to farms which rely on machines for planting, weeding and harvesting.

Our ancestors had a great deal of knowledge of plant combinations which protected each other. Most of this knowledge has

MIXED CROPPING-  
PUMPKIN-COWPEA-  
MAIZE



been lost but in recent years researchers have become aware of the value of this skill and are gathering the relevant information together. Some combinations which have been found to reduce crop losses to pests and diseases are: cowpea with cassava or sorghum; maize with sunflower, potato and mustard; okra with tomato, ginger and mung bean; kale with tomato and tobacco; pumpkin, okra, cowpea, bambara nut, melons and watermelons with maize, sorghum or millet; maize and cowpea; and cotton with cowpea, maize or sorghum. Nematode damage on citrus trees has been reduced by planting legumes underneath (cowpea being mentioned in particular); and fruit trees have been protected from the same pest by planting acacia trees nearby.

Even mixing together varieties of the same crop has been found to reduce crop losses to pests and diseases.


Lantana, cosmos, caryopteris, marigolds, asters, chrysanthemums, garlic, tomatoes, onions and most herbs, are known to protect other plants.

iii INTERCROPPING

Intercropping is similar to mixed cropping except that the chosen crops are planted in alternate rows, sometimes in pairs of rows.

The wide spacing between the rows of plants of the same kind give some protection from the spread of insects but pests can still travel

INTERCROPPING:  
MAIZE-SUGAR BEANS



easily along the crop rows. Nevertheless intercropping has been found to reduce crop losses compared to monocropping. For instance, intercropping maize with cowpea has been found to reduce the incidence of legume pod borer on the cowpea. Similarly, insect damage on cotton has been reduced by intercropping cotton with cowpea, maize or sorghum; and certain beans have been found to reduce powdery mildew on cassava, while the cassava protects the beans from angular leaf spot.

#### iv STRIP CROPPING

The advantages of rotations, mixed cropping and intercropping have been combined in a new system called no-till strip-cropping.


In this system no ploughing is done. A scratch mark 50mm deep is made to plant the row crop (e.g. maize) and the underplanted crop (e.g. pumpkin, cowpea, dolichos lab lab) is put into a hoe-hole positioned between rows.

Plant diversity is increased by having the entire rotation on the field at the same time laid out in strips along the contour, between each pair of contour ridges.

A rotation of maize-soyabean-finger millet is shown in the sketch below as it may appear on the field in the first year. In the second year, the bottom crop moves up to the top of the field in place of the top crop and all the other strips move down-slope to make room for it.

Every year the same method is followed with the bottom strip

STRIP CROPPING:  
M=MAIZE  
S=SOYABEANS  
F=FINGER MILLET



moving up to the top of the field.

In this way, not only does the entire rotation appear on the field at once but each strip of soil receives a different crop from year to year.

The maize strips can be underplanted to a variety of suitable crops (like pumpkin, cowpea, watermelon, squash and beans) to increase plant diversity and to obtain all the other benefits of mixed cropping. The possibilities are almost endless. Even tomatoes have been planted successfully in maize, with the tomato plants tied up to the maize stems.

#### v PERMACULTURE DESIGNS

Plant diversity can be further increased by growing together a wider range of plants of varied growth habits and habitat preferences than envisaged under any of the previously-mentioned systems. The plants are chosen not simply because they provide food, but because they will protect the food crops from insect attack, or give the right soil condition, air temperature, sunlight and humidity to maximise production of the food crops planted among them.


Permaculture designs may include several heights (stories) of vegetation from trees to shrubs to creepers to small plants arranged so that each creates the environment required by the others. Plants requiring or tolerant of shade (like sweet potato or dust berries) are grown under the trees and shrubs, while those needing sunlight occupy the open spaces.

Care is taken not to have too many similar plants growing next to each other and to ensure sufficient numbers of multi-purpose plants for soil improvement (legumes) and protection against insects. Strong-smelling plants such as tomato, onion, garlic,

PRACTISE MIXED,  
INTER & STRIP  
CROPPING.

APPLY  
PERMACULTURE  
DESIGNS WHERE  
POSSIBLE.

PERMACULTURE  
DESIGNS





marigold and so on, are planted among or around other plants such as cabbage, rape and lettuce to protect them from insect attack.

Other plants attractive to pests are deliberately grown to trap the pests.

#### 4 Use natural remedies to replace agro-chemicals

Widespread dependence on agro-chemicals has a detrimental effect on the on-farm and off-farm ecosystems, ultimately on production levels and inputs as well as on human health and welfare.

Soluble agricultural fertilizers cause loss of soil organic matter and structure, deplete numbers of useful living organisms in the soil and encourage acidification and loss of essential trace elements. We should rely more on natural fertilizers such as organic matter and manures.

Herbicides are designed to kill off unwanted plant life on the soil surface but also destroy microscopic plant life in the soil responsible for breaking down plant residues and for maintaining the pest/predator balance. Fungi, for example, predate on nematodes. Wherever possible, weed control by mechanical cultivation, mixed cropping techniques and rotational practices (particularly growing dense cover crops) are preferable to chemical treatments.

Newly introduced insecticides are highly efficient at first in eradicating pests but the effectiveness reduces with time due to two factors. On the one hand the predators which would normally reduce insect pest numbers are themselves effectively killed off, sometimes more effectively than the target pest. Secondly, the pests surviving the treatment are resistant to the chemical used and continue to breed. The new off-spring are also resistant to the insecticides and numbers of insect pests increase.

The counter-strategy of the past has been to either increase the number of times the crop is sprayed, to increase the dosage or to devise new, costly and more potent chemicals. Research has shown that these strategies merely produce an increasingly resistant insect population. Weeds also build up resistance to herbicides and much the same strategies have been tried without success.

The ineffectiveness of these strategies can be judged from surveys such as the one which found that compared to thirty years ago crop losses to insect attack have doubled in spite of ten times more insecticides being used.

The indirect effects of insecticides on crop production levels are more difficult to assess but yields are said to have been adversely

affected by the loss of large numbers of beneficial insects involved in pollinating crops, eating pests, maintaining soil fertility and in supporting the ecosystem in general. In addition, food quality has deteriorated from contamination of the edible material.

Not only has the use of pesticides affected crop quantity and quality but one survey found that an estimated eleven million people are hospitalised each year in Africa alone due to insecticide poisoning. Also there is mounting evidence overseas that recommended levels of agro-chemicals (fertilizers, insecticides and herbicides) can lead to serious pollution of surface and groundwater supplies.

Various initiatives have been taken to reduce the dangers of insecticide use. Governments have banned many of the more poisonous chemicals and restricted the use of others to prevent further pest-resistance build-up and regulations have been tightened up. These initiatives are emergency measures and do not get to the root of the problem: the ecological damage caused by poisons over the decades.

On the more positive side, management systems are being developed to bring about significant reductions in the amounts, potency levels and frequency of insecticide use, the best known being Integrated Pest Management (IPM). So far the developed Integrated Pest Management systems are relatively sophisticated and more applicable to large-scale commercial enterprises than to small-scale subsistence farming. These systems include: more frequent use of crop rotations and leys; biological controls such as breeding and release of predators or pollinators or the deliberate spread of diseases; planting of resistant crop types and varieties; and genetic engineering techniques to introduce sterility into the pests, to improve the aggressiveness of predators or to increase the resistance of plants.

The main aim of early Integrated Pest Management systems was to lower pest control costs rather than to protect the environment but recent emphasis has shifted towards environmental protection, with pesticide use being relegated to providing back-up when all else has failed.

The development of these systems is clearly beneficial but the insecticides themselves and the Integrated Pest Management packages developed so far are rarely within the budget or skills of the small-scale farmer so an alternative, subsistence-level approach is required to entirely safeguard the environment.

This book promotes an alternative technology called 'the natural method'. It is based on observing and applying nature's own methods backed up by simple remedies available at no or low cost.

APPLY IPM SYSTEMS  
AS A TRANSITIONAL  
STAGE WHERE  
APPROPRIATE.

WHEREVER POSSIBLE  
REPLACE ALL  
CHEMICALS WITH  
NATURAL REMEDIES.

## Supportive Strategies

### Hygiene

Cleanliness is extremely important in preventing the spread of insects and diseases, especially diseases.

Firstly make certain that the seedbed soil does not contain harmful numbers of pests and diseases. A soil which has been treated correctly over the years will be free of them but suspect soils may be sterilised by burning branches on the prepared surface (the ash helping to fertilize the soil or it can be used in pest control), watering the ground with liquid made by crushing wild basil (*Ocimum canum*) leaves in water, steaming the soil or solar heating the ground under plastic sheeting (Part III A29 ii).

A disease-free seedbed is essential to prevent disease outbreaks among the plants in the nursery or after transplanting them.

Seeds and transplant material should be free of pests and disease otherwise they will not only suffer themselves but will infect other plants in the field. One old tactic still in use is to store the seed with burnt cow dung or wood ash so that it is disease free at planting.

Should outbreaks occur, all infected plant material should be removed. This material can be composted if adequate heat in the compost heap can be guaranteed, failing that it can be fed to livestock or buried well below ground. Burning the material will solve the problem too but would be a waste of valuable organic matter unless the ash can be used for insect control.

Hands and tools should be cleaned after the infected plant material has been removed. If this is not done the problem will be spread to other plants. Remember also that the spores of plant diseases can be carried on clothing and footwear.

The water used for liquid manures, for sprays and infusions or for watering the plants, should be from a non-infected source. Water in which hands, tools and plants have been washed, or placed, will be infected.

### Select resistant varieties

Research organisations are breeding plants resistant to specific pests and diseases. However, growers can do their own plant breeding simply by collecting the seed from healthy plants in the field which have not suffered from insects or diseases.

The seeds from these plants will produce vigorous growth and the plants growing from them will have a good chance of being

NEVER LEAVE  
INFECTED MATERIAL  
LYING AROUND.

ADOPT THE GOOD  
HABIT OF  
CLEANLINESS.

better suited to the local environment and of being more resistant to pests and diseases. Indeed, in the past, through keen observation of nature, traditional farmers have played an important part in developing varieties of crops resistant to a wide spectrum of pests and diseases. Since natural systems are dynamic and not static, selection for breeding resistance must be an on-going process, otherwise the hard-gained resistance will be lost. In the case of subsistence farmers, breeding can best be done by the farmers themselves in their own fields.

### Timing of planting

Planting can be timed to minimise attack by pests and diseases. To do this effectively the grower needs to know the life cycles of the pests and the conditions favouring the spread of diseases.

For example, vegetable crops do better in winter than in summer because many of the pests and diseases are either dormant or less active in the cold weather. The major pest, root knot nematode, is relatively inactive at this time.

Another way of improving plant survival is to grow seedlings in protected nurseries and plant them out in the field only when they are large enough to withstand attack. Damage by snails, cutworms, beetles and hoppers can be minimised in this way.

Many insects after remaining hidden in the soil through winter and spring, emerge with the first rains. Damage to summer crops can be minimised by either establishing the crop earlier or by planting a few weeks after the rains.

### Plant spacing

Plant spacing can be varied to some extent to hinder the spread of pests and diseases, particularly diseases.

In some cases control can be achieved by increasing the humidity of the air through growing plants closer together. The activity of certain insects, red spider mites for example, is reduced by dense planting.

On the other hand, most fungi thrive better in the humid atmosphere created between closely-grown plants and plants should be spaced further apart where fungi may be problematical.

However, in the absence of mulch, wide spacing reduces overall yield, exposes the soil to the damaging effects of erosion and sunlight and encourages weed growth. Therefore, rather than resort to wider spacing, it would be better to ensure vigorous plant growth so that the plants have maximum resistance to attack, and to control the fungi by the methods described later in this book. Diseases

DEVELOP THE SKILL  
OF GROWING YOUR  
OWN SEED.

TIME YOUR  
PLANTING TO AVOID  
THE PESTS.

PLANT CLOSE AND  
INSPECT THE CROP  
OFTEN.



especially are more easily controlled if caught in the early stage therefore regular and frequent inspections should be carried out.

### Insect barriers and decoys

Plants can act as physical barriers to the movement of pests. Hedges hinder the movement of aphids into the garden; a few rows of maize can protect a bean crop against aphids; and a row of pigeon peas has been used to protect tomato, potato and cabbage crops against red spider mite.

Another approach is to trap insects on decoy plants. The decoy may consist of weeds allowed to grow in the crop or may be susceptible plants grown in rows around the food crop. The insect prefers the decoy plant and infests it, leaving the cash crop untouched. The infested plant can then be removed and composted or fed to the livestock. For example, aphids are attracted to milkweed and a food crop will be protected from attack if a few of these weeds are left to grow in the land.

Beans have been planted as decoys in rows around cabbages or cotton to protect these cash crops from red spider mite. The infested bean can then be fed to the livestock or composted or chopped up and left on the surface as a mulch. This crop serves as an ideal decoy for it provides three functions: insect control (decoy), improvement of the soil (legume) and food for livestock or material for mulching or compost.

Barriers and decoys can also provide suitable habitats to encourage the predators of field pests.

### Encourage predators

In the early stages of establishing the natural order, there will be a deficiency of predators. Most predators will have been killed off by current farming practices and wholesale destruction of their natural habitats.

In western countries predators are bred on insect farms and sold to the farmers in bulk for them to release into their lands. This practice is spreading into Africa and elsewhere but is an unnecessary expense. Mostly these predators survive for only a short time before they are killed off by the very farming practices that eradicated them in the first place. The correct approach is to create conditions conducive to increasing existing predator numbers to levels which mean they can carry out their task of keeping crop-eating insects in check.

The value of reinstating the soil's fertility, improving plant diversity and of avoiding the use of agro-chemicals to re-

CONSIDER USING  
MULTI-PURPOSE  
BARRIERS AND  
DECOYS.

establish the natural order (and thereby increase predator numbers) has already been mentioned, as has the importance of providing diverse habitats on contour ridges, rocky outcrops and at field edges. To re-establish the natural order is perhaps the safest approach to pest control as the subjective judgement of whether a certain insect or animal is or is not beneficial is avoided.

Many texts provide lists of beneficial creatures so that farmers can avoid killing them. The selection is often arbitrary as some eat insects at one stage of their lives and vegetation at another. Without doubt most insects are beneficial at some stage. For instance in one study of the 86,000 insects identified on farms, 76,000 were found to be friends of the farmer.

Purists might argue that all insects are beneficial since every one of them forms part of the food chain. Nevertheless it will do no harm and maybe some good to list those considered to be more beneficial than others because they play an obvious and direct role in controlling pests which feed on plants. The ones most commonly identified are: anthocorid bugs, ants, assassin bugs, bats, bees, birds, black-kneed capsids, branchid wasps, chamelions, dragon flies, dung beetles, earthworms, frogs, ground beetles, hawk moths, hedgehogs, hoverflies, ichneumon flies, lacewings, ladybirds, lizards, mice, moles, nematodes, praying mantises, spiders, stick insects and toads.

The provision of bird baths and crumb or seed trays will encourage a large variety of birds back into the neighbourhood and, provided the habitat is suitable for them, they will remain in the vicinity to do their job of keeping problem insect numbers in check.

Another important step in improving and maintaining predator numbers is to refrain from killing them for short-sighted reasons, as this is what created the problem in the first place. For example a farmer might notice that birds of prey are killing his chickens. Instead of protecting his hens with wire mesh, 'scare-crows' or tree cover, he kills the hawks and destroys their nests and habitat. Numbers of rats, mice and seed-eating birds which the hawks would have kept in check then increase and consume the farmer's grain both in the field and in storage. As a result he and his family suffer from hunger and poverty simply because he did not think carefully enough before he acted. This is only one small example.

For decades, through annual ploughing, destruction of the natural habitat, indiscriminate use of agro-chemicals and thoughtless destruction of insect life in many other ways, we have upset the delicate but beneficial balance of nature and are now suffering the consequences.

INITIALLY  
ENCOURAGE AN  
INCREASE IN  
PREDATOR NUMBERS.

ULTIMATELY AIM TO  
RESTORE THE  
NATURAL ORDER.

## Summary

### LET NATURE CONTROL PESTS.

The only sensible approach is not to eliminate all pests but to restore the natural balance. Learn from nature how to maximise food production and minimise crop losses.

- 1 Avoid using pesticides. They are poisons. They pose grave threats to human health, upset the natural order and balance and, in the long term, cause increased crop losses.
- 2 Replace agro-chemicals with natural remedies.
- 3 Encourage predators to breed. Do not kill them. Rather provide them with suitable habitats.
- 4 Imitate nature by providing as much plant diversity as possible.
- 5 Dig less, mulch more, compost always. The basis of plant protection is to take care of the soil. A healthy soil is less likely to harbour pests and diseases at harmful levels and produces healthy crops better able to resist attack.
- 6 Be hygienic. Do not assist diseases by spreading them.

These are the foundation blocks of good pest and disease management. By practising them the number and severity of attacks by pests and diseases will decrease.

It is accepted, however, that infestations will occur from time to time, particularly in the early stages when growers are attempting to restore nature's balance. Do not give in to the temptation to reach for that can of insecticide or else you may well undo all the progress you have made so far.

Rather select one of the natural remedies, identified in the rest of the text, appropriate to the problem pest to be controlled. With few exceptions, these natural remedies are simple to use, safe, and cost next to nothing.

The key to success is to use even natural chemicals only in an emergency. And when you do, avoid killing off the useful predators. Therefore scout regularly, spray selectively. Learn to recognise friendly insects and encourage them to stay.

In addition, it is important to recognise that the development of any skill depends entirely on the degree of care and concentration the farmer or gardener devotes to cultivation practices: to keen observation and attention to the needs of the soil, the plants and the insect and bird life. All must be in complete harmony with the growers and so the growers' attitudes are a major factor governing the success of their attempts to co-operate with those subtle but powerful forces at the seat of all creation.

## PART II

# PESTS, DISEASES AND CONTROL MEASURES



## The Advantages of Natural Pest and Disease Control

Major advantages to the methods of pest and disease control described in this book are:

- 1 It is more difficult for pests and diseases to develop resistance to combinations of plant-derived chemicals than to single synthetic chemicals.
- 2 Growers can choose whichever method suits them and the locality best.
- 3 The plant preparations and other methods described in this book are simple, easy to prepare and safer to handle than current pesticides.
- 4 The methods given here can be made by the growers on their home ground at little to no cost.

Most of the substances listed in this book do not give dramatic 'kills'. Indeed complete eradication of insects would disrupt the natural balance and be highly undesirable. The objective of 'control' should be merely to reduce numbers of the problem insect to levels resulting in insignificant or acceptable crop losses.

The few natural remedies that give dramatic kills can be poisonous to human beings and animals, as well as to insects and should be handled with extreme caution. For safety purposes, these remedies have been identified by the words 'See Warning' placed directly after the subheadings in Part III Section B of the text.

### THE CORRECT APPROACH

The correct approach to pest and disease control is as follows:

- 1 Practise the techniques given in Part I.
- 2 Assess whether the infestation is serious enough to warrant any action or whether it is at a harmless level appropriate to the natural balance.
- 3 If the outbreak is considered serious enough to warrant prompt action, identify the pest concerned from the descriptions given in the following pages and apply the recommended remedies. If the pest cannot be exactly identified then identify it by its general form, for example, beetle, caterpillar, fly, eggs and so on, and use the control methods given for a similar form of insect detailed in this text.
- 4 Consider carefully the cause of the outbreak and using the information given in Part I as a guideline, seek out and correct the fault. Some helpful check notes are given at the end of each section dealing with specific pests.

This section lists the host plants of each selected pest, describes the damage done by the pest, gives its life cycle and some details to help the user identify the pest or disease. Suggestions or remedies for controlling the pest are then given. These have been drawn from the literature and from sources of local experience. Finally, check notes are provided to help the farmer avoid future outbreaks.

Unnecessary repetition has been avoided by a simple code which enables the user to find the detailed remedies given in Part III. The code consists of the Key Name (the first word in the descriptive heading), followed by the section and subsection numbers. For example, the remedy to the code **Clay etc. (A10 i)** can be found in Part III under the heading **Clay, Lime & Other Powdery Materials** given under Section **A item 10 Subsection i** which begins: 'Dust the fine material....'.

Before making the plant preparations detailed in Part III B, read *Guidelines for the use of botanical sprays* given immediately below the main heading, 'B Plant Materials'.

## Common Pests

### AMERICAN BOLLWORM (*HELIOTHIS ARMIGERA*)

#### Host Plants

Barley, beans, brinjals, coffee, cotton, crucifers, cucurbits, garden flowers, grapes, groundnuts, lucerne, maize, peas, potatoes, sorghum, soyabean, sunflowers, tobacco, tomatoes, wheat, citrus and deciduous fruit trees.

#### Damage

The *Heliothis* bollworm larvae feed on the buds and young bolls of the cotton plant. It is capable of destroying the entire crop if not controlled. The caterpillars also feed on the buds, blossoms and fruit of a wide variety of plants, particularly those listed above. Very young fruit may be completely destroyed while large holes may be eaten into mature fruit.

#### Life cycle

The female moth lives for 3 weeks and lays up to 1500 eggs on the leaves, buds and flowers of host plants. The small whitish-yellow eggs are laid only at night. They hatch in 3–6 days into caterpillars which live for approximately 3 weeks in warm weather passing through 5 stages. The caterpillars then drop to the ground to form pupae which live for 2–3 weeks in the ground before developing into moths. Lacewing larvae and spiders predate on the bollworm.

#### Identification

The caterpillars vary in colour, being dark beige when young but later becoming green, brown or even reddish pink. They grow to about 35mm long and have a characteristic pale yellow longitudinal stripe down each side.

#### RECOMMENDED REMEDIES

**Moths:** Aromatic plants (A1); Biological (A4 i); Light traps: Various insects (C17 i); Repellent sprays (B6). The bodies of the moths caught in the light traps (Various insects: C17) can be used as a biological spray (A4 i).

**Eggs:** Ash (A2 i, ix, x, xi); Clay etc (A10 i, v); Hand picking (A15 ii); Oil-mineral (A21 i); Pyrethrum (B 1.6); Tobacco (B 1.17); Rain tree (B 2.19); Cork bush (B 2.20); Tephrosia (B 2.26).

**Caterpillars:** Biological (A4 i, ii); Glues (A13); Hand picking (A15 i); Hens etc. (A16); Oil-mineral (A21 i — see warning); Potassium permanganate (A22); Salt preparations (A23 ii); Smoke etc. (A24 i); Soap solutions (A25 ii); Insecticidal plants (B5).

Of the above, Hand picking and Hens are likely to be the most appropriate for removing caterpillars from small areas; while for larger



AMERICAN BOLLWORM





areas. Mineral oil and Potassium permanganate may be more suitable. In addition, for both large and small areas, out of the insecticidal plants listed under B5 useful sprays can be made particularly out of Pyrethrum (B 1.6), Tobacco (B 1.17), Cork bush (B 2.20), the Rain tree (B 2.19) and Tephrosia (B 2.26).

**Pupae:** Hens etc. (A16) to dig the pupae from the soil; and strong sprays from Pyrethrum (B 1.6), Tobacco (B 1.17), Cork bush (B 2.20), the Rain tree (B 2.19) and Tephrosia (B 2.26) should prove useful.

#### Checks

Ensure an adequate habitat for owls, nightjars, bats and other night creatures that predate on the moth; and for birds that will feed on the caterpillars during the day. Many other insects feed on the eggs and larvae of this pest so encourage as wide a range as possible of insect life both above and below ground. Plant diversity is also vital.

#### ANTS (*FORMICIDAE*)

##### Host

Honeydew and insects.

##### Damage

The small black ants which encourage aphids are those which most interest the farmer or gardener. They are attracted to the honeydew produced by the aphid and are said to transport young aphids from one location to another thereby spreading diseases carried by the aphids. The ants also scare away predators. Many other species of ants assist the gardener by killing off insect pests such as cutworm, stalk borer, spider mites, termites and many others.

##### Life cycle

Ants are small insects which nest underground, usually making use of existing cavities under rocks and among roots. A colony consists of a king and a queen and a large number of workers. The queen lays clusters of small, white eggs which are then moved to safe locations by the workers. In addition to feeding on the honeydew secreted by aphid and scale insects, ants feed on a wide variety of other insects many of which are potential pests (like cutworms, mites, termites, slugs and snails).

##### Identification

All ants live in colonies. They are small, six-legged creatures with their bodies divided into three segments by two thin waists. There are between 8,000 to 14,000 different species of ants in the world. Most ants build permanent nests in the soil and have poor eyesight; however, they have a very keen sense of smell and vibration which enables them to operate in a highly efficient manner. Ants release chemicals which enable them to recognise companions from the same colony, to raise the alarm or to sting enemies.



ANTS

#### RECOMMENDED REMEDIES

**Eggs:** if the ant problem has become severe, search for their nests which will be close at hand and destroy the eggs. Birds and Hens etc. (A16) will also help to control ant and egg numbers.

**Ants:** different species of ants respond to different treatments and, since it is almost impossible for the layman to distinguish between the thousands of species, users need to determine which of the treatments below give the best results in their circumstances.

Those with the most general application are:

Aromatic plants (A1); Biological (A4 i); Clay etc (A10 iii); Grease (A14 i or ii); Garlic (B 1.2); Chilli (B 1.4); Tomato (B 1.12); Syringa (B 1.15); Tobacco (B 1.17); Rhubarb (B 1.19); African marigold (B 1.20); Blackjack (B 2.8); Lantana (B 2.15); Lippia (B 2.17); Mexican marigold (B 2.25); Rain tree (B 2.19); Cork bush (B 2.20); Wild tobacco (B 2.22); Tephrosia (B 2.26)

The small black garden ants which harvest the aphid have been found to be most effectively deterred by a spray made from blackjack seeds or from Mexican marigold leaves. The whole Mexican marigold plant dug into the soil at planting will deter the ants from seedlings. Sprays made from the leaves of rhubarb, tomato and tephrosia will exterminate all ants.

Others mentioned in the literature are: Soyabean (B 1.10); Oleander (B 1.16); Mexican poppy (B 2.5); Wormseed (B 2.10); Sodom apple (B 2.16); Castor oil (B 2.24).

Small grains sprinkled around vegetable seeds have been found to lure red ants away from the seed which would otherwise be eaten. Penny royal and mint can be planted as barriers to prevent red ants from eating the roots of other plants.

Worth considering are general insecticides and repellants B5 and B6. Also some species of house-loving ants have been repelled effectively by baby powder and lime.

##### Checks

Ensure that numbers of aphids and scale insects are under control as the ants are attracted to the honeydew secreted by them.

#### APHIDS (*APHIDIDAE SPP*)

##### Host plants

Most plants, shrubs and trees, particularly the tender young shoots.

##### Damage

Aphids are sap feeders. In large numbers they can remove sufficient sap to kill off leaves and the growing tips. They are very important vectors of plant viruses. The honeydew they produce attracts a large number of other pests (like ants and fruit flies).

##### Life cycle

Young aphids are produced directly from the mother without the egg



APHIDS

stage. There are no males in the warmer areas of the world. These insects are usually wingless but will produce wings and fly to other plants when the host plant becomes overcrowded. Flying aphids can be carried long distances by the wind. Young aphids are said to be spread by the ants feeding on their honeydew. Hoverfly larvae and ladybirds are said to eat up to 250 aphids per day. Other predators are lacewing larvae, parasitic wasps, the praying mantis and small birds.

#### Identification

Adult aphids are only a few millimetres long. They are round and plump and the colour of each species is different, varying from yellow, green, brown, grey, white and black. The woolly aphid looks like a small piece of cottonwool. All species tend to cluster close together on the tender parts of the plant. They are soft-bodied and can be squashed easily with the fingers. Aphids form wings only when they are overcrowded and need to migrate.

#### RECOMMENDED REMEDIES

Growers are strongly recommended to grow healthy plants and to use compost in preference to manures, including liquid manures. Healthy plants develop their own in-built resistance to aphid attack.

Manures and mineral (artificial) fertilizers, particularly nitrates, produce sappy and fleshy plant tissue attractive to aphids.

Aphids can be controlled successfully by increasing plant diversity on the land particularly by mixed, inter and strip cropping, and by Permaculture designs (see *Part I, 3 Re-introduce plant diversity ii, iii, iv, v*); and by trap cropping and barriers (see *Part I, Supportive strategies*).

Garlic, chives, marigold, nasturtiums, parsley, onions and many herbs grown as companion plants are said to repel certain species of aphid, particularly greenfly. Nasturtiums grown as companion plants are recommended for repelling woolly aphid or as a spray to kill them, and pigeon pea is said to repel green and grey aphids.

Milkweed, sowthistle and black nightshade are good trap or decoy plants to attract aphids away from the main crop.

#### Supplementary measures

Because aphids prefer young tender shoots, take care to ensure concentrations of the natural remedies do not harm the plant growth. Supplementary remedies with the most general application are:

Ash (A2 i, ix, xi); Clay etc. (A10 i, ii, iv, v); Manure etc. (A17 i, ii, iii); Oil-mineral (A21 i); Potassium permanganate (A22); Salt preparations (A 23 i, ii); Soap solutions (A25 i, ii); Vinegar (A30); Water (A31 i); Garlic (B 1.2); Chilli (B 1.4); Pyrethrum (B 1.6); Sweet potato (B 1.11); Tomato (B 1.12); Tobacco (B 1.17); Sweet basil (B 1.18); Annona squamosa (B 2.3); Custard apple (B 2.4); Blackjack (B 2.8); Lantana (B 2.15); Lippia (B 2.17); Wild basil (B 2.23); Mexican marigold (B 2.25); Tephrosia (B 2.26); Leadwood (B 4.9); Cucumis anguria (B 7.1);

African sandalwood (B 7.2); Various insects (C17 iii).

Aphids and ants go together. Ants are said to spread the aphids and to scare off predators. The key to controlling aphids is to get rid of the ants as well. Thus the best control is usually given by clay and lime sprays, blackjack seeds or Lippia, Mexican marigold, rhubarb, tomato and Tephrosia leaves (see also *Ants*).

Tobacco in dust and liquid spray forms is said to be effective. Others mentioned in the literature are:

Glues (A13); Onion (B 1.1); Tea (B 1.3); Pawpaw (B 1.5); Soybean (B 1.10); Cassava (B 1.13); Feverfew (B 1.14); Syringa (B 1.15); Rhubarb (B 1.19); Catunaregam (B 2.9); Rubber hedge (B 2.13); God's tobacco (B 2.18); Castor oil (B 2.24); Plant mixtures (B 3.1 to 3.6).

A strong solution of onions poured around carrots, lettuce, parsley, green beans and groundnuts when the soil is moist is said to protect them from root aphid.

Worth considering are general insecticides and repellants B5 and B6.

#### Checks

The presence of large numbers of aphids on a plant is an indication that the plant is not healthy. Ensure that the plants are strengthened by applying the right quantity, quality and frequency of compost and water to the soil. Check the soil for phosphorus deficiency. Also ensure that the plants are grown from good seed and show a vigorous growth right from the start. Feed them with Compost (A12 i) rather than Manure etc. (A17 i, viii) foliar sprays. Encourage the presence of hoverfly, praying mantis, spiders and ladybirds.

#### ARMYWORM (SPODOPTERA EXEMPTA)

##### Host Plants

Barley, oats, wheat, sugar cane and grasses/pastures, citrus, deciduous fruit trees and grapes, beans, brinjals, cotton, crucifers, cucurbits, ground-nuts, maize, millets, peas, potatoes, sorghum, soyabean, tomatoes and flowers.

##### Damage

A major pest in outbreak years, these caterpillars move in swarms and can do great damage to crops and grassland. They feed mainly on the grass family of plants but will eat any green plant material when hungry, including cotton bolls. They move and feed during the day and at night.

##### Life cycle

The parent insect is an inconspicuous night-flying moth with brown marked forewings and off-white hindwings. The eggs are laid in clumps of about 200 or more on food plants on the underside of the leaves (grasses and millets, being preferred) and are covered with down from the body of the mother. The eggs hatch in 3 to 4 days. There are 5-6 caterpillar stages; development is completed in about 18-21 days in summer. The caterpillars occur in two forms: the



ARMYWORM





gregarious outbreak phase, which feed voraciously, and the solitary phase, which feed less heavily. In the solitary phase the caterpillars occur in small numbers, are sluggish, do little damage and can be found feeding at the base of plants of the grass family. A feature of the gregarious phase outbreaks is their unexpected suddenness. Large areas may be found to harbour swarms of caterpillars virtually overnight. When fully grown to about 25mm, the caterpillar enters the soil, constructs a thin-walled earthen cell and pupates for 12–14 days. The whole life cycle from egg to adult moth may be completed in a little over a month. After a single generation has come and gone, an infected area may be clear of them as the moths migrate once their food source has been depleted. In this way, from generation to generation, they may migrate long distances. Many birds eat the caterpillars while flies and wasps parasitise the caterpillars, eggs and pupae. Ladybirds are said to eat armyworm eggs. Guinea fowl are particularly good predators.

#### Identification

The armyworm is a caterpillar which crawls over the surface of the ground. They may occur in vast numbers (gregarious phase) mostly in December or January and less often up to April. The young caterpillars are smooth and vary considerably in colour. In the gregarious phase the young caterpillar is greyish-green, becoming black above with thin blue lines down the middle of the back; and several yellow lines along the sides. The underparts are mostly apple green. They have the tendency to drop to the ground when disturbed. In the solitary phase the caterpillars are greenish or brownish in colour.

#### RECOMMENDED REMEDIES

With armyworm perhaps more than with any other pest it is vital to keep numbers down to a minimum at all stages of the life cycle. Thus, ensure an adequate habitat for owls, nightjars, bats and other night creatures that predate on the moth; and for birds that will feed on the caterpillars and pupae during the day. Many other insects feed on the eggs and larva of this pest, so encourage as wide a range as possible of insect life both above and below ground. Plant diversity is also vital.

Low value grain crops like finger millet can be used as trap crops which can be destroyed along with the armyworm infesting it. This is perhaps a more preferable strategy than keeping the lands free of all host plants including weeds (as recommended in some texts) because the hungry caterpillars will attack almost any green plant in the absence of their preferred food.

Interplanting with stagger weed, a minty legume, has been found effective.

Regular and careful scouting can reveal eggs and young caterpillars on the field edges or on the contour banks. If the grass is thick enough there may be sufficient food off the arable land for it to

complete its life cycle without damaging the crop. To prevent future problems, however, the armyworm should be destroyed by any of the supplementary methods outlined below.

The two most important preventive measures are :

- 1 Avoid burning the grassland at all costs.**  
Burning the veld in winter not only degrades the soil and grass in the long term, but also is a prime cause of outbreaks of large swarms. As soon as the temperatures begin to rise, eggs are laid in very large numbers on the fresh new grass sprouting from the burned veld.
- 2 Avoid overgrazing.**  
Overgrazing in summer removes the natural food store of the caterpillar and it has then no option but to invade cropland in search of food and habitat.

#### Supplementary measures

**Moths:** Aromatic plants (A1); Biological (A4 i); Light traps: Various insects (C17 i); Repellent sprays (B6).

The bodies of the moths caught in the light traps (Various insects: C17) can be used as a Biological repellent (A4 i) spray.

**Eggs:** Ash (A2 ix, x, xi); Clay etc. (A10 i, v); Hand picking (A15 ii); Oil-mineral (A21 i); Pyrethrum (B 1.6); Tobacco (B 1.17); Rain tree (B 2.19); Cork bush (B 2.20); Tephrosia (B 2.26). Make sure to spray under the leaves.

**Caterpillars:** Biological (A4 i, ii); Soap solutions (A 25 ii); Garlic (B 1.2); Syringa (B 1.15); Clausena (B 4.8) and Trenching (C1) have been particularly mentioned in the literature. But Glues (A13); Hand picking (A15 i); Hens etc. (A16); Oil (A21 i, ii — see warning); Potassium permanganate (A22); Salt preparations (A23 ii); Smoke etc. (A24 i); and plants listed as Insecticidal (B5) are other possibles, particularly sprays made from Pyrethrum (B 1.6), Tobacco (B 1.17), the Rain tree (B. 2.19), Cork bush (B 2.20) and Tephrosia (B 2.26).

Of the above, Hand picking and Hens are likely to be the most appropriate for small numbers of caterpillars, removal of which will prevent swarms in future. The caterpillars vary in colour when there are only one or two of them, being greenish or brownish.

When the caterpillars are moving in large swarms it is important to stop the swarms before they reach the arable land.

**The following measures are highly recommended:** spraying with vegetable oils (away from the crop) or mineral oil (on or near to the crop); liberally dusting the caterpillars and their food with lime; since many diseased specimens can be found trailing behind a large swarm, Biological sprays (A4 ii) to spread the disease are a useful alternative; Trenching (C1) can be applied as an additional emergency measure along with crushing them underfoot, repeatedly driving animal herds over the swarm, beating them with stiff branches, setting fire to the grassland if it is dry enough, or setting fire to dry grass and branches placed over them.

**Pupae:** Hens etc. (A16) to dig the pupae from the soil; and strong sprays from Pyrethrum (B 1.6), Tobacco (B 1.17), the Rain tree (B 2.19), Cork bush (B 2.20) and Tephrosia (B 2.26), should prove useful when watered thoroughly into the soil.

#### Checks

Check that as many preventive measures as possible have been implemented to prevent swarms; unburnt grassland, no overgrazing and restoration of the pest/predator balance. Take steps to eradicate small infestations as these are the origin of future swarms.

### BEAN STEM MAGGOT OR BEAN FLY (*OPHIOMYIA PHASEOLI*)

#### Host Plants

The maggots attack beans of various species (french, haricot, dolichos and ratic) and other leguminous plants.

#### Damage

Young plants can be seen wilting and dying off. Investigation will reveal decayed, swollen and hollow stems at ground level with a small maggot inside. The leaves of the plant may show yellow spots where the adult (fly) has laid her eggs. On mature plants the grub may be located higher up the stem. Velvet beans (*mucuna*) are not attacked. No major attacks on cowpeas have been reported.

#### Life cycle

The female is a tiny black fly which pierces holes in the leaves of the plants and lays eggs inside the tissues. The eggs hatch in 2–4 days. The maggots eat their way down the stem to just above ground level. There they complete their development. The grown maggots then pupate and become adult flies in about 20 days. Under favourable conditions, the entire life cycle from egg to fly takes 21 days. Many of the insects spend winter as pupae and emerge as adults in November. The greatest damage is caused from November to January inclusive. A parasitic wasp called *Opius liogaster* can suppress the pest from February onwards.

#### Identification

The best way to identify the insect causing the damage is through the type of damage: wilting of the plant with hollowing and rotting of the stem at ground level and the presence of the maggot or pupa in the stems. However, early warning of the presence of this pest is given by the presence of yellow blotches on the leaves forming around small punctures made by the fly when it laid its eggs.

#### RECOMMENDED REMEDIES

The old saying that prevention is better than cure applies well to this pest. The best approach is to prevent the fly from laying its eggs by companion planting with aromatic plants, mixed cropping, or by routine spraying with repellants made from aromatic plants; or by



BEAN STEM MAGGOT



egg-laying bean fly

planting outside the period November–January. Planting very early in summer will avoid the worst period for attacks.

An old remedy is to build up the soil around the plant stems to encourage roots to form above the already damaged part at original ground level but this is a 'rather-too-late' tactic.

#### Supplementary measures

**Fly stage repellants:** Manure etc. (A17 ii); Garlic (B 1.2); Tomato (B 1.12); Goat weed (B 2.2); Apple of Peru (B 2.21); Mexican marigold (B 2.25).

**Fly stage predators:** fly numbers can be dramatically reduced by bird life so encourage birds with a bird bath and seed tray; Hens etc. (A16) will also do their part if allowed to roam through the arable areas. Bantams are particularly fond of insects and although they may peck occasionally at leaves, they do far more good than harm during summer when insects are plentiful. Nevertheless do not give them access to seedlings.

**Eggs:** Ash (A2 ix, x, xi); Clay etc. (A10 i, v); Oil-mineral (A21 i); Pyrethrum (B 1.6); Tobacco (B 1.17); Rain tree (B 2.19); Cork bush (B 2.20); Tephrosia (B 2.26).

**Maggots and pupae:** because the maggot burrows into the plant material as soon as it has hatched, it is not possible to treat it directly; however, future numbers of the fly can be reduced effectively by taking off and destroying leaves showing the yellow blotches where the fly has laid its eggs. This should be done by pinching through the leaf stem as close to the main stem as possible, thus intercepting the maggot before it has had time to reach the main stem. In addition, remove and burn all plants which show signs of serious stem damage from maggots. Also the stems of damaged plants should not be left in the field over winter otherwise the pupae will survive, but should be composted or fed to livestock.

#### Checks

Ensure an adequate habitat for bird life that predate on the fly and that there is a wide variety of insects to predate on the eggs, maggots and pupae. Have you given enough thought to planting date, rotations and to planting resistant varieties? Some local cowpea cultivars can tolerate this pest by thickening the undamaged strands of tissue in their stems.

### BEETLES — BLISTER: (*MYLABRIS SPP.*)

#### Host Plants

Beans, groundnuts, maize, peaches, peas and a wide range of garden flowers.

#### Damage

The beetles destroy the flowers of plants, especially on the pea and bean family, thus preventing pods and fruit from forming. They have been recorded attacking maize silks and tassels. They are beneficial,



BEETLES—BLISTER



however, in the pre-adult phases as they eat the eggs of locusts and grasshoppers. The large colourful CMR beetle is about the best known blister beetle.

#### Life Cycle

Blister beetles are common throughout Africa. There are many different species but they are all rather large, varying from 20–50mm in length. The female lays about 20–30 large, oval, white eggs in hard, dry soil. She lays the eggs in batches in different sites until she has exhausted her supply, whereupon she dies. The eggs hatch in 3–4 weeks into tiny, active, six-legged larvae 1.5mm long, white with a large head and two large bristles on the tail. These larvae run over the surface of the soil seeking places where locusts and grasshoppers have laid their eggs. They then burrow down to the eggs and spend the rest of their lives feeding off them. Having found the egg store, a larva sheds its skin turning into a fat white maggot with very short legs. It continues to feed on the locust or grasshopper eggs and grows rapidly. Towards the end of summer it becomes an inert larva and passes the winter in this state. The skin of the larva is tough and white and its short legs are still evident. It sheds its skin once more when summer arrives, eats some more of the eggs and darkens in colour. In 3–4 weeks it sheds its skin for the last time and emerges from the soil as an adult beetle.

#### Identification

These large, colourful beetles can be seen eating flowers during the summer months and can be identified readily from their large size (20–50mm) and bright colours. Most of the beetles are black and yellow, black and red or all black. They walk and fly deliberately. A chemical in their bodies blisters the skin. They may also be readily identified in the active juvenile stage when running over the surface of the ground, or in the inert larval stage when resting in the soil.

#### RECOMMENDED REMEDIES

**Beetles:** these large beetles, not usually appearing in large numbers together, are best treated by simple methods: Aromatic (A1) to repel them; Biological (A 4 i); Hand picking (A15 i); Beetles (C3) and (C4); Blister (C6). Hens usually avoid them.

Hand picking (wear gloves) or trapping usually prove sufficient. The CMR beetle is said to be particularly attracted to blue and red colours.

**Eggs:** mulches, cover crops and keeping the soil surface damp all discourage egg laying but farmers should avoid destroying the eggs as the larvae are very helpful.

**Larvae:** do not destroy as they do much good — see *life cycle*.

#### Checks

These insects are beneficial on the whole and efforts should be made to provide alternative habitats for them such as flowering borders to

the fields, trap crops etc. When they invade the crop do not destroy all of them but keep numbers in check.

#### BEETLES — LEAF (CHRYSOMELIDAE)

##### Host Plants

Apples, cowpeas, creepers, gooseberries, sunnhemp and a variety of garden plants.

##### Damage

These beetles skeletonise the leaves leaving only the main veins untouched. Others chew shot-holes in the leaves (like flea beetles). Some species such as the gooseberry beetle also eat fruit, roots or flowers. Leaf eaters often show a distinct preference for certain plants, some preferring creeping plants such as cowpea. Generally both the larva and adult beetle damage plants.

##### Life cycle

Some types of leaf beetle lay their eggs on the leaves while others lay them in the soil. After the eggs have hatched, the grubs feed ravenously on the plant material. The larvae then pupate, again either on the leaves or in the soil and emerge later as adults. The eggs typically hatch in about 6–12 days, the larvae reach full size in about another 10–12 days and the pupa form can take a week. Five or six generations can be produced in a year.

##### Identification

This very large family of beetles is difficult to characterise as they vary greatly in shape and colour. However, most of them are small. They are best identified simply by observing the beetle eating the vegetation. Most are very active, smallish insects (6–9mm) which fly away rapidly when disturbed. Some are a rather ordinary colour: brown, dull yellow or black, while others, like the fool's gold beetle, are brightly coloured.

#### RECOMMENDED REMEDIES

Since these beetles are small and can appear in large numbers, spraying is more practical than the simpler methods recommended for the blister beetles.

**Beetles:** Aromatic (A1); Ash (A2 ix, xi); Biological (A4 i); Clay etc (A10 i) — particularly lime; Hens etc (A16).

Insecticides recommended in the literature as effective against various beetles are: Garlic (B 1.2); Syringa (B 1.15); Oleander (B 1.16); African marigold (B 1.20); Goat weed (B 2.2); Mexican poppy (B 2.5); Blackjack (B 2.8); Wormseed (B 2.10); Lantana (B 2.15).

\* Also worth trying are any of the insecticidal plants mentioned in B5 and particularly sprays from Pyrethrum (B 1.6), Tobacco (B 1.17), the Rain tree (B 2.19), Cork bush (B 2.20) and Tephrosia (B 2.26).

Routine spraying with mixtures of repellent plants from B6 may prove a useful preventive measure.



BEETLES — LEAF

**Eggs on the leaf:** Ash (A2 i, ix, x, xi); Clay etc. (A10 i, v); Hand picking (A15 ii); Oil-mineral (A21 i); any of the insecticides listed in B6 particularly those \* mentioned previously for controlling the adults.

**Eggs in the soil:** Ash (A ii); Hens etc. (A16) to dig the eggs from the soil; and strong sprays from Pyrethrum (B 1.6), Tobacco (B 1.17), the Rain tree (B 2.19), Cork bush (B 2.20) and Tephrosia (B 2.26), should prove useful when watered thoroughly into the soil.

**Pupae on leaves:** Ash (A2 ix, x, xi); Clay etc. (A10 i, ii, v); Hens etc. (A16); any of the insecticides listed in B6 particularly those \* mentioned previously for controlling the adults.

**Pupae in the soil:** Ash (A2 ii); Hens etc. (A16); Mulches (A19 i) aromatic; and again particularly those \* mentioned above for controlling the adults.

#### Checks

Ensure an adequate habitat for bird life that predate on the beetles and that there is a wide variety of insects to predate on the eggs and pupae. Have you given enough thought to planting date and rotations? Tree, hedge and plant barriers will often prevent the beetles from flying into the field. For the same reason intercropped or mixed cropped plants suffer less.

### BEETLES — SNOUT (*CURCULIONIDAE SPP*)

#### Host Plants

Apples, aloes, citrus, coffee, grapes, groundnuts, maize, millet, peaches, pears, plums, soyabeans, strawberries and many others.

#### Damage

The beetles of this very large family chew the edges of the leaves leaving ragged margins. The stems of young branches can be eaten causing the fruit to drop or the insects may create shallow lesions on the fruit itself. The larvae (grubs) of some species may bore into stems or bulbs, or burrow under the bark of trees causing the plants to die.

#### Life cycle

The life cycle of one of the maize snout beetles serves as an example. Adults emerge from the soil 4–6 weeks after the first substantial rains and feed off the edges of the leaves at night. They become sexually mature after about 10 days and batches of about 30 eggs are laid in the leaf folds. These hatch approximately one week later into tiny, white, legless grubs which drop to the ground and enter the soil where they feed on roots. The grubs hibernate in earthen cells during the winter. With the onset of the next rainy season, the grubs once again become active for about three weeks and damage the hypocotyls of recently-germinated maize plants, causing them to remain stunted and to exhibit symptoms of phosphate deficiency. The grubs then pupate in the soil, emerging as adults some two weeks later. The adults are unable to fly. They feed at night and, during the day, hide under clods of soil or in the funnels of the maize plants.



BEETLES — SNOUT

#### Identification

Snout beetles are part of the extremely large weevil family with 60,000 known species. There are several types of snout beetles varying in size from 5–25mm long. All have the characteristic elongated snout of the weevil family, hard bodies and dark colours: grey, brown or black. The antennae are generally short and bulbous at the end.

### RECOMMENDED REMEDIES

Good weed control combined with planting the crop three weeks later is said to give maize good protection against the maize snout beetle.

Weed fallows can be used as trap crops and cowpea and velvet beans in rotation are said to control the insect.

**Beetles:** the larger beetles of this family can be best dealt with by Hand picking (A15) in the early morning, late evening or on cloudy days, by Hens etc. (A16), or by catching them at night in light traps: Various insects (C17 i). The smaller beetles are more troublesome and can appear in very large numbers. Their numbers can be depleted substantially by light traps: Various insects (C17 i). Hens etc. (A16) are also useful for getting rid of those appearing during daylight hours. All sizes can be repelled by use of strong-smelling plants Aromatic (A1) or by the Biological (A4 i) method. Damage to the leaves can be reduced by dusting the leaves with lime (Clay etc A10 i) and those living in the soil can be trapped by dusting the soil surface around the plants: Ash (A ii), Clay etc. (A10 ii, iii).

The stronger Soap solutions (A25 ii) have been found useful against certain small beetles.

Insecticides recommended in the literature as effective against various beetles are: Garlic (B 1.2); Syringa (B 1.15); Oleander (B 1.16); African marigold (B 1.20); Goat weed (B 2.2); Mexican poppy (B 2.5); Blackjack (B 2.8); Wormseed (B 2.10); Lantana (B 2.15).

These would have to be sprayed or dusted (as per the recipe) on the plants at night or watered thoroughly into the soil during the day.

\* Also worth trying are any of the insecticidal plants mentioned in B5 and particularly sprays from Pyrethrum (B 1.6), Tobacco (B 1.17), the Rain tree (B 2.19), Cork bush (B 2.20) and Tephrosia (B 2.26).

Routine spraying with mixtures of repellent plants from B6 may prove a useful preventive measure.

**Eggs on the leaf:** Ash (A2 i, ix, x, xi); Clay etc. (A10 i, v); Hand picking (A15 ii); Oil-mineral (A21 i); any of the insecticides listed in B6 particularly those \* mentioned above controlling the adults.

**Eggs in the soil:** Ash (A2 ii); Hens etc. (A16) to dig the eggs from the soil; and strong sprays from Pyrethrum (B 1.6), Tobacco (B 1.17), the Rain tree (B 2.19), Cork bush (B 2.20) and Tephrosia (B 2.26), should prove useful when watered thoroughly into the soil.

**Pupae/larvae on leaves:** Ash (A2 ix, x, xi); Clay etc. (A10 i, ii, v); Hens etc. (A16); any of the insecticides listed in B6 particularly those \* mentioned previously for controlling the adults.



**Pupae in the soil:** Ash (A2 ii); Hens etc. (A16); Mulches (A19) aromatic; and again particularly those \* mentioned previously for controlling the adults.

**Pupae/larvae in the stems, bark or roots:** dead or seriously damaged plant parts should be removed/dug up and destroyed. Old bark can be scraped off trees and builders' lime brushed on (A7 ii); or the exposed surface sprayed with either agricultural lime (A10 iv, v) or any of the insecticides listed in B6 particularly those \* mentioned previously for controlling adults.

#### Checks

Ensure an adequate habitat for bird life that predate on the beetles and that there is a wide variety of insects to predate on the eggs and pupae.

#### CUTWORMS: (NOCTUIDAE)

##### Host plants

Cutworms attack young seedlings (mainly) of beans, brinjals, coffee, cotton, crucifers, cucurbits, groundnuts, maize, onion, peas, potatoes, soyabeans, tobacco, tomatoes and many other plants and crops.

##### Damage

This smooth, greyish caterpillar emerges from the soil at night, encircles the plant with its body and cuts through the stem of young plants just above ground level, leaving the wilting top to die. They will climb up mature plants and cut off lateral growth but usually do minimal harm at that stage of the plant's development. They are general feeders and will attack almost any kind of plant. A single caterpillar may cut down several seedlings in one night.

##### Life cycle

The dull-coloured moth emerges in spring and lays its eggs on the soil or on the foodplants. A single female may lay 600–800 eggs which hatch in approximately thirteen days into tiny black caterpillars. The young caterpillars initially feed on plant leaves for ten days before burrowing into the soil. Thereafter they live in the soil during the day, lying in a curled up position usually within 10–25mm of the surface, emerging only at night to feed. The fully grown 30mm long caterpillar burrows approximately 25mm and turns into a smooth reddish-brown pupa. It remains in the soil for 2–3 weeks before emerging as a moth. There may be up to four generations. Many insects predate on the cutworm at all stages in its life cycle. For example, a small parasitic wasp lays its eggs in the caterpillar and the wasp larvae gradually eat the cutworm.

##### Identification

The sight of the seedling top or severed laterals lying on the ground surface is often the first hint of the presence of the cutworm. A small hole may be found in the soil near the severed plant which, when excavated carefully, will reveal the curled up smooth greasy-grey caterpillar.



CUTWORMS



#### RECOMMENDED REMEDIES

A wide variety of simple but effective measures are available for controlling this pest.

**Moths:** Aromatic plants (A1); Biological (A4 i); Clay etc. (A10 x); Mulches (A19i, ii); Light traps (Various insects: C17); Repellent sprays (B6).

The bodies of the moths caught in the light traps; Various insects (C17) can be used as a Biological (A4 i) spray.

Liming acid soil is said to deter the moth from laying its eggs or to reduce egg fertility. Mulching also deters moths from laying eggs on the soil.

**Eggs:** Ash (A2 i, ii, ix, xi); Clay etc. (A10 i, v); Hand picking (A15 ii); Oil-mineral (A21 i); Pyrethrum (B 1.6); Tobacco (B 1.17); Rain tree (B 2.19); Cork bush (B 2.20); Tephrosia (B 2.26); Leadwood (B 4.9); African sandalwood (B 7.2). Make sure to spray or dust both the upper surface and underside of the leaves.

**Caterpillars:** Ash (A2 i, ii); Clay etc. (A10 iii); Hand picking (A15 iv); Hens etc. (A16); Manure etc (A17 i, iii); Mulch (A19 i, ii); Smoke etc (A24 i); Chilli (B 1.4); Pawpaw (B 1.5); Finger millet (B 1.8); Thorn apple (B 2.11); Castor oil (B 2.24); Sodom apple (B 4.16); Combretum imberbe (B 4.9); Spirostachys africana (B 7.2) and Cutworm (C8) have been specially identified in the literature as effective means for controlling the cutworm.

One of the simplest and most effective control measures is to concentrate hens or bantams (Hens A16) on to the beds before planting.

The squashed ripe fruit of Sodom apple steeped in water and dug into the soil is said to control cutworm.

Plants listed as Insecticidal (B5) are other possibles, particularly sprays made from Pyrethrum (B 1.6), Tobacco (B 1.17), the Rain tree (B 2.19), Cork bush (B 2.20) and Tephrosia (B 2.26).

The tin can and stick methods described in Cutworm etc. (C8 iii and iv) are effective emergency methods for small and medium size areas and very simple to apply. For larger areas, the best emergency approach is to water well into the soil insecticides made from the plants mentioned previously.

For small infestations dig the cutworm out the soil with a sharp stick. Carefully search around the base of the cut plant to find the small entrance hole to its hideout. Dig along the tunnel until the grub is found.

Another simple technique is to delay transplanting seedlings until the stems are too wide for the cutworm to encircle, too hard or woody for it to cut, or too strong-smelling to attract it.

Repeated disturbance of the soil surface is also said to reduce caterpillar numbers but, as this can have a detrimental effect on soil fertility, it should be a last resort.

Several sources recommend baits as alternatives. These can be made out of maize flour and water with plant poisons, pyrethrum and

rotenone (C8 ii) added in.

Another source has recommended watering around the plants with a mixture of one teaspoon of grated turnip having cooled after one litre of boiled water has been poured over it.

#### Checks

Check the pH of the soil; acidity is said to encourage the pest. Is there sufficient mulch? Ensure an adequate habitat for owls, nightjars, bats and other night creatures that predate on the moth; and for birds that will dig up the caterpillars during the day. Many other insects feed on the eggs and larva of this pest, so encourage as wide a range as possible of insect life both above and below ground. Plant diversity is not so vital as the caterpillars feed on almost all cultivated plants.

#### DIAMOND-BACK MOTH (*PLUTELLA XYLOSTELLA*).

##### Host plants

Cabbages, cauliflowers, other members of the cabbage family (crucifers) and some garden plants.

##### Damage

The small greenish caterpillars about 8mm long eat numerous small holes into the plant leaves which seriously affects the appearance and vigour of the plant. The leaves may be reduced to a network of veins in the case of severe infestations. It is active particularly in the dry season.

##### Life cycle

The ash-grey female moth lays her tiny yellowish-green eggs either singly or in small groups on the upper surface of the leaves. They hatch in a few days into small green caterpillars which move to the underside of the leaves to feed. They grow rapidly, attaining full growth in about 10–30 days, depending on temperature. The 12mm long caterpillars spin silk cocoons under the leaves in which they pupate. The adult moth emerges about one week later. There may be as many as ten generations a year and the female may lay 50 or more eggs. As a result their numbers can multiply very rapidly when conditions are favourable. Green and brown lacewings, parasitic wasps, spiders and larvae of hover flies are important predators of the caterpillars; while fungal diseases play their part in reducing the numbers of these pests during the rainy season.

##### Identification

Numerous small holes in the plant leaves are usually the first signs of the pest. The small, green caterpillar can then be seen on the underside of the leaves, lying under a protective web. The caterpillar has the characteristic habit of wriggling actively when touched. The moth is a small grey insect. When the moth is resting, a pattern of three diamonds can be seen along the line where the wings meet over the back.

#### RECOMMENDED REMEDIES

Summer rains and frequent irrigations reduce the mating of the moths



DIAMOND-BACK MOTH



and wash off young caterpillars and the pupae. Strongish sprays need to be directed on to the underside of the leaves to remove pupae.

Growing resistant cultivars, planting mustard as a decoy or trap crop and biological control through diseased specimens, have worked well.

Rotations effectively control moth numbers. Numbers of caterpillars have been reduced considerably by breaks of over 6 weeks during which no crucifers have been grown.

Dusting or spraying leaves with agricultural lime (Clay etc. A10) effectively controls eggs, caterpillars and pupae.

#### Supplementary measures

The following remedies have been specifically recommended in the literature for general control of the pest in all phases: Potassium permanganate (A22); Salt preparations (A23 ii); Garlic (B 1.2); Pyrethrum (B 1.6); Tomato (B 1.12); Feverfew (B 1.14); Syringa (B 1.15); Oleander (B 1.16); Tobacco (B 1.17); Sweet basil (B 1.18); Custard apple (B 2.4); Desert date (B 2.6); Catunaregam (B 2.9); Tephrosia (B 2.26).

**Moths:** Aromatic plants (A1); Biological (A4 i); Light traps (Various insects C17 i); Repellent sprays (B6).

The bodies of the moths caught in the light traps: Various insects (C17) can be used as a Biological (A4 i) spray.

**Eggs:** Ash (A2 i, ix, xi); Clay etc. (A10 i, v); Hand picking (A15 ii); Milk (A18 i); Oil-mineral (A21 i); Pyrethrum (B 1.6); Tobacco (B 1.17); Rain tree (B 2.19); Cork bush (B 2.20); Tephrosia (B 2.26). Make sure to spray particularly the upper surface of the leaves.

Treat crucifers routinely at regular intervals to prevent the eggs from hatching if unsightly holes in the leaves are to be avoided.

**Caterpillars:** Ash (A2 ix, x, xi); Biological (A4 i, ii); Clay etc. (A10 i, v); Glues (A13); Hens etc. (A16); Oil-mineral (A21 i-see warning); Potassium permanganate (A22); Salt preparations (A23 ii); Smoke etc. (A24 i); Soap solutions (A25 ii); Garlic (B 1.2); Syringa (B1.15); and Clausena (B 4.8) have been mentioned in the literature as possible controls for caterpillars. Plants listed as Insecticidal (B5) are other possibilities.

Once the caterpillars have emerged, spraying immediately with Oil-mineral (A21 i), potassium permanganate (A22), or with sprays made out of Pyrethrum (B 1.6), Tobacco (B 1.17), the Rain tree (B 2.19), Cork bush (B 2.20) and Tephrosia (B 2.26) should prevent the appearance of unsightly holes in the leaves.

If a few holes are not a problem, liberally dusting the caterpillars and their food with lime is likely to be the most convenient alternative but the lime should be washed off before the goods are marketed.

**Pupae:** Ash (A2 ix, x, xi); Clay etc. (A10 i, ii, v); Hens etc. (A16); and any of the insecticides listed in B6 particularly those already mentioned for rapidly controlling the caterpillars. Thoroughly spray the underside of the leaves to clear the pupae.



**Checks**

Ensure an adequate habitat for owls, nightjars, bats and other night creatures that predate on the moth; and for birds that will feed on the eggs, pupae and caterpillars during the day. Many other insects feed on the eggs and larva of this pest, so encourage as wide a range as possible of insect life. Pay adequate attention to rotations to break the continuity in the growing of crucifers. A break of 6–8 weeks in which no crucifers at all were grown in a market garden reduced pest numbers considerably.

**FRUIT FLY: (TEPHRIDAE)****Host plants**

Apples, apricots, avocados, citrus, figs, grapes, guavas, mangoes, olives, pawpaws, peaches, pears, plums and quinces.

**Damage**

The maggots of the fruit fly eat into the flesh of the fruit and promote rotting.

**Life cycle**

In summer the female cuts a hole in the skin of the fruit and inserts a number of eggs. The eggs hatch in 2–4 days. The maggots feed on the tissues of the fruit for about 2 weeks. During this period the fruit is likely to rot and drop to the ground. At the end of this period the maggots are fully grown (10mm long) and leave the fruit to burrow into the soil and pupate. The adult flies emerge after 2–3 weeks. In winter the flies tend to congregate on the underside of leaves of certain trees like orange and loquat. The honeydew secreted by scale insects and aphids attracts them.

The female fly also lays her eggs in the growing tips of the cabbage tree. The maggots then tunnel down the branches keeping just beneath the bark, causing a slimy mess under the bark and die-back of the shoot.

**Identification**

There are several species of fruit fly. This small fly can be recognised by its shiny eyes and striped wings. The body is light brown with darker markings: stripes or dots. It holds its wings at a wide angle to the body and appears to trail the back edge of them as it rests or walks. The maggots are small, legless and creamy white. The first signs of infestation are tiny brown punctures in the skin of the fruit from which clear gum exudes.

**RECOMMENDED REMEDIES**

The eggs and maggots, being buried in the fruit, are difficult to treat. The fly and pupa stages are the most vulnerable.

**Fly:** the following have been mentioned specifically as controls of the fly: Manure etc. (A17 ii); Finger millet (B 1.8); Flies (C9); Various insects (C17 iii).

Spraying with residues of finger millet soaked in water has been found effective against fruit fly.



FRUIT FLY

Susceptible fruit should be picked early, when green if possible, to ripen in storage.

The flies can be repelled by spraying the fruit with the plant remedies listed in B6.

Timely and routine spraying of the fruit with the plant remedies listed in B5 will help to reduce numbers of flies and eggs. Those identified as treatments for flies in general may be worth trying first: Garlic (B 1.2); Chilli (B 1.4); Pyrethrum (B 1.6); Tomato (B 1.12); Feverfew (B 1.14); Oleander (B 1.16); Tobacco (B 1.17); Goat weed (B 2.2); Apple of Peru (B 2.21); Mexican marigold (B 2.25) and Woman's tongue (B 4.2).

*A strong recommendation is to keep numbers of aphids and scale down to a minimum as fruit fly are attracted by the honeydew secreted by these insects.* This can be done most economically by encouraging ladybirds and bird life into the orchard.

**Maggots and Pupae:** fallen fruit containing the maggots should be collected, buried deeply (50–100cm), composted or fed to livestock immediately. Grease bands have been recommended to prevent the larvae from crawling up and down the tree trunk. Hens etc. (A 16) will dig maggots out of the fruit and pupae from the soil. Hens etc. and orchards go well together. They can be assisted in their task by disturbing the soil under the fruit trees: Soil (A26 ii).

The Cabbage tree is a very useful trap crop for the pest. The infested branches should be cut off and destroyed as soon as the leaves begin to wilt under the maggot attack.

The pupae are said to pass the winter in loquats and particular care should be taken to remove the fallen fruit from under this tree.

**Checks**

For this pest particularly it is best to attack from as many possible angles at once. Hence spray, trap or repel flies, harvest early, collect and destroy fallen fruit and also get the hens to eat the pupae hiding in the soil under the trees. Are there enough Cabbage trees as decoys in the area? Are you encouraging bird life to come into the orchard with bird baths and so on?

**GRASSHOPPERS: (ACRIDIDAE)****Host plants**

Garden flowers, herbs, shrubs and vegetables.

**Damage**

The leaves and even the soft stems of young shoots will be eaten, impairing growth and destroying seedlings. They do the most damage to gardens in early summer before the surrounding bush can offer them sufficient green vegetation.

**Life cycle**

The adults lay their eggs in the soil, just below the surface. In some species of grasshopper the eggs may lie in the ground all through



GRASSHOPPERS

winter, hatching when the weather warms up. In this way the insects can survive prolonged dry periods during which food may be scarce.

#### Identification

Most grasshoppers can jump well and can fly strongly. Their hindwings are often brightly coloured. Their hindlegs are powerfully developed and many species have sharp spines on them. The 30mm long, brown, short-horned grasshopper is perhaps the most common grasshopper in gardens but locusts are included in the *Acrididae* family.

#### RECOMMENDED REMEDIES

Planting members of the Marigold (*Tagetes*) family as companion plants is said to give some protection against grasshopper attack.

**Adults:** the following remedies have been mentioned in the literature: Ash (A2 i, ii); Biological (A4 i, ii); Hand picking (A15 iii); Hens etc. (A16); Manure (A17 i); Mulch (A19 i); Tomato (B 1.12); Custard apple (B 2.4); Catunaregam (B 2.9); Leadwood (B 4.9); African sandalwood (B 7.2); Beetles etc. (C3); Various insects (C17 i).

Of these using hens, light traps (Various insects C17 i) and hand picking or, preferably, nets (C3) are simple and effective.

In addition some success in reducing adult numbers has been claimed by spraying and dusting with the stronger plant remedies like Pyrethrum (B 1.6), Tobacco (B 1.17), Rain tree (B 2.19), Cork bush (B 2.20) and Tephrosia (B 2.26).

**Eggs in the soil:** since the grasshopper likes to lay its eggs in bare soil, ensuring dense living ground cover (crops and grassland) and Mulching (A19 i) will significantly reduce pest numbers. Hens etc. (A16) will help by digging up any eggs laid in patches of bare soil; and strong sprays from Pyrethrum (B 1.6), Tobacco (B 1.17), the Rain tree (B 2.19), Cork bush (B 2.20) and Tephrosia (B 2.26), should prove useful when watered thoroughly into the ground.

#### Checks

Avoid destroying all blister beetles, their eggs and larvae as the larvae eat the eggs of grasshoppers. Check on the amount of vegetative cover in the garden and in the surrounding bush. Overgrazing and burning of the grassland, and particularly areas of ploughed but unplanted land, provide bare areas which are ideal breeding areas for the hoppers. Encourage an abundant bird life.

#### MAIZE STALK BORER: (*BUSSEOLA FUSCA*)

##### Host Plants

Maize, millet, sorghum and grasses.

##### Damage

The insect attacks members of the grass family. The leaves are perforated and the caterpillar bores into the stem causing loss of yield. The plant may dry up and die if the infestation is severe or the plant is weak.



MAIZE STALK BORER

#### Life cycle

The mature insect is a night-flying moth with a wingspan of about 35mm. The female moth lays her eggs in a cluster between the leaf sheaths and the stalk. The eggs hatch from 7–9 days after being laid. The young larvae make their way to the top of the plant and feed between the unopened leaves, perforating them. As they increase in size they become over-crowded. They then migrate to other maize plants and bore into the stems. A full-grown larva measures about 30mm. It pupates within the stem. The adult moth emerges after 2–3 weeks. Maize grown under the natural rainfall is attacked by 2 main broods of the moth. The first brood emerges shortly after the first rains in November/December, and causes a lot of damage. The second brood appears in February/March, tends to attack the tassels and grain but the damage is much less severe. This brood passes the winter as a larva within the maize plants and gives rise to the spring brood of moths. Many insects predate on the moth, egg, caterpillar and larval stages of this pest including ants, ladybirds, parasitic wasps, spiders, birds and many others.

#### Identification

The moth can be seen hovering around the maize plants on moonlit nights. Its forewings are reddish brown and the hindwings yellowish brown. The eggs can be seen clustered between the leaf sheath of the plant and the stalk. The grubs are dark coloured at first and can be seen by splitting the sheath downwards until they are exposed. Blobs of rotting vegetation are excreted by the grub. Later the grubs become pale in colour and exit holes can be seen in the lower stem.

#### RECOMMENDED REMEDIES

Stalk or stem borers can be effectively controlled by composting all infected plant residues or feeding them to the cattle. The compost/manure can then be returned to the field.

The pest's life cycle can also be broken by practising rotations, with wide gaps between members of the grass family.

Intercropping, mixed cropping, strip cropping and Permaculture designs reduce moth attack because the moth apparently recognises the host plant by its silhouette (outline). Thus tall intercrops such as tomato, sunflower, climbing beans, and so on, give better protection than short intercrops.

Another tactic which is said to reduce the incidence of attack in a field is to plant a susceptible maize hybrid, with the first rains, around the edges of the field to act as a trap crop, which is then cut for fodder. These methods, however, are far less satisfactory than composting or rotational practices mentioned previously.

Maize grown for silage which is planted at the start of the rains can act as a decoy or trap crop. The caterpillars or larvae are then killed off when the maize is turned into silage. Similarly, silage maize can be planted in mid-January to divert the second brood from the grain crop.



As the moths are most active during the full moon and are attracted to plants 3-4 weeks old, the best time to plant is between two full moons. It is also possible to plant at a higher density than optimal and chop out the infected plants later for fodder. Full moon is also the best time to catch the moths in light traps as there are more of them about at that time.

Other recommended planting strategies are either to plant 3-4 weeks after the start of the rains when moth activity has lessened or to plant 1 month before the start of the rains using irrigation.

One farmer reports that he controls stalk borer in sorghum by cutting off and destroying the central shoot of the plant (containing the stalk borer) before the plant flowers. Lateral shoots grow to replace the main one.

**Moths:** Aromatic plants (A1); Biological (A4 i); Gum tree (B 1.9); Light traps: Various insects (C17); Repellent sprays (B6).

The bodies of the moths caught in the light traps: Various insects (C17) can be used as a Biological (A4 i) spray.

One farmer reports that spraying plants (especially the tops and bottoms) with powdered gum tree leaves proved an effective repellent to the moth particularly when the days were sunny.

**Eggs:** Ash (A2 i, ix, xi); Clay etc. (A10 i, v); Hand picking (A15 ii); Manure etc. (A17 ix); Oil-mineral (A21 i); Pyrethrum (B 1.6); Tobacco (B 1.17); Rain tree (B 2.19); Cork bush (B 2.20); Tephrosia (B 2.26). Make sure to spray particularly around the leaf sheaths.

**Caterpillars:** Various preparations can be poured down the funnel of maize plants when they are about knee high or when first signs of leaf damage occur. Ash (A2 i) and Soil (A26 i) are mentioned particularly in the literature; Tephrosia (B 2.26) gave effective kills in a field trial; Tobacco (B 1.17) performed well but recommended concentrations should not be exceeded if plant damage is to be avoided; Thorn apple (B 2.11), Lantana (B 2.15) and Red sunflower (B 4.18) were only partially effective. Others which have shown promise are Pyrethrum (B 1.6), Rain tree (B 2.19) and the Cork bush (B 2.20).

The above liquid preparations can also be sprayed on the leaves to kill eggs and the newly-hatched caterpillars.

These preparations are most effective when the stalk borer caterpillar is small and resting either on the leaf or in the leaf sheath before boring into the stem.

The crop should be treated at least twice and preferably three times at 7-day intervals once the infestation is evident.

**Pupae:** pupae can best be destroyed by composting the crop residues or feeding them to livestock. Alternatively, when the residues are to be retained as mulch the method given in Sun (A29 i) should be followed.

By far the simplest method of controlling maize stalk borer are rotations and removing all the residue from the land and turning it into compost or feeding it to the cattle. For the small number of plants still attacked, pouring soil into the funnel gives very effective control at no cost and little effort.

### Checks

Ensure that the residues of all host plants (sorghum, maize and millets) are free of borer larvae or are turned into manure or compost. Encourage surrounding growers to follow similar practices. And, of course, do everything possible to reinstate a wide variety of bird and insect life. Ants are fond of the pupae and will find them by going down the borers tunnels in the maize stem.

### NEMATODES: (NEMATODA)

#### Host Plants

Apples, avocados, citrus, figs, garden flowers, grapes, pawpaws, peaches, pears, strawberries, vegetables — particularly: bambara nut, beetroot, brinjals, beans, carrots, celery, cowpeas, cucumbers, cucurbits, gourds, potatoes, leeks, lettuce, melons, okra, onions, parsley, peas, pumpkin, sorghum, squash, sugar bean, sweet pepper, swiss chard, tomatoes, tsetse; and cotton, millets, sunflowers, tobacco.

#### Damage

Nematodes attack a very wide range of plants causing poor growth, wilting, poor development of leaves and fruit, yellowing of leaves and other nutrient deficiency symptoms, with potentially dramatic loss of yield. The ends of branches can die in tree crops. Nematodes which leave open wounds in the root tissue encourage the occurrence of plant diseases. However, the majority of nematodes are essential to good soil structure as they break down organic matter into humus.

#### Life cycle

Nematodes live mainly in the soil. They are worm-like creatures which feed on soil bacteria, fungi, algae, other nematodes and some feed on the roots of plants. Root-knot nematodes are microscopically small. The young root-knot nematode burrows into the root tissue of the plant where it matures and lays eggs, causing localised swellings on the roots. The young emerge from the eggs and move into the soil in search of another suitable plant. The life cycle from egg to adult is about 4 weeks and the juvenile nematode can survive for a couple of years without finding a suitable plant host. Other nematodes do not live in the roots but live in the soil, lay their eggs there and feed on the roots without entering them. Some nematodes are beneficial organisms as they keep each other in check and parasitise the larvae of many insects including armyworm, beetles, bollworm and leaf miner.

#### Identification

Most plant-parasitic nematodes cannot be seen with the naked eye. The presence of these parasites must therefore be identified from the damage to the plant and its roots. Apart from the above-ground evidence of damage as described, which is common to all nematode attack, the cause should be confirmed by examination of the plant roots. The root-knot nematode causes localised swellings on the roots, lesion and ring nematodes cause dead spots to appear around their feeding points and

nematode root knot damage



NEMATODES

daggernematodes cause swelling and distortion of the roots on plants.

### RECOMMENDED REMEDIES

Nematodes are always in the soil but in balanced, harmless numbers. Numbers of root-knot and other crop-damaging nematodes increase dramatically as soon as the soil is ploughed and/or organic matter decreases. The continuous growing of host crops adds to the problem by ensuring adequate food supplies for the inflated nematode population.

The most permanent solution is therefore to minimise soil disturbance, to ensure as high a level of soil organic matter as possible and to include nematicidal plants in the rotation.

**Raising soil organic matter:** soil organic matter can be raised by no or minimal digging, by mulching, and by including crops in the rotation which add much organic matter to the soil through their dense root networks (preferably) and/or prolific above-ground vegetation which can be used as residue. For example: millets, maize, many types of beans both bush and creeping, sunnhemp, lucerne, silver leaf and other fodder grasses; also direct additions of Compost (A12 ii), Manure etc. (A17 viii) and Mulch (A19 i) to the soil surface.

Sunnhemp has the potential to not only improve soil organic matter but also to act as a trap crop for nematodes by preventing the female from maturing.

**Plants with nematicidal properties:** garlic, cassava, leeks, mustard, onions, shallots, pawpaw, African and French marigold, Mexican poppy, Mexican marigold and spear grass.

**Rotations:** apart from the nematicidal plants already mentioned which can also be introduced into rotations, several grasses are known to be non-hosts or very poor hosts to root-knot nematode. These are *Eragrostis curvula* (Weeping Love grass), *Chloris gayana* (Katambora Rhodes grass), *Panicum coloratum* (Bambatsi Panicum), *Cynodon dactylon* (Couch), *Paspalum notatum* (Paraguay grass) and *Panicum maximum* (Sabi Panicum). When grown in extended rotation (several years of grass included) nematode numbers are decreased significantly.

Among the crops said to be poor hosts or resistant to nematodes are garlic, leek, onion, early planted green maize and sweet corn, and shallots. Those said to be able to tolerate attack are: broccoli, brussels sprouts, cabbage, cauliflower, kale, rape, radish, leaf mustard, turnip, sweet potato and chillies.

In addition, it is said that traditional vegetables are usually tolerant or even resistant to indigenous pests and diseases including root-knot nematodes, and could be included in the rotation to advantage; whereas the ones of European origin are more susceptible.

Cropping strategies recommended by experts are:

- Plant susceptible intercrops such as pumpkin, okra, cowpea and bambara nut into a field of maize, sorghum or millet.
- Grow winter (irrigated) crops of European-type vegetables in

strict rotation with crucifers or alliums. Grow green maize on this same land during the early summer to reduce the nematode population.

- Sterilise seed-beds by burning wood or other materials on the soil and change/rotate the position of the seedbed, selecting previously uncultivated ground wherever possible.
- Flood seedbeds and plots with standing water for a minimum of 4 weeks prior to planting.
- Plant known resistant plants or varieties and nematode-free roots, tubers, seedlings and cuttings.
- Plant susceptible crops in winter during periods when the soil temperatures are below 15°C as the nematodes are relatively inactive at lower soil temperatures.
- Remove infested roots immediately after harvest, dry and burn them. The tops can be composted but don't use the infected roots as nematodes can survive in the cooler parts of the heap.
- Add copious amounts of organic matter to the soil like compost, mulches, legumes (particularly sunnhemp), manures and through the root systems of uninfected plants left to decompose in situ. High levels of organic matter which encourages a healthy plant and a good pest/predator balance will control nematode numbers.

**Others:** Citrus trees are said to be protected from nematode attack by planting legumes underneath (like cowpea) or nearby (acacia trees).

### Emergency tactics

Other methods specifically recommended in the literature are: Ash (A2 iv, vii); Burning (A9); Sugar (A27); Garlic (B 1.2); Pawpaw (B 1.5); Tomato (B 1.12); Cassava (B 1.13); Syringa (B 1.15); Goat weed (B 2.2); Sugar apple (B 2.3); Mexican poppy (B 2.5); Thorn apple (B 2.11); Castor oil (B 2.24); Chinese lantern (B 4.10); Spear grass (B 4.12).

Sugar is particularly useful for destroying root-knot nematodes in seed potatoes before planting by soaking them in sugar water and then rinsing. It may be a practical method for potting and small garden plots but not for large areas.

### Checks

Check that soil disturbance is minimal; that existing soil humus levels are high; that there is an adequate programme to maintain organic matter levels; and that nematicidal or resistant crops and soil-improving plants are part of the rotation.

### RED SPIDER MITE: (*TETRANYCHUS SPP.*)

#### Host plants

Apples, apricots, beans, brinjals, carnations, carrots, citrus, cotton, crucifers, cucurbits, deciduous fruits, grapes, nectarines, peaches, pears, plums, potatoes, roses, soyabeans, strawberries, sweet potato, taro, tobacco, tomatoes, and others.





RED SPIDER MITE

#### Damage

There are several kinds of mites of different colour but the red spider mite is probably the most widespread. These insects suck the sap from plants causing retarded growth, leaf-fall or death.

Feeding initially causes yellow spots to form on the leaves which gradually spread until the whole leaf is yellow and drops prematurely. They do not like wet conditions and outbreaks tend to occur in warm, dry conditions.

#### Life cycle

The female lays her eggs on the underside of plant leaves protected by the fine web spun by the insect. She lays about 15 eggs a day for about 3 weeks. In hot weather it takes only 10 days from egg to adult. This rate of reproduction can result in a very rapid build-up in numbers. It is said that they can remain dormant in the soil or on plant residues for a long time before colonising plants. Ants, lacewing larvae, ladybirds, praying mantis and other mites predate on the red spider mite.

#### Identification

The red spider mite is a small oval insect 0.5–0.7mm in length. It has four pairs of legs and is reddish brown in colour. The first sign of infestation often is the presence of yellow spots on the upper surface of the leaves. In heavy infestations they can be clearly seen massing on the underside of the leaves of the host plant, the crop having a reddish colour. They spin silken webs over themselves, hence the name red spider mite.

#### RECOMMENDED REMEDIES

Rotations are particularly important in the control of this insect. The situation is currently particularly serious on lands planted to cotton, crucifers and tomatoes without due attention to rotating with non-host crops. Overuse of modern agricultural chemicals has built up a large resistant population while effectively removing most of the predators of the red spider mite. These predators such as ladybirds/bugs, mantises, lacewings and ants should be allowed to build up to re-instate the natural pest/predator balance.

Maintaining the humidity in the crop as high as possible (A4 iii) will repress numbers of spider mite as they do not like damp conditions. This can be achieved by close plant spacings, mulching, spraying the plant leaves with a water jet and frequent irrigations.

Increasing the plant diversity will also improve protection. A hedge of perennial pigeon pea around the field is recommended as a breeding site for other mites which predate on the red spider mite, and increased humidity in the garden will further reduce pest numbers.

Interplanting with garlic, basil, onions and other aromatic plants is said to give some protection.

#### Supplementary measures

The following measures are said to control the insect at all stages in the

life cycle: Ash (A2 i, ix, xi); Biological (A4 iii); Clay etc. (A10 i, ii, iv, v, vi); Glues (A13); Hand picking (A15 ii); Manure etc. (A17 iii); Milk (A18 i); Oil-Mineral (A21 i) — see warning; Potassium permanganate (A22); Soap solutions (A25 i, ii, iii); Sulphur (A28); Water (A31 i); Onion (B 1.1); Garlic (B 1.2); Pyrethrum (B 1.6); Tomato (B 1.12); Syringa (B 1.15); Tobacco (B 1.17); Sweet basil (B 1.18); Blackjack (B 2.8); Euphorbia (B 2.13); Rain tree (B 2.19); Cork bush (B 2.20); Wild basil (B 2.23); Castor oil (B 2.24); Plant Mixtures (B 3.3, 3.6).

In addition, any of the Insecticidal plants (B5) should be tested. The effectiveness of repellants on red spider mite (B6) is as yet unknown; but the ones mentioned as useful in interplanting: basil, chives, garlic and onions, should be appropriate.

#### Checks

Ensure plant diversity is sufficient through practising rotations, mixed, inter, strip and trap cropping and Permaculture designs. Consider using cultivars known to be resistant to red spider mite. Check that spraying is done as infrequently as possible and spot spray to avoid killing off predators.

#### SCALE INSECTS: (COCCIDAE)

#### Host Plants

Apples, apricots, avocados, citrus, coffee, deciduous fruits, figs, granadillas, grapes, mangoes, mulberries, olives, peaches, pears, plums, quinces; protea, roses, sisal and many other orchard trees and garden plants.

#### Damage

Scale insects suck the sap of plants and may have a toxic effect causing loss of health or death of the plant. They are mostly a pest of orchard trees and other woody plants. Palms, ferns, herbaceous plants and sisal are also attacked.

#### Life cycle

Armoured scales protect themselves by secreting a waxy shield which grows in size as the insect matures. The waxy substance is further protected by a hard surface made up of their own moulted skins. The male eventually emerges from the shield and flies off. The females remain under the waxy cover and produce their young beneath it, usually as eggs (a few species produce young directly). The young larvae hatch, emerge and enjoy a brief, active period during which they seek new places to settle. They then moult, becoming adults, then secrete the waxy protection. Soft scales do not have a hard cover, the soft coating being made of solidified watery wax. Naked scale varieties do not secrete wax but are protected by their own horny backs. The natural controls for these insects are birds, parasitic wasps and larvae. Beetles of the Coleoptera family have been reported eating scale insects.

wax scale — acacias



SCALE INSECTS

**Identification**

Armoured scale insects commonly seen grouped on the leaves of mango trees are small, white and elongated. Blemishes on the leaf caused by toxic reaction can clearly be seen. The adults of the species look like small, white, grey or reddish-brown shells glued to the branches or leaves of the plant. Lines of ants can be seen on their way to collect the honeydew secreted by soft scales. Armoured scales do not secrete honeydew.

**RECOMMENDED REMEDIES**

The incidence of scale can be reduced by pruning heavily infested branches in winter. Controlling the ant population which feeds off the honeydew secreted by soft scales will allow predators to do their job — see *Ants*.

The following are specifically recommended in the literature; Ash (A2 ix, xi); Baking soda (A3); Clay etc. (A10 ii); Oil (A21 i, ii, iii)—see warning; Soap solutions (A25 ii); Sulphur (A28); Onion (B 1.1); Tobacco (B 1.17); Catunoregam (B 2.9).

In addition, any of the Plant Mixtures (B3) and Insecticidal plants (B5) should be tested.

The effectiveness of Repellants (B6) is as yet unknown.

For scale on woody branches or plants with naturally oily leaves (mango, sisal), brushing or spraying with vegetable or mineral oils is very effective. Water-mixable mineral oils will give equally good results on vegetation with non-oily leaves.

**Checks**

Scale is normally kept in check by bird life, parasitic enemies and by ensuring healthy plant growth. Therefore when plants are infested with scale, pay particular attention to those factors ensuring healthy plant growth: soil fertility, watering and seedling vigour. Provide a suitable habitat for birds: fruit trees, shrubs and permanently-filled bird baths.

**SLUGS AND SNAILS: (PHYLUM MOLLUSCA)****Host plants**

Almost any plant except some of the aromatic bitter ones.

**Damage**

These small animals eat the leaves off plants, produce unsightly holes in fruit and can completely destroy seedlings leaving only a short stem protruding from the earth.

**Life cycle**

Slugs and snails favour damp places. Their round, white eggs are laid in the soil, under leaves or under any object in contact with the soil.

**Identification**

Slugs and snails are soft-bodied animals, snails with shells and slugs



SLUGS AND SNAILS

without. They grow up to about 7–8cm long and leave slimy, silvery trails behind them, which are detectable for days afterwards. Large snails with conch-shaped shells can be found sometimes. They are said to prey on their smaller cousins.

**RECOMMENDED REMEDIES**

Remedies in the literature recommended for controlling both slugs and snails are: Ash (A2 ii); Builder's lime (A7 i); Clay etc. (A10 iii); Coffee etc. (A11 i); Hand picking (A15 iv); Hens etc. (A16); Mulch (A19 i, ii); Salt (A23 iii); Smoke etc. (A24 i); Soap solutions (A25 ii) with paraffin added; Tea (B 1.3); Chilli (B 1.4); Finger millet (B 1.8); Tobacco (B 1.17); Desert date (B 2.6); Barbados nut (B 2.14); Wild tobacco (B 2.22); Slugs and snails (C14).

Those mentioned for slugs only are: Biological (A4i), Clay etc. (A10 i) and under Caterpillars etc. (C7). Spinach planted in lines is said to repel slugs.

Slugs and snails can survive only in permanently wet conditions. Their numbers can be controlled by allowing infested areas to dry out thoroughly over winter.

Dusting around the plants with finely ground-up tobacco leaf scrap or agricultural lime are very effective methods but need repeating often. Too much lime will make the soil too alkaline.

Dusting floors with builder's lime (avoid the plants) is an efficient way of removing slugs and snails from nurseries but having the legs of nursery tables sitting in tins full of water does the trick too.

Hand picking is slow, labour-intensive and needs repeating often; wet days are the best time. Similar problems are experienced with traps. These methods reduce adult numbers but do not get to the root of the problem. New generations keep emerging.

Allowing free range to bantams is thoroughly recommended as these useful creatures prefer insects for food and do little damage to plants. On the other hand, hens and ducks love greens and should be fenced into the orchards where they can do no damage and moved on to the garden to clean up unplanted areas between crops in the rotation.

A useful tactic is to delay planting out seedlings until they are large enough and the growth sufficiently vigorous to withstand attack.

**Checks**

Keep the ground clear of non-essential materials lying on the soil which provide perfect hiding places for slugs and snails and their eggs: boards, bricks, sacks, tins, pots, stones and so on, unless they have been deliberately placed there as traps. Small animals such as toads, moles, mice, hedgehogs and large predatory snails, as well as many insects are said to feed off both the adults and the eggs.

Birds are essential to control numbers of these troublesome pests.



STINK OR SHIELD BUGS — ANTESTIA (ANTESTIOPSIS SPP.), BAGRADA. (BAGRADA HILARIS) AND GREEN STINKBUG (NEZARA VIRIDULA)

*Host plants*

A wide variety of plants are attacked. Antestia bugs attack coffee and orchard trees and protea. Bagrada bugs prefer pawpaws, crucifers and garden flowers, while their nymphs attack a wide range of food plants. Green stinkbugs are known to be able to feed on about 200 different plants including grapes, millets, soyabeans and tobacco.

*Damage*

The bugs suck the sap from the plants causing wilting and death of the leaves, growing tips or, in the case of heavy infestations, the whole plant may die.

*Life cycle*

The life cycle of the antestia bug will serve as an example. The female lays her eggs on the twigs and leaves of the chosen food plant in clusters of 10–14. She lays several batches and then dies. The eggs are about 1.5mm long, pearly white and shaped like short, stout barrels with lids on the top through which the young appear. The eggs darken and hatch in 10–14 days into nymphs which are black and wingless but look similar to the adult bug. They scatter and feed on sap for the next 12 days, then rest and moult. Their colour changes to bright yellow and black spots. As they continue to feed they shed their skin another 4 times whereupon the adult stage is reached. The whole life cycle from egg to death of the adult takes about 100 days, giving three generations per year. The eggs are parasitised by a tiny black wasp. The green stinkbug cycle is identical to the antestia bug but the female bagrada bug lays its small, barrel-shaped, greenish-white (at first) to pink mottled (later) eggs in the soil near to the host plant or on the leaves. They hatch in about 5–7 days. The subsequent stages, however, are the same as for the other two bugs.

*Identification*

There are well over 4,000 known species of shield or stink bug. Their bodies are flattened and shield-shaped. If roughly handled they give off a strong, unpleasant odour. They do not have biting jaws like beetles but a slender tubular probe which is carried folded back against the underside of their bodies. The Antestia bug is about 10mm, usually yellowish-green in colour with orange and black markings but can be green with red, yellow and blue markings. Bagrada bugs are often seen in pairs. They are about 6mm long, black, with a few bright yellow or orange markings, the nymphs being bright red. The so-called green stinkbugs can be green or brownish-green and are about 15mm long.

**RECOMMENDED REMEDIES**

**Large beetles:** the larger types of beetles, not usually appearing in

adult bagrada bug



STINK OR SHIELD BUGS

great numbers together, are best treated by simple methods: Aromatic (A1) to repel them; Biological (A4 i); Hand picking (A15 i); Hens etc. (A16); Beetles (C3).

**Small Beetles:** Aromatic (A1); Ash (A2 ix, xi); Biological (A4 i); Clay etc. (A10 i, ii, ix) — particularly lime; Hens etc. (A16); Manure (A17 i); Soap solutions (A25 ii); Tea (B 1.3); Pawpaw (B 1.5); Syringa (B 1.15); Mexican poppy (B 2.5); Lantana (B 2.15); God's tobacco (B 2.18); Castor oil (B 2.24); and the Tung tree (B 4.3) have been specifically identified in the literature.

A solution of green bar soap in water has been found effective against Bagrada bugs.

Insecticides recommended in the literature as effective against various beetles are: Garlic (B 1.2); Syringa (B 1.15); Oleander (B 1.16); African marigold (B 1.20); Goat weed (B 2.2); Mexican poppy (B 2.5); Blackjack (B 2.8); Wormseed (B 2.10).

\* Also worth trying are any of the insecticidal plants mentioned in B5 particularly sprays from Pyrethrum (B 1.6), Tobacco (B 1.17), the Rain tree (B 2.19), Cork bush (B 2.20) and Tephrosia (B 2.26).

Routine spraying with mixtures of repellent plants from B6 may prove a useful preventive measure.

Trials with different coloured containers as traps (C4) and light traps: Various insects (C17 i) may yield useful results.

**Nymphs on leaves:** Ash (A2 i, ix, x, xi); Clay etc. (A10 i, ii, v); Hens etc. (A16); any of the insecticides listed in B6 particularly those \* mentioned previously for controlling the adults.

**Eggs on the leaf:** Ash (A2 ix, xi); Clay etc. (A10 i, v); Hand picking (A15 ii); Oil-mineral (A21 i); any insecticides listed in B6 particularly those \* mentioned previously for controlling the adults.

**Eggs in the soil:** Ash (A2 ii); Hens etc. (A16); and strong sprays from Pyrethrum (B 1.6), Tobacco (B 1.17), the Rain tree (B 2.19), Cork bush (B 2.20) and Tephrosia (B 2.26), should prove useful when watered thoroughly into the soil.

*Checks*

Ensure an adequate habitat for bird life that predate on the beetles and that there is a wide variety of insects to predate on the eggs and nymphs. Tree, hedge and plant barriers will often prevent the beetles from flying into the field. For the same reason intercropped or mixed cropped plants suffer less.

**TERMITES: (TERMITIDAE)**

*Host plants*

Fungus-growing termites feed on dead grasses, young woody plants, the old bark off mature trees and any dead organic material. They will consume live plant material when no other food is available, particularly groundnuts, maize and millets. Harvester termites will collect green material stripping the veld of grass and the leaves from shrubs.



TERMITES

### Damage

Hundreds of species of termites exist in Africa alone, of which a few are considered important because they damage plants. Of these few, the fungus-growing termites are probably the most troublesome. Fungus-growing termites are known to attack the roots of young shrubs and trees during the first two years after they have been planted, causing retarded growth or death. They do no harm, however, when feeding on the dead bark of established trees. These termites are also known to attack field crops in semi-arid regions but usually only do so when plants are already weak (for example, wilting) or when no other food is available to them (for example, no other sources of organic matter such as mulch and soil humus).

### Life cycle

Most termites that damage crops and trees live in nest systems constructed well below ground. The fungus-growing termites, which cause the most damage to trees and crops, live on wood and other plant material (usually dead). The collected plant material is taken back to the nest for the construction of fungus gardens, which are visculated with a special species of fungus (*Termitomyces*). This fungus breaks down the plant cellulose and enables the termites to use the plant material as food. The queen and king live in a thick-walled, mud cell in the centre of the termitary, usually a high mound or vertical shaft. Workers carry away the eggs as she lays them. The queen and king never move away from the cell; but each year during the rainy season winged termites (potential kings and queens) appear in the colony and fly off. They then pair off and form new colonies. Termite numbers are controlled particularly by ants, birds, reptiles, amphibians and mammals.

### Identification

The presence of the fungus-growing termites can be detected from the mud tunnels they build en route to and around the material to be attacked. The fungus-growing termites are about 10mm long, white-bodied with brown heads and the soldiers have prominent pincers.

### RECOMMENDED REMEDIES

Soil disturbance upsets the pest/predator balance among termites and favours those which are more likely to attack plants.

Termite attack on crops has been observed to be related to the loss of organic matter in the soil. Therefore all means should be taken to improve soil organic matter content.

Generally termites will attack only weakened plants under stress, particularly those under moisture stress thus moisture conservation techniques (conservation-tillage), which reduce surface run-off, should be applied at all times and the encouragement of healthy plants through compost and manure additions.

When there is no other food available to them as is often the case on arable land, termites have no option but to attack green plants. If mulch,

dead-grass, compost, manure or litter is present they will feed off that in preference to green material. So avoid having bare, dry, disturbed, residue-free soil around the plants.

Damage to crops can be lessened by harvesting the crop as soon as possible and drying it off the land.

Certain trees are resistant to termite attack: *Acacia* sp, *Cassia* sp, *Colophospermum mopane*, *Markhamia* sp, *Meliaceae* sp (*Mahogany* family), *Pterocarpus angolensis* and *Terminalia* spp.

Other strategies to protect young trees are: avoid seedling containers made of materials attractive to the termites like bark, paper or cardboard, take only the base of the plastic bag off the seedling plant and leave the top of the bag jutting above ground level and plant just after a heavy storm when the soil is moist.

### Supplementary measures

Supplementary measures specifically recommended in the literature are: Ash (A2 v, vi); Biological (A4 i); Mulch (A19 i); Smoke etc. (A24 iii); Water (A31 i); Garlic (B 1.2); Tea (B 1.3); Pawpaw (B 1.5); Syringa (B 1.15); Tobacco (B 1.17); Sweet basil (B 1.18); American aloe (B 2.1); Mexican poppy (B 2.5); Blackjack (B 2.8); Catunaregam (B 2.9); Wormseed (B 2.10); Rubber hedge (B 2.13); Lippia (B 2.17); Wild basil (B 2.23); Castor oil (B 2.24); Mexican marigold (B 2.25); *Albizia anthelmintica* (B 4.1); Swallow wort (B 4.6); Finger grass (B 4.11); Red mahogany (B 4.13); Sodom apple (B 1.16).

Liquid from blackjack seeds and tobacco leaves has been found very effective locally as has the use of smoke and lemon-flavoured herbs such as lemon grass, vetiver grass and penny royal.

The leaves of *Aloe graminicola* crushed and the leaves or berries of *Oximum* applied to planting holes or watered into the soil around plants are said to protect plants from termite attack. In addition the leaves of *Cassia siamea* can be used as a protective mulch; and extracts from the leaves of *Commifora africana* (Corkwood) and from the roots of *Adenia gummifera* (Monkey rope) are said to have potential in termite control.

Castor oil cake soaked for 2 weeks in the latex from aloes can give eight months' protection to trees which have had the mixture painted around the base of their trunks.

Diesel, paraffin and used engine oil have also proved effective in repelling termites from their termitaries and from houses and earth dams; and have been painted onto the trunks of trees or mixed with the soil at planting time to protect them against termite attack. These products should not be used in the garden as they are harmful to plant and soil life.

### Checks

Make sure that the soil is being disturbed as little as possible, that there is plenty of humus in the soil and dead material to feed the termites, and that the soil is kept as moist as possible. In short, avoid having bare, dry, disturbed, organic-deficient, residue-free soil. Allow time for the pest/predator balance to be restored.



**THRIPS: (THYSANOPTERA)***Host plants*

Apples, avocados, beans, brinjals, citrus, crucifers, cucurbits, garlic, grapes, green peppers, guavas, mangoes, onions, peas, tomatoes and many garden plants.

*Damage*

Due to their feeding activities these very small sap-sucking insects cause scars and blemishes on leaves, fruit and pods of a wide range of plants. Onion thrips attacks many vegetables and may cause plants to wilt and die. Onion thrips is a vector of tomato spotted wilt virus.

*Life cycle*

The adults are very small (1–3mm long), yellowish, slender insects having four oar-like wings fringed with long hairs. They breed throughout the year although they favour warm, dry weather. The female lays her eggs in the leaf tissue and they hatch in about 4 days. In warm weather the cycle from egg to adult is about 16 days. The insect tends to hide in the leaf sheaths or under the leaves. Some species pupate in the soil.

*Identification*

Few of these insects exceed 2mm in length and nearly all of the 5,000 or more species are black, brown or yellow. Signs of damage to plants is usually the first evidence of their presence, although a keen observer might detect a deposit of tiny black specks on the plant leaves. Arising from feeding, the insects leave white or silvery patches and streaks on leaves, fruit and pods. Citrus thrips causes grey or brown blemishes around the stem and navel ends of ripe fruit. Heavy infestation on beans causes a roughened silvery texture on the pod. Onion thrips gives a silvery appearance to the leaves of many vegetables. The leaves of grapes become silvery and the fruit may be scarred brownish red.

**RECOMMENDED REMEDIES**

Remedies specifically recommended in the literature are: Ash (A2 ix, xi); Clay etc. (A10 i, iv, v); Glues (A13); Hens etc. (A16); Manure etc. (A17 iii); Mulch (A19 i, ii); Oil (A21 i) — see warning; Soap solutions (A25 i, ii, iii); Water (A31 i); Onion (B 1.1); Tobacco (B 1.17) — see warning; and Various insects (C17 iii).

Potassium permanganate (A22) should be worth trying and, in addition, any of the Plant Mixtures (B3) and Insecticidal plants (B5) should be tested, particularly strong sprays from Pyrethrum (B 1.6), Rain tree (B 2.19), Cork bush (B 2.20) and Tephrosia (B 2.26).

Spray plants in blossom in the evening to avoid killing off bees and other useful insects.

The effectiveness of Oils (A21) should be investigated, particularly light, water-mixable mineral oils.

Adult thrips can be trapped by hanging blue or yellow sticky boards



THRIPS

in the nursery or garden. A light coating of clean engine oil can be used in place of glues.

Hens can clear pupa from the soil or the soil can be sprayed with rotenone products and liquid tobacco or dusted with dried tobacco.

Any of the measures mentioned for aphid and whitefly also have the potential to control thrips, though thrips are more difficult to treat because of their tendency to hide in buds and leaf sheaths and to lay their eggs in slits in the plant leaves. Spraying with lime/sulphur mixes, however, has been recommended.

The effectiveness of repellants (B6) is as yet unknown but routine spraying with repellants may have potential.

*Checks*

Since thrips feed on aphids and on their own kind they should not be eradicated entirely but a certain level of damage accepted as necessary to maintain the pest/predator balance. As is the case with all insects, going to the extreme of trying to kill off all members of a particular species often recoils in the form of an outbreak of other insects. Some species favour particular plants, and rotations should be a useful way of containing the problem.

**TIP OR TWIG WILTERS: (ANOPLOCNEMIS CURVIPES)***Host plants*

Beans, brinjals, cucurbits, granadillas, guavas, mangoes and many garden plants. Tip wilters are a major pest of cowpeas in Africa.

*Damage*

These bugs suck sap from stems, leaves, pods, buds and growing tips of a wide variety of plants causing them to wilt and die back.

*Life cycle*

Over 200 species of tip wilter have been recorded. The eggs are cylindrical and barrel-shaped, deposited in rows of about 15 on the stems of young or healthy plants. They hatch in about 14 days and, like shield bugs, the nymphs moult 5 times in the process of becoming adults. Two to four generations can be produced per year.

*Identification*

The adults are usually large, 25mm long, angular, usually pale grey to dark brown, elongated bugs which exude an unpleasant odour when handled roughly. They are winged and, in most species, the hind legs of the male are enlarged.

**RECOMMENDED REMEDIES**

**Adults:** remedies specifically recommended in the literature are: Biological (A4 i); Clay etc. (A10 ii); Hand picking (A15 i).

These large beetles can often be caught in nets: Beetles etc. (C3) and use of coloured traps: Beetles (C4) and light traps: Various insects (C17) may prove useful.



TIP OR TWIG WILTERS

Now Blooms  
use 20% B  
with 20%

\* Also worth trying are any of the insecticidal plants mentioned in B5 particularly sprays from Pyrethrum (B 1.6), Tobacco (B 1.17), Lantana (B 2.15), the Rain tree (B 2.19), Cork bush (B 2.20) and Tephrosia (B 2.26).

Nothing appears to be known concerning the effectiveness of Repellants (B6).

**Eggs on the leaf:** Ash (A2 i, ix, x, xi); Clay etc. (A10 i, v); Hand picking (A15 ii); Oil-mineral (A21 i); any of the insecticides listed in B6 particularly those \* mentioned previously for controlling the adults.

**Nymphs on leaves:** Ash (A2 ix, xi); Clay etc. (A10 i, ii, v); Hens etc. (A16); any of the insecticides listed in B6 particularly those \* mentioned previously for controlling the adults.

#### Checks

Ensure an adequate habitat for bird life that predate on the beetles and that there is a wide variety of insects to predate on the eggs and nymphs. Tree, hedge and plant barriers will often prevent the beetles from flying into the field. For the same reason intercropped or mixed cropped plants suffer less.

#### WHITEFLY: (ALEYRODIDAE)

##### Host plants

Whiteflies favour beans, brinjals, cotton, crucifers, cucurbits, okra, potatoes, pumpkin, oranges, sunflowers, tobacco and tomatoes.

##### Damage

The nymphs suck the sap causing yellowing of the leaves which weakens the plant. In the case of severe infestations the leaves may dry and drop off.

##### Life cycle

The female generally lays the oval white eggs in a circle on the underside of tender young leaves. Each egg is attached to the leaf by a short stalk. These hatch into flattened oval nymphs that can creep about at first but once a feeding site is found, they lose their legs and become static. The nymph moults 3 times and then becomes a pupa before turning into an adult.

##### Identification

The adults are very small, white, winged insects which are often clustered together on the plant. They can be seen easily by eye usually hiding on the underside of plant leaves. The whiteflies will dart away when the plant is shaken. The nymphs appear as minute yellow dots among the adults.

#### RECOMMENDED REMEDIES

Planting African marigolds, apple of Peru, nasturtiums or rhubarb next to or near susceptible plants is said to deter whitefly.

Mint and pigeon pea are useful decoy or trap crops.



WHITEFLY

Remedies specifically recommended in the literature are: Ash (A2 ix, xi); Clay etc. (A10 i, iv, v); Glues (A13); Hand picking (A15 ii); Oil (A21 i, ii, iii)— see warning; Potassium permanganate (A22); Salt (A 23 i); Soap (A25 i, ii, iii); Water (A31 i); Onion (B 1.1); Chilli (B 1.4); Tomato (B 1.12); Rhubarb (B 1.19); Blackjack (B 2.8); Lippia (B 2.17); Whitefly (C15).

Tobacco has been applied successfully both as a dust and as a spray (with soap added).

Good results have been obtained locally with Tephrosia (B 2.26) but any of the insecticidal plants mentioned in B5, particularly sprays from Pyrethrum (B 1.6), Tobacco (B 1.17), the Rain tree (B 2.19), Cork bush (B 2.20) and Tephrosia (B 2.26) should be worth testing.

Spray particularly on the underside of plant leaves.

Nothing appears to be known concerning the effectiveness of repellants (B6).

#### Checks

The little whitefly is windborne and has a strong preference for certain plants particularly tomato, mint and beans, therefore barriers and trap cropping are good methods of controlling them. They prefer humid, windless corners, nurseries and so on. Many small and large insects predate on them and the presence of a varied insect life should be encouraged. Check the soil for phosphorus deficiency.

#### WHITE GRUB: (SCARABAEIDAE)

##### Host plants

The grubs feed on roots of many plants. Attacks on cotton, crucifers, groundnuts, maize, protea, tobacco, tomatoes and grass roots have been recorded. The adult beetle of a large species called *Eulipida Mashona* has a preference for the leaves of *Julbernardia globiflora* and *Brachystegia spiciformis* trees.

##### Damage

The most damage is done between February and April. The plants wilt and die as the roots are eaten. At the end of the rains the grubs have a tendency to cut through the main anchor roots of maize plants causing them to dislodge. The adult beetle rests in the soil during the day and emerges at night to feed on plant leaves, making irregular holes. In the case of heavy infestations, whole plants may be defoliated.

##### Life cycle

The white grub is the larval stage of the Scarab 'chafer' beetle. It lives underground feeding on humus and plant roots. The adult beetle lays 30-60 eggs in the soil usually in November or December. The grub is fully grown by early winter and penetrates more deeply into the soil as the dry season advances. It then constricts an earthen cell in which to pupate and the adult beetle emerges with the first soaking rains. The whole life cycle takes about one year to complete.



WHITE GRUB



**Identification**

The grub is a large, fleshy, C-shaped insect with a brown head, three pairs of forelegs and blueish hindparts. It is often seen when digging into heaps of manure, compost or humus-rich ground.

Adults are light to dark brown chafer beetles.

**RECOMMENDED REMEDIES**

This insect is a classic example of a creature becoming a pest when its main source of food in the grub stage, soil organic matter and decomposing vegetation, is reduced or removed by farming/gardening activities. Further loss of the favourite food of the adult *Eulipida Mashona*, *Brachystegia spiciformis* and *Julbernardia globiflora* leaves, has resulted from the short-sighted policy of clearing all trees from arable lands. Loss of bird and small animal life which predate on the beetle has also contributed substantially.

The insect is attracted particularly by buried organic matter and less so by organic amendments applied to the soil surface. This is another reason for applying zero or minimal tillage techniques — not turning-in residues but leaving compost, manures and crop residues on the surface.

However, in the case of heavily-infested ground (presumably because of heavy amounts of organic matter buried previously), turning the soil to expose the grubs and then hand picking them might be the most practical solution.

**Beetle:** the beetles are large, hard-shelled and hide underground during the day. Nevertheless this is perhaps the most vulnerable stage of the insect as it can be readily caught at night by light traps: Various insects (C17). Hens etc. (A16) are useful for digging them out during the day.

**Eggs:** during the egg laying period November–December Hens etc. (A16) can easily dig the eggs from the soil. In addition spray the following well into the soil: Pyrethrum (B 1.6), Tobacco (B 1.17), the Rain tree (B 2.19), Cork bush (B 2.20) and Tephrosia (B 2.26).

**Grubs:** remedies specifically recommended in the literature for grubs are: Hens etc. (A16); Garlic (B 1.2); Tomato (B 1.12); Syringa (B 1.15) and Mexican poppy (B 2.5). A strong mixture of Pyrethrum and Tephrosia should be tried if all else fails.

The grubs are large, hardy and protected by the soil in which they hide. They are most vulnerable in summer as they bury more deeply into the soil as the winter advances.

**Checks**

Ensure sufficient soil humus, particularly old root systems such as can be encouraged by zero or minimum disturbance (tillage) of the soil. Are there enough *Brachystegia spiciformis* and *Julbernardia globiflora* trees in the vicinity to attract the Mashona beetle and to act as wind breaks? Encourage the return of birds particularly and tolerate a certain level of activity of predatory small animals such as moles, rats, hedgehogs, mongooses and so on.

**Diseases****BACTERIA****Host plants**

The leaves, fruits, stems and roots of a wide variety of plants. The following have been mentioned in local texts:

**Bacterial blight:** beans, potatoes, tomatoes.

**Bacterial canker:** apricots, cherries, peaches, plums, tomatoes.

**Bacterial spot:** almonds, apricots, cherries, mangoes, peaches, plums, beans, crucifers, lettuces, peppers, tomatoes.

**Common scab:** potatoes.

**Crown gall:** apricots, asters, cherries, peaches, plums, chrysanthemums, roses.

**Damage**

Bacteria cause browning and yellowing of the leaves, stems and fruit of plants, with rotting of the affected parts for example, blight, canker and bacterial spot. Other bacteria attack the root systems for example, common potato scab and crown gall.

**Life cycle**

Bacteria are minute organisms whose cells remain separate or may join up into microscopic groups or strands. They cannot be seen by the naked eye. Parasitic plant bacteria obtain their food from the host plant and thereby weaken it. Most bacteria reproduce by a cell dividing into two parts but a few kinds reproduce by a parent cell producing a new cell. Some types of bacteria live in the soil while others live on the above-ground parts of plants. Soil-borne bacteria can be spread by surface water (erosion) or simply by soil being splashed up on to plant leaves by raindrops or irrigation sprays. Infected soil may be moved from one place to another when filling seed boxes, transplanting, or by carrying it about on dirty boots, tools, and so on. Bacteria living above ground can be transmitted by wind, insects and infected tools and plant material.

**Identification**

The presence of bacterial attack can be identified through the declining health of the plant, taking into account the absence of insect damage and the presence of unsightly blemishes on plant parts or abnormal growths.

**RECOMMENDED REMEDIES**

Bacterial diseases can be best prevented by growing healthy plants (*Part 1; Section 1 Reinstate Soil Fertility*), by rotations (*Part 1; Section 3 Re-introduce Plant Diversity*) and by hygiene and other good husbandry practices (*Part 1; Supportive Strategies*).

If serious infections are allowed to build up it may be necessary to leave the land fallow for long periods (5–10 years).

Mulching will help to minimise the spread of diseases by preventing their transmission through soil splash. For the same reason, surface irrigation is better than overhead irrigation.

Ensure that irrigation water is free from diseases and has not been contaminated by diseased plants.

Windbreaks will minimise the spread by reducing the ability of wind to carry the disease.

A treatment of seed to prevent the spread of diseases is given in Part IV A1.

#### Supplementary treatments

These supplementary measures have been mentioned in the literature for the control of bacterial diseases: Ash (A2 viii, xii); Burning (A9); Compost (A12 i, ii); Manure etc (A17 i, iii); Sun (A29 ii); Water (A31 iv); Garlic (B 1.2); Chilli (B 1.4); Tomato (B 1.12); Goat weed (B 2.2).

Regular routine spraying with compost extract, manure and urine mixes are considered to be among the most effective all-round supplementary measures, resulting in removal or suppression of the disease, increased plant health and greater resistance to other diseases.

#### Checks

Check garden hygiene, and the use of resistant plants, time of planting, and plant spacing to minimise the spread of bacterial diseases. Check mulch quantity and use of windbreaks. Scout often and apply the supplementary treatments as necessary.

## FUNGI

#### Host plants

The leaves, fruits, stems and roots of most plants, including grasses. The following have been recorded on crops locally.

*Alternaria*: apples, carrots, citrus, cotton, onion, pears, tobacco.

*Anthraco*: avocados, beans, coffee, grapes, guavas, mangoes, cucurbits, tomatoes, sorghum, tobacco, roses.

*Botrytis Rot*: grapes, groundnuts, guavas, strawberries, tomatoes.

*Common Scab*: potatoes.

*Crown Rot*: carnations, groundnuts, onions.

*Damping off*: (seedlings) beans, cotton, crucifers, cucurbits, carrots, peas, groundnuts, tomatoes.

*Fusarium Blight & Rot*: beans, cabbages, carnations, onions, potatoes, sorghum, tomatoes.

*Early & Late Blight*: potatoes, tomatoes.

*Leaf curl*: deciduous fruits.

*Leaf Spots (Septoria & Cercospora)*: coffee, cotton, granadillas, groundnuts, lettuces, maize, pears, spinach, sorghum, sunflowers, tomatoes.

*Powdery & Downy Mildew*: apples, apricots, crucifers, cucurbits, deciduous fruit, grapes, mangoes, onions, peaches, pears, peas, peppers, plums, sorghum, sunflowers, tomatoes, vines, and many other garden plants.

*Root Rot*: avocado, citrus, parsley, sorghum, protea and other garden plants.

*Rot (fruit)*: apples, citrus, deciduous fruits, peaches, pears, plums, vines.

*Rusts*: apricots, beans, coffee, cucurbits, deciduous fruit, maize, millet, peaches, plums, sorghum, sunflower, and many other garden plants.

*Scab*: apples, citrus, deciduous fruits, pears, pecans, beans, cucurbits, deciduous fruit.

*Smuts*: barley, millet, sorghum, wheat.

#### Damage

Very many fungal diseases attack all parts of plants with loss of appearance of the produce, declining yields or loss of the plant. Most plants are susceptible to fungal diseases of one type or another, each type of fungus having a specific range of host plants.

#### Life cycle

Fungi are organisms which obtain their nutrients by drawing them from other plants. Fungal diseases are transmitted mainly by spores which may be carried to new sites by wind, water, soil, insects, birds, animals and by man *via* his tools and activities. Many fungi favour moist conditions.

#### Identification

Fungi can often be seen as moulds on the surfaces of fruit and leaves but may also appear as black or brown spots and blemishes on leaves, growing tips, blossoms and fruit, the fruit eventually rotting. Root rot can be identified by inspecting part of the root system when a plant wilts without any apparent cause.

## RECOMMENDED REMEDIES

Fungal diseases can be prevented best by growing healthy plants (*Part I: Section 1 Reinstate Soil Fertility*), by rotations (*Part I: Section 3 Re-introduce Plant Diversity*) and by hygiene and other good husbandry practices (*Part I: Supportive Strategies*).

Mulching will help to minimise the spread of diseases by preventing their transmission through soil splash.

Take care not to apply a greater amount of irrigation water than necessary. When overhead irrigation is being applied, apply the required amount of water in frequent small doses to avoid the leaf being wet for long periods.

Surface irrigation (including drip) is better than overhead as overhead irrigation wets the leaves and thereby creates an ideal environment for spore development. Timing the irrigation so that the plants do not remain wet overnight will help to contain the disease.

Ensure that irrigation water is free from diseases and has not been contaminated by diseased plants.

Select well-drained lands for planting particularly susceptible



crops like potatoes and tomatoes; otherwise grow the plants on graded ridges or beds.

Windbreaks will reduce the spread by reducing the ability of wind to carry the disease.

Other useful cultural techniques are to plant mixed varieties of the same crop on the same field, or to stagger the planting dates — not planting all the crop at the same time, plant the seed moving along the field into the prevailing wind, and choosing the right plant for the particular soil in the field, for example citrus and perennial crops generally prefer soil with less than 25 per cent clay.

A treatment of seed to prevent the spread of diseases is given in Part IV A1.

#### Supplementary treatments

The following remedies are recommended in the literature: Ash (A2 viii, x, xii); Baking soda (A3); Biological (A4 iii); Bordeaux mixture (A5); Burgundy mixture (A8); Burning (A9); Compost (A12 i, ii); Manure etc. (A17 i, iii); Milk (A18 i, iii); Mulch (A19 i); Oil-rape seed (A21 i, ii); Potassium permanganate (A22); Soap Solutions (A25 i, ii) — some soaps only; Sulphur (A28); Sun (A29 ii); Water (A31 ii, iii, iv); Onion (B 1.1); Garlic (B 1.2); Chilli (B 1.4); Pawpaw (B 1.5); Sunn hemp (B 1.7); Sweet potato (B 1.11); Tomato (B 1.12); Syringa (B 1.15); Oleander (B 1.16); Tobacco (B 1.17); Sweet basil (B 1.18); Rhubarb (B 1.19); African marigold (B 1.20); Mexican poppy (B 2.5); Thorn apple (B 2.11); Asthma weed (B 2.12); Rubber hedge (B 2.13); Wild tobacco (B 2.22); Wild basil (B 2.23); Castor oil (B 2.24); Mexican marigold (B 2.25) and Pigweed (B 4.4).

Bordeaux mixture has been found useful in the treatment of rusts on coffee.

Regular routine spraying with compost extract, manure and urine mixes are considered to be among the most effective all-round supplementary measures, resulting in removal or suppression of the disease, increased plant health and greater resistance to other diseases.

The infected leaves and fruit should be taken off the plant and burned, buried deeply or composted. Seed from diseased fruit or vegetables should not be re-sown as they will be infected with the disease.

#### Checks

Check garden hygiene, the use of resistant plants, time of planting, plant spacing and watering regime to minimise the spread of fungal diseases. Check mulch quantity and use of windbreaks. Scout often and apply the supplementary treatments as necessary.

### VIRUSES

#### Host plants

*Maize Streak Virus*: maize.

*Mosaic*: beans, crucifers, cucurbits, potatoes, tomatoes, roses and many other garden plants.

*Tomato Spotted Wilt*: chrysanthemums, dahlias, tomatoes.

#### Damage

Leaves and stems of plants may become discoloured or distorted affecting photosynthesis and nutrient uptake with loss of yield or death of young plants. Fruit may ripen unevenly and be deformed or blemished.

#### Life cycle

Unlike bacteria and fungi, viruses cannot reproduce themselves. They multiply by causing the cells of the host plant to produce new copies of the virus. In doing so, the virus produces toxic symptoms in the host plant and depletes it of nitrogen compounds. Plant viruses are usually transmitted from one plant to another by sucking insects both below and above the ground but can also be spread by grafting, by seeds and pollen, and via infected cutting and pruning tools.

#### Identification

The removal of nitrogen compounds by the virus may result in the first symptoms being a yellowing of the leaves as in mosaic or maize streak viruses. But toxic effects caused by the presence of the virus may show up first, in which case brown spots may be seen on the leaves, for example, tomato spotted wilt.

### RECOMMENDED REMEDIES

Viral diseases are extremely difficult to control once they are present.

If serious infections are allowed to build up it may be necessary to leave the land fallow for long periods (5–10 years).

They can be prevented best by growing healthy plants (*Part I: Section 1 Reinstiate Soil Fertility*), by rotations (*Part I: Section 3 Reintroduce Plant Diversity*) and by hygiene and other good husbandry practices (*Part I: Supportive Strategies*); including the use of virus-free plants and seeds.

In addition the number of sucking insects such as aphids, leafhoppers, mites, scale, tip wilters, thrips and whitefly should be kept strictly under control (not eliminated entirely); and their movement restricted by windbreaks.

In the case of viruses carried within the soil by nematodes and fungi, the best policy is to ensure a well-balanced soil-life population by minimal soil disturbance, compost additions, mulching and rotations.

#### Supplementary treatments

The following supplementary measures have been mentioned in the literature for the control of viral diseases: Ash (A2 viii); Burning (A9); Compost (A12 i, ii); Manure etc. (A17 iii); Milk (A18 i, ii); Mulch (A 19 i); Oil-mineral-paraffins (A21 i, ii); Chilli (B 1.4); Tobacco (B 1.17); Asthma weed (B 2.12); Wild tobacco (B 2.22); Snow berry (B 4.15).

Regular routine spraying with compost extract, manure and urine

mixes will help to build up plant resistance.

In the case of mild attacks sprays made with ash, milk or any of the plants mentioned above can be tried.

In case of serious infection or even mild infections failing to respond to the supplementary treatment, all infected material should be removed and burned.

#### Checks

Prevent outbreaks by checking that the soil is in optimal condition, the crops are healthy, that the varieties selected are resistant to the prevalent viral diseases; and that there is sufficient plant diversity in the form of rotational and other practices. Minimise the spread of diseases through garden hygiene, time of planting, and plant spacing. Check mulch quantity and use of windbreaks. Scout often and apply the supplementary treatments as necessary. Remove and burn badly-infected plants.

## Other Pests and Diseases

The following list gives the references to the treatments mentioned in Part III for pests not covered in detail in the foregoing text. In addition, references for specific diseases are given.

### PESTS

- Animals (general): A17 iv-vii; A20; B 1.2.  
 Ants (Leaf Cutting): B 2.24.  
 Aphid (Woolly): B 1.3; B 1.10; B 4.3.  
 Baboons: A17 vi; C5.  
 Beetle (Asparagus): B 1.12.  
 Beetle (Bean Leaf): B 2.16.  
 Beetle (Black Carpet): B 2.6.  
 Beetles (Flea): B 1.17.  
 Beetles (general): A10 i-iii; A25 ii; B 1.2; B 1.15; B 1.16; B 1.20; B 2.5; B 2.8; B 2.10; B 2.15; C3; C4.  
 Beetles (Mexican Bean): B 2.21.  
 Beetle (Red Pumpkin): B 4.6.  
 Beetles (stored produce): B 2.2.  
 Birds (general): A20; B 1.2; C5.  
 Bilharzia snails: B 1.17; B 2.6; B 2.14; B 2.22.  
 Blackflies: C17 iii.  
 Blowflies: B 2.25.  
 Buck: A17 iv-vii; A18 iv; B 2.1; C5.  
 Butterflies (general): C3.  
 Bugs (general): B 1.5; B 1.15; B 2.5; B 2.18.  
 Cabbage butterfly/worm: A18 i; B 1.1; B 1.12; B 3.5; B 4.6.  
 Cabbage root fly: A2 iii; A23 ii; B 1.4; B 1.15; B 2.8; B 3.6; C8 iii.  
 Canker moths: A26 ii.  
 Caterpillars (general): A2 i; A10 i-iv; A13; A17 iii; A21 i, ii; A22; A23 ii; A24 i; A31 i; B 1.2; B 1.5; B 1.12; B 1.14; B 1.15; B 2.5; B 2.6; B 2.17; B 2.18; B 2.24; B 2.25; B 3.1; C7.  
 Caterpillars (hairy): B 2.9; B 2.23.  
 Caterpillar (lawn): A19 ii; B 1.17.  
 Caterpillars (small): B 1.19.  
 Cockroaches: B 1.12; B 4.13.  
 Codling moth: B 1.10.  
 Cotton stainer: B 2.2; B 2.16.  
 Crawling pests (general): A14; A31.  
 Crickets (armoured): A25 i-iii.  
 Crickets (general): B 2.8; C8 ii.  
 Dusty surface beetle: C8 ii.  
 Eggs (general): A7 ii, iii; A10 i, v; A15 ii; A21 i-iii; A31 i, ii.  
 False codling moth: B 1.2.  
 False wireworm: C8 ii.  
 Fleas: A21 v; B 1.17; B 2.22; B 2.24; B 2.25; B 4.1.  
 Flies (general): A 17 ii; A21 v; B 1.12; B 1.14; B 1.16; B 1.18; B 2.21; B 2.23; B 2.25; B 4.2; C9; C17 iii.  
 Flies (house): B 1.2; B 2.2; C9.  
 Flies (vinegar): B 2.2.  
 Gnats: C17 iii.  
 Grubs (general): B 1.2; B 1.12; B 1.15; B 2.5; C7.  
 Hares: A17 vi; C13 iv.  
 Head lice: B 2.3.  
 Hoppers (general): B 1.15.  
 Larvae (general): A7 ii, iii; A10 i-v; A21 i-iii; A31 i-iii; B 1.12; B 2.10; B 4.2.  
 Larvae (leaf-eating): B 2.14.  
 Leaf hoppers: B 1.6; B 2.4; C17 iii.  
 Leaf miner: A25 ii; B 1.17; B 2.15; B 2.26; C10.  
 Lice: B 2.24; B 4.13.  
 Locusts: A20.  
 Maggots (general): B 1.18; B 1.19; B 2.23; B 2.25.  
 Maggots (onion): B 1.4; B 3.1-3.4.  
 Mealy bugs: A17 iii.  
 Melon worm: B 2.6; B 2.16.  
 Mice: B 1.1; B 1.2; C13.  
 Mites (general): A10 vi; A17 iii; B 1.12; B 2.24; B 3.6.  
 Mites (on chickens): B 4.1.  
 Moles: A20; B 1.1; B 1.2; B 1.15; B 2.24.  
 Mosquitoes: A21 v; B 1.2; B 1.18; B 2.10; B 2.13; B 2.23; B 2.25; B 4.2.  
 Moths (general): A19; C3; C12.  
 Peach borer: B 1.2.  
 Potato moth: B 1.18; B 2.23.  
 Psyllids: A25; A17 xi.  
 Rats: B 4.13; C13.  
 Rabbits: A17 vi; C13 iv.  
 Rodents (general): B 1.16.  
 Root maggots: A2 iii.  
 Seedbed pests (general): B 2.23.



Soft-bodied insects (general): A2 i; B 1.14; B 1.19.  
Shield bugs: B 2.26.  
Squash bug: A7 iv; A10 ix; A15 ii; B 1.3; C14.  
Stemborers (general): B 1.15; B 2.18.  
Ticks: A21 v; B 1.1; B 1.2; B 1.17; B 2.22.  
Tomato fly: B 1.1.  
Tomato hornworm: A15 i; B 1.12; B 2.21.  
Wild pig: C16.  
Weevils (banana): C2.  
Weevils (general): A2 iv-vii; B 1.15; B 1.16; B 2.5;  
B 4.3; C17 i.  
Weevils (maize): B 2.2.

#### DISEASES

Anthraxnose: B 2.24.  
Apple scab: A17 iii.  
Apple mildew: A17 iii.  
Bacterial wilt: B 1.12.  
Bean rust: B 1.2.  
Black spot: A5.  
Botrytis: A22; A28.

Brown patch: B 2.24.  
Canker: A10 viii.  
Club root: A2 iii; A10 x; B 1.19.  
Coffee berry disease: B 2.25.  
Damping off: A29 ii; A31 iii; B 1.1; B 2.24.  
Downy mildew: A5, A17 iii.  
Gooseberry mildew: A17 iii.  
Late blight: A12, B 1.1; B 1.20; B 2.25.  
Leaf curl virus: A5; A12; B 1.17; B 2.22.  
Leaf spot: B 1.1; B 1.5; B 1.11; B 1.16.  
Mildew (general): A2 x; A3; A8; A12; A17 i, iii;  
A18 iii; A22; A29; B 1.2; B 1.19; B 2.25.  
Mosaic virus: A18 i.  
Powdery mildew: A5; A8; A12; A17 i, iii; A28; B  
1.5.  
Rice blast fungus: B 1.11.  
Root rot: B 2.24.  
Rusts: A3; A5; A8; A17 iii; A28; B 1.5; B 1.17; B  
2.22.  
Scab: A17 iii; B 1.2.  
Tomato blight: A5; A8; A17 iii; B 1.2.  
Vine powdery mildew: A17 iii.

## PART III

# REMEDIES FOR PROTECTING FIELD CROPS

## A Miscellaneous Substances and Methods

Where spraying equipment is not available, the liquid can be flicked on to the plant with paint brushes, brooms or brushes made from local plants.

### A1: AROMATIC PLANTS

#### Materials

Leaves of any strongly-smelling plants like gums, lantana, khaki weed, tomato or any herb.

#### Targets

Any insect.

#### Methods

Make up a spray from the leaves of strongly-smelling plants either using 1 plant or mixing 2 or more together. The strong smell will repel pests. Garlic, chilli and onion sprays are recommended particularly.

The plant material is dried, ground to a powder, mixed with boiling water and allowed to cool before spraying. Recommended mixes vary from 20–500g per litre of water.

#### Frequency

Spray just before peak infestation is expected in early summer and repeat as often as necessary depending on the number of pests evident that year. More applications will be needed during the rain as the substance will be washed off the plants.

#### Warning

Some crops are damaged by juice from other plants, therefore watch out for signs of damage (usually deformed leaf growth), try to identify the problem plant and leave it out in future; or simply avoid using it on that type of crop.

### A2: ASH

#### Targets

Soft-bodied insects including aphids, beetles, cabbage root fly, caterpillars, cutworms, grasshoppers, eggs, larvae, nematodes, potato moths, pupae, root maggots, slugs, snails, squash bugs, stalk borer, termites, weevils and insects in general; diseases with mildew and club root being specified; and other pests.

#### Materials

Wood ash from plant parts. Ash from specific vegetation has been found to be more effective against specific insects. Ash from acacia nilotica, casuarina, cyprus, combretum imberbe, eucalyptus, mango wood, finger millet, rice trash, spirostachys africana and tamarind have been found to be particularly effective as disinfectants or as general insect repellents — see *warning* that follows.

#### Method

- i Dust ashes evenly on to vegetables to dehydrate soft-bodied insects or into the funnel of young maize plants to kill stalk borer. Dusting crop leaves with the ashes of combretum imberbe and spirostachys africana are said to control aphids and grasshoppers.
- ii Spread ashes thickly around plants or seed beds to deter soil pests: cutworms, eggs, larvae, pupae and so on; preferably in a 150–200mm wide trench 20–50mm deep to deter slugs and snails. The ash of combretum imberbe and spirostachys africana is said to control cutworm when mixed with the soil in the planting holes.
- iii Spread around the roots of radish, onions, cabbage and other crucifers and cover lightly with soil as a protection against cabbage root, fly maggots and club root.
- iv Spread over the soil surface as a mulch to control nematodes and weevils.
- v Incorporate into the soil during planting of any plants including in forestry nurseries to protect seedlings against termite attack. This treatment has been found to protect sweet potato from the sweet potato weevil.
- vi Heap ashes around the base of tree trunks to prevent them from termite attack.
- vii Ashes mixed with compost and dug into the soil can control weevils and nematodes and greatly improve soil fertility.
- viii Dip cuttings into ashes mixed in water to reduce diseases and increase the percentage take.



- ix Spray with wood ash mixed with soapy water and/or lime as a general insecticide.
- x Spray with a mixture of 1 heaped tablespoon of wood ash stirred vigorously into 1 litre of water, leave overnight, strain to remove solids, mix with 1 cup of sour milk and add 3 litres of water. This treatment has been found effective against mildew — see also *milk treatments* — and a range of pests and their eggs.
- xi Wood ash mixed in lime sprays is said to increase its potency against certain insects like squash bugs. Allow mixture to stand for 1–2 days before use.
- xii Soaking seed in wood ash and water for 24 hours before planting gives protection against fungal and bacterial diseases.
- xiii Soak seeds in ashes mixed with water or with the juice or leaves of any strongly aromatic plant added to the water to deter birds from eating the seed (Maize seed has been identified). Test the effect of new aromatic plant material on germination using small quantities of seed before using it as a general treatment; some strong-smelling plants can affect the viability of seed.

**Frequency**

Apply as frequently as necessary. More applications will be needed for surface ashes (required as a dry barrier) during wet or windy periods. Dusts are said to be more effective applied to plant leaves when they are moist from rain, irrigation or dew.

**Warning**

Wood ash contains variable amounts of caustic potash so it should not be used on hot days and on tender plant growth. Do not allow ashes to touch the stems of plants, particularly seedlings.

Ashes should be allowed to cool before use. Do not use coal ash.

The milky latex from *spirostachys africana* should not touch the skin or enter the eyes. Protect eyes from sawdust and ashes.

**A3: BAKING OR WASHING SODA****Materials**

Baking soda or washing soda, water and soft soap.

**Targets**

Fungal diseases (mildew and rust mentioned particularly) and scale.

**Method**

Spray with a mix of 100g of baking or washing soda with 50g of soft soap. Dilute with 2 litres of water or more.

**Frequency**

Spray only once preferably or leave as long a gap as possible (several months) between applications.

**Warning**

Use immediately but never during very hot weather. Avoid spraying the leaves and flowers as the spray is known to damage plant tissue. For the control of scale, spray onto fruit trees when they are dormant (leafless).

**A4: BIOLOGICAL CONTROLS****Materials**

Cultural methods detailed in Part I should be applied. See summary at the end of Part I for quick reference. Apart from cultural methods, some simple biological controls can be applied as follows.

**Targets**

Insects in general.

**Methods**

i Spray with a mixture made up of a handful of the pest insects crushed and mixed into 10 litres of water. Add a little liquid soap. Leave for 12–24 hours. Pests of the same species have been found to evacuate crops treated in this way. The insect bodies remaining after sieving the liquid can be left in open containers (jars, tins etc.) in the garden as the smell will continue to repel the insects.

This treatment is said to be effective on armyworm and other caterpillars, millipedes, saw fly, slugs and various bugs; but less successful on grasshoppers and locusts.

ii Spray with a mixture made up of a handful

of the insects which are showing signs of distress from a natural disease. This spreads the disease to the healthy insects of the same species, or even to other species.

iii Control the humidity by altering the spacing distance between plants. Close spacing discourages red spider mite while wide spacing discourages fungal and other diseases.

**Frequency**

Treatment i may have to be applied several times, particularly during the rains. Treatment ii normally need only be applied once but monitor the rate of spread of the disease and assist it with further spraying if necessary.

**Warning**

Wear plastic or rubber gloves as certain insects contain harmful substances, for example, blister beetles.

**A5: BORDEAUX MIXTURE (SEE WARNING)****Targets**

Fungi; powdery mildew, downy mildew, tomato and potato blight, black spot on beans, leaf curl on peaches, rust and as a general fungicide.

**Method**

Mix 90g of blue copper sulphate with 4.5 litres of water in a non-metallic container. In another non-metallic container, mix 125g of slaked lime (builder's lime) with 4.5 litres of water. Stir and make sure the lime is well dissolved. Mix the two solutions and stir well. Test the mixture by putting an old nail in it for 30 seconds. If it turns blue there is not enough lime in the mixture or the lime is not sufficiently dissolved in the water. Allow the lime to soak longer, otherwise the mixture will burn the leaves. Use the mixture immediately.

**Frequency**

Apply twice at 7-day intervals but only if necessary.

**Warning**

Do not use regularly in the same place as it kills many beneficial fungi and pest predators as well.

Some plants are sensitive to copper and sulphur so test first if uncertain. At high concentrations the mixture is toxic to soil organisms, insects, plants, animals and humans.

**A6: BRUSHING****Materials**

A stiff brush.

**Targets**

Crawling insects and swarms.

**Methods**

- Swarms of beetles and caterpillars can be swept into piles with a stiff broom and destroyed.
- Insects, larvae and eggs can be swept from leaves, branches and trunks of fruit trees with a stiff brush.

**Frequency**

Carry out as frequently as necessary.

**Warning**

Protect the eyes (wear dark glasses) from falling dirt, insects and so on. Some insects can seriously injure the eyes.

**A7: BUILDER'S LIME (SEE WARNING)****Materials**

Builder's lime (quicklime), which is called slaked lime after water has been added.

**Target**

Eggs, larvae, slugs and snails, and squash bugs.

**Method**

- Dust over nursery floors, or where there is no plant material, to kill slugs and snails.
- Brush freshly-mixed lime and water on to the hardened bark of mature trees to kill eggs and larvae on the trunk. Old bark under which insect larvae and eggs can be hiding should be scraped off first with a wire brush.
- Spray lime and water mixtures onto tender plant parts (leaves etc.) only after it has been allowed to stand for a few days until the heat of hydration has worn off — see also *Bordeaux mixture*.

- iv Mix builder's lime and wood ash with water to the consistency of thin soup, allow to stand for 1-2 days before spraying on squash bug.

**Frequency**

Treatment i may need to be repeated every time the slug and snail population increases. Treatment ii can be applied once only in winter but should be re-applied every few months during the rainy season.

**Warning**

Do not allow builder's lime-dust onto the skin or into the eyes as it burns on contact with moisture. Do not apply as a dust to tender plant parts such as leaves for the same reason.

## A8: BURGUNDY MIXTURE (SEE WARNING)

**Materials**

Copper sulphate, washing soda and water.

**Targets**

Fungal diseases.

**Method**

Dissolve 200g of copper sulphate in 9 litres of water. Then dissolve 30g of washing soda in 9 litres of water. Mix the two and add another 4.5 litres of water. Spray on affected plants.

**Frequency**

Use only once a year at the most.

**Warning**

Use with even greater caution than Bordeaux mixture. Apply only to dormant fruit trees or vines i.e. woody species not in leaf. Some plants are sensitive to copper and sulphur so test first if uncertain. At high concentrations the mixture is toxic to soil organisms, insects, plants, animals and humans.

## A9: BURNING (SEE WARNING)

**Materials**

Infected plant material.

**Targets**

All pests.

**Method**

Burn all affected plant material when dry or mix with dried inflammable material if green.

**Frequency**

Burn until all infected material is reduced to ashes. The ashes will then be safe to use for further pest and disease control both in the field and in storage.

**Warning**

Ensure that the fire does not spread. Great environmental damage is caused to the pest/predator balance (see *Armyworm* in Part II), to the soil (loss of organic matter and soil life), and to the atmosphere (greenhouse effect) by both intentional and unintentional bush fires. Even small fires may be prohibited in future should the situation worsen. Keep small children away from the fire.

## A10: CLAY, LIME AND OTHER POWDERY MATERIALS

**Materials**

Any fine powdery substance such as clay, crushed laterite, agricultural lime, flour, chalk, rock dust and so on.

**Targets**

Ants, aphids, beetles, canker, caterpillars, codling moths, cutworm, eggs of insects, larvae (any), mites, scale, slugs, snails, squash bugs, stink bugs, thrips, tip wilters and whitefly; club root.

**Methods**

- i Dust the fine material onto the leaves, the insects (aphids, mites, thrips, whitefly) or their eggs, to suffocate them. Lime dusted over loopers worked very well; lime dusted onto small black slugs was effective and has been recommended also for controlling small beetles.
- ii Leaves can be dusted with clay and laterite to protect them from attack by sucking insects (aphid, mites, scale, stink bugs, tip wilters and so on), but the treatment is only temporary, especially during the rains.
- iii Dust around the base of the plants to protect them from any insect trying to crawl up the stems (ants, slugs, snails, cutworm, beetles and so on); or trying to lay eggs in the soil near to the plant.
- iv Spray with a mixture of the powdery material in water. Add material (clay etc.) and

stir thoroughly until the mix is like very thin soup. Allow to stand for a few minutes and then pour the liquid into another container so leaving any solids behind. Stir in a little soap to give a slight froth on the surface. Apply with a watering can (with sprayhead) or splash on with a large paint brush or bush-broom. Do not use sprays with fine nozzles as they will get blocked. Apply during a dry spell so that the liquid has time to evaporate and the fine dry material hardens around the pest. A lime spray worked well on loopers — see also *dusting i* — and flour sprays have worked well on aphids, mites, whitefly and thrips.

- v Spray with a mixture of 4 cups of flour, half a cup of sour milk and 20 litres of water to destroy eggs, larvae and adult aphid, mites, thrips and whitefly.
- vi Some old gardening books recommend a lime-water spray for controlling mites such as red spider mite. Lime water is the clear water remaining on top after all the lime itself has settled to the bottom of the bucket.
- vii Mix clay with fresh cattle manure to form a paste and paint the mixture on trunks of fruit trees against codling moths and other pests, and to seal the fresh cuts on the trees after pruning.
- viii Fill wounds in trees, including those caused by cutting out canker, with moist clay rammed in tight (a little cement, putty or builder's lime can be added for durability) and the surface smoothed over.
- ix Wood ashes mixed in lime sprays into a thin soup is said to increase its potency against certain insects like squash bugs. Allow mixture to stand for 1-2 days before use.
- x Liming acid soil is said to reduce the incidence of cutworm and club root.

**Frequency**

The treatments are effective for short periods and must be repeated as often as necessary, depending mainly on the weather and the presence of insects. Lime dusts are said to be more effective when applied to the plant leaves when they are moist from rain, irrigation or dew.

**Warning**

Frequent use of clay preparations will have no harmful effect on soil with lots of organic matter in it but clay and other powders may increase surface crusting on soils in poor condition. Builder's lime can be used mixed with water as a spray and applied after several days when the heat of hydration has worn off; but it should never be used as a dust on leaves or they will be scorched.

Application of agricultural lime will at first benefit acid soils, particularly if inorganic fertilizers have been used in past years, but the soils may become overly alkaline if heavy dusting is repeated too often.

Do not allow lime dust to touch the stems of seedlings.

## A11: COFFEE GROUNDS AND TEA LEAVES

**Materials**

As above, either waste or fresh.

**Targets**

Insects in general, slugs and snails.

**Method**

- i Spread on the surface of the soil around the plants to protect plants from slugs and snails.
- ii As sprays, fairly strong brews made from tea leaves and coffee grounds are said to have general insecticidal properties.

**Frequency**

Use as frequently as necessary.

**Warning**

Prolonged use may create acid conditions in the soil.

## A12: COMPOST

**Materials**

Well rotted-down compost made with plants and animal manures.

**Targets**

Most pests and diseases (by giving the plants greater resistance to attack). Rusts are said not to respond to compost sprays.

**Method**

- i Spray the plants and soil with a mixture of 1



shovelful of compost in 20 litres of water after the mix has been standing for 3 days to several weeks. For smaller amounts put 1 sock full of compost (about 0.5 kg) in a 20 litre bucket. Apply as a foliar spray to supply the plant with much-needed nutrients to help it withstand attack and to increase yield. The longer the mix is left to stand, the stronger and more effective it becomes. The spray should be applied once a week when the plants are young.

ii Increasing the compost level in the soil protects the plant against all pests and diseases. The compost is best applied to the soil surface for reasons discussed in Part 1.

#### Frequency

Apply the compost spray as a preventive measure and as a regular weekly treatment once the disease appears. Feed the plants with a palm full of compost at planting, before fruiting and after every harvest or cutting.

#### Warning

None known.

### A13: GLUES

#### Materials

Any water-soluble glue, particularly those obtained from plants. Water from boiling potatoes and cassava, if it contains sufficient concentration of starch.

#### Targets

Aphids, caterpillars, spider mites, thrips and whitefly.

#### Method

Spray a very weak solution of a water-soluble glue to suffocate the insects. The strength of mixes vary widely according to the strength of the glue available but the diluted solution should leave a thin skin coating on the plant when the solution has dried.

#### Warning

None known.

### A14: GREASE

#### Materials

Any thick grease.

#### Targets

Ants and any crawling pest in the orchard or nursery like fruit fly larvae, slugs, snails, beetles, caterpillars and so on.

#### Method

- i Wrap tin foil or durable plastic around the tree trunk and smear it with grease forming a barrier ring around the trunks of fruit trees. Fold over the top of the foil or plastic to form an overhang to protect the grease from being washed off by rainfall.
- ii Put a wide band of grease directly onto the wood or metal of nursery tables to prevent crawling pests from reaching the seedling trays.

#### Frequency

Check the grease every week to ensure that it is intact.

#### Warning

Do not place the grease directly on to the surface of the trunk and branches of young trees otherwise they will be damaged and may even die.

### A15: HAND PICKING

#### Materials

Gloves are sometimes advisable — see warning.

#### Targets

Many insects and chrysalises.

#### Method

- i Large insects can be picked off the foliage by hand and fed to poultry, destroyed or thrown into the bush.
- ii In the case of heavy infestations by small insects, eggs or larvae, whole leaves can be nipped off and disposed of or the insects crushed on the plant by rubbing over them with the fingers.
- iii Children were able to fill a 500ml jam jar each per day with grasshoppers that they chased and caught by hand — see also *Beetles, butterflies etc C3* for use of nets.
- iv Cutworms, slugs and snails can readily be seen and hand picked at night by torch or lamplight.

#### Frequency

The treatment is repeated as often as possible.

#### Warning

Wear thick gloves when dealing with blister beetles, hairy caterpillars or any stinging insect.

### A16: HENS, BANTAMS AND DUCKS

#### Materials

Hens, bantams or ducks, a mobile coop, some chicken wire and string.

#### Targets

Almost all insects.

#### Method

Hens and bantams will catch and eat the most troublesome pests including beetles, caterpillars, cutworm, grasshoppers, and snails. They can run quickly to catch flying insects and their scratching will uncover the larvae, pupae and eggs of many pests hiding in the soil. While feeding on the pests, they also manure the ground.

There are a wide variety of ways in which fowl can be used. They can be allowed to roam freely in orchards, or on lands not yet planted, or where hardy crops are growing which will not be damaged by the hens either feeding from them or scratching around them. Alternatively, they can be cooped up in portable pens built to the exact size of the garden beds; or a mobile fence can be used to confine the hens to the required area. The services of hens and bantams may be required urgently away from home with no time to organise fencing. In this case they should be transported to the site as humanely as possible. Once there, a 3 metre length of string is tied to one leg of each hen, which enables them to be easily caught at the end of the day.

Ducks are particularly useful for removing slugs and snails.

#### Frequency

The hens/bantams can be confined on garden beds to clear them of insects prior to planting. They can be concentrated around the foot of fruit trees in early spring and throughout summer to clear fruit fly larvae from the soil and the grub from the fallen fruit.

#### Warning

Hens and ducks, particularly, can do considerable damage to seedlings by eating them or by digging them up during their scratching activities. Therefore do not allow them near seedbeds. Bantams usually prefer insects to vegetation and a flock which does not eat any leaves can be developed simply by getting rid of any bantam showing a taste for 'greens'. They are great scratchers, however, and should be kept away from young and/or delicate plants.

In winter when green material and insects are scarce, fowl can do much damage to crops. They require 'herding' at such times.

### A17: MANURE, DUNG AND URINE

#### Materials

Cow, donkey, goat, predatory animal and poultry manure; human and animal urine.

#### Targets

Animals, aphids, bagrada bugs, birds, caterpillars, codling moths, cutworm, fruit flies, grasshoppers, mealy bugs, mites and thrips, fungal, bacterial and viral diseases.

#### Method

i Spray with a mixture made up of 1 shovel full of dry cow manure mixed thoroughly into 10 litres of water after it has been stirred once daily for 14 days. Clay dust can be sprinkled into the bucket to reduce smells and to add useful minerals. Dilute the mixture 3-5 times before spraying. This fermented dung water will repel aphids when applied to plant leaves and protect plants from cutworms when sprayed on the soil surface around seedlings. It also acts as a foliar nutrient when sprayed on leaves thus overcoming nutrient deficiencies in the plant and thereby increasing the plant's resistance to bacterial and fungal diseases. The liquid also acts as a fertilizer when sprayed on the soil.

A shovelful of fresh donkey manure mixed in a bucket of water and left overnight is said to be effective as a spray for aphids, bagrada bugs and grasshoppers.

- ii Spray leaves and fruit with a mix of 1 part burnt cow dung to 2 parts water to repel aphids and flies (including fruit flies).
- iii Spray with a mixture of 1 part urine (human or animal) to 1 part water to prevent aphid, caterpillars, cutworm, mealy bugs, mites, thrips, and for protection against fungal, bacterial and viral diseases. Spray on a warm day and use as a preventive measure. Weaker sprays of 1 part human urine to 4 parts water have been recommended for fungal diseases: apple scab, apple mildew, gooseberry mildew, downy mildew, vine powdery mildew and others.
- iv Paint a mixture of clay and cattle manure onto the trunks of fruit trees to protect them from codling moths and other pests. Wood ash can also be added to this mixture. The mixture can be applied to protect fresh cuts after pruning and to repel animals from eating the bark or browsing on the leaves.
- v Paint tree trunks with a thick soupy mixture of fresh dung (of the animal to be repelled) stood in water for 3 days.
- vi Paint tree trunks with a mixture of goat, cow, poultry manure, clay dust and animal urine at monthly intervals to repel baboons, hares, buck, goats and rabbits.
- vii Spray plants with liquid goat manure to repel hares and buck; or hang up bags of manure obtained from predatory animals (cats, dogs, lion and so on).
- viii Manure, dung and urine added to the soil improves soil fertility and protects plants from many pests and diseases.
- ix Spraying with cow urine that has been standing for two weeks and diluted with 6 parts of water is said to control stalk borer (presumably the eggs and new caterpillars).
- x Soaking seeds for 24 hours in cow dung mixed with water is said to prevent birds from digging them up afterwards.

**Frequency**

Apply as often as required. The protection is needed most in the winter when there is a shortage of green

material in the surrounding bush.

**Warning**

Vegetables or fruit sprayed with these mixtures must be washed thoroughly before consumption. Undiluted urine will burn plant leaves and will render soil toxic if used too often.

**A18: MILK****Materials**

Milk, paraffin, soil and water.

**Targets**

Fungal and viral diseases in general with blights, mildews, mosaic viruses on tomato, tobacco, sugar cane and sorghum being mentioned particularly; spider mites, caterpillar eggs on crucifers, and buck.

**Method**

- i Spray every 10 days with a mixture of 1 litre of milk to 10–15 litres of water to control mites, blights, mildew, mosaic virus and other fungal and viral diseases on a wide range of plants. This spray is said to also control spider mites and caterpillar eggs on the crucifer family when sprayed every 3 weeks.
- ii Sprays of skimmed milk have been used to control many viral diseases.
- iii Spray with a mixture of 1 heaped teaspoonful of wood ashes stirred vigorously into 1 litre of water, leave overnight, strain to remove solids, add 1 cup of sour milk and add 3 litres of water. This treatment has been found effective in controlling mildew.
- iv Painting a mixture of soil, sour milk and paraffin to halfway up tree trunks has been found to deter buck from eating the bark. This treatment was effective for 2–3 weeks. Protect the mix from being washed off by heavy rain.

**Frequency**

Frequencies are given above as appropriate.

**Warning**

None known.

**A19: MULCH****Materials**

Dead plant material: crop residues, hay, leaves, weeds etc. and tobacco leaves.

**Targets**

Cutworm, grasshoppers, ground beetles, lawn caterpillar, moths, nematodes, slugs and snails, termites, thrips and a wide variety of pests and diseases.

**Method**

- i Cover the soil with a thick layer of mulch, one finger deep preferably. The mulch should consist of as wide a variety of plant material as possible (dry). To repel termites, include banana stems and mixtures of aromatic plants and weeds such as lantana, Mexican marigold, gum tree leaves and so on. Mulches of aromatic herbs are recommended for protecting plants from attack by cutworm, ground-beetles, thrips, slugs and snails. Covering bare soil with mulch discourages grasshoppers, moths and certain beetles from laying their eggs in the soil. Building up humus reduces numbers of harmful nematodes.
- ii A mulch of tobacco leaves or dust will control cutworm, lawn caterpillars, thrips, slugs and snails and many other insects; but does not harm many beneficial ones such as earthworms.

**Frequency**

Add to the mulch whenever its depth decreases. It may need to be added to about 4–6 times per year.

**Warning**

Before applying the mulch, make raised beds or ridges on any soil prone to waterlogging. Do not use green plant material on seedbeds otherwise the humic acids will acidify the soil and inhibit germination of some seeds.

**A20: NOISE AND VIBRATION****Targets**

Locust, birds, moles and wild animals.

**Method**

Banging on tins and drums has been used for genera-

tions to scare pests away from crops.

Moles are particularly sensitive to vibrations in the soil. They can be driven away by frequent hammering on the ground. A simple windmill can be built with a pole attached to one arm. The pole is placed so that it thumps the ground at every turn of the windmill.

A noisy radio playing beat music is said to keep away baboons and other animals from cropland.

Birds can be deterred by putting a few dry beans or maize seeds into paper bags and tying the bags to a string line so that they rattle in the wind.

**Frequency**

Loud continuous banging is the most effective. Birds soon get used to infrequent bangs especially if the interval between bangs is regular; however, consider your neighbours!

**Warning**

Constant exposure to loud noises can impair hearing and upset the nerves; however, in pest control, noise is unlikely to be used with sufficient volume and frequency to be a danger.

**A21: OIL****Materials**

Light grade water-mixable mineral (petroleum) oil or vegetable cooking oil.

**Targets**

Aphid, caterpillars, fleas, flies, insect eggs and larvae, mites, mosquitoes, red spider mite, scale, thrips and whitefly, viruses (mineral oil from paraffins) and fungal diseases (rape seed oil).

**Method**

- i Spray with a mix of 1 litre of ordinary vegetable cooking oil with 100g of bar soap (or 100ml of liquid soap) and 15 litres of water. Stir thoroughly to give a cloudy emulsion — see warning below. Light grade water-mixable mineral oil can be used as an alternative at 30ml of oil to 1 litre of water.
- ii Spray on vegetable or light mineral oils undiluted — see warning below.
- iii Spray with a solution of sufficient fatty soap made from vegetable oil to give a frothy



surface to the water.

- iv Brush 2ml of mineral or vegetable oil on to the end of the maize silk after it has wilted but before it has begun to dry off in order to protect the cob from insect attack.
- v Citrus oil is said to repel fleas, flies, mosquitoes and ticks.

#### Frequency

Apply as infrequently as possible. The treatment is very effective.

#### Warning

Take care to avoid beneficial predators. Do not spray vegetable oil (either diluted or undiluted) onto leaves unless you have first tested the reaction. Some shiny leaves like mango and sisal are not affected whereas many others are destroyed by the oil. Avoid spraying on hot days or even shiny leaves may be scorched. Spray deciduous fruit trees only in winter when they are dormant. Light grade mineral oils do not appear to harm plant leaves when the oils have a paraffin base.

## A22: POTASSIUM PERMANGANATE

#### Materials

Potassium permanganate and water.

#### Targets

Aphids, armyworm, caterpillars, diamond-back moth, spider mites, whitefly, botrytis and mildews.

#### Method

Dissolve 30g of potassium permanganate in 9 litres of water and apply as a spray. The effects last for only 24 hours but the spray feeds the soil with potassium.

Potassium permanganate dusted on to plants is said to control mildew and botrytis.

#### Frequency

Several applications may be necessary for maximum effect.

#### Warning

Avoid predators. Do not use regularly or in large quantities otherwise the soil and plants will suffer from manganese toxicity.

## A23: SALT PREPARATIONS

#### Materials

Salt, vinegar, water and soap.

#### Targets

Aphids, cabbage worm, caterpillars, slugs, snails and whitefly.

#### Method

- i Spray aphid and whitefly with a mixture of 1 teaspoonful (5ml) of salt to 1 tablespoonful (20ml) of vinegar and mix with 1 litre of water. Dissolve half a teaspoon (2.5ml) of liquid soap.
- ii The above mixture without vinegar has been recommended for use on aphids, cabbage worm and caterpillars; and as a general repellent.
- iii Sprinkle a few grains of dry salt on slugs and snails.

#### Frequency

Apply the salt and vinegar spray twice at 5-7 day intervals. Apply sprinkled salt as often as necessary to keep snail population to low numbers. At night or during damp cloudy days are the best times for hunting slugs and snails.

#### Warning

Do not spray the same area too frequently with salt solutions unless sufficient extra water is being applied to flush the salt out of the soil. Apply dry salt even less frequently to prevent the fertility of the soil from being damaged.

## A24: SMOKE AND SOOT

#### Targets

Caterpillars, cutworm, locusts, slugs, snails and termites.

#### Method

- i Dust the soot onto vegetables to control caterpillars or spread it as a barrier on the soil around seedlings to protect them from cutworms, slugs and snails.
- ii Smoke fires in the field have been used to repel swarms of locusts and other insects but have minimal effect.

- iii Tobacco smoke is effective in repelling termites and other insects.

#### Frequency

Apply as often as necessary.

#### Warning

Smoke is an irritant to the eyes and lungs. People who constantly expose themselves to smoke will be more susceptible to eye and lung infections.

## A25: SOAP SOLUTIONS

#### Materials

Liquid soap of the kind used for washing the dishes, solid bar soap and soap powders (modern detergent soap powders are not acceptable to the Zimbabwe Organic Producers' Association). Soft soaps made from potassium compounds are preferred as they have been found to be the most effective against insects and also to add potassium to the plant through the leaves and the soil.

#### Targets

Aphids, armoured crickets, armyworm, caterpillars, leaf miner, mites, psyllids, white fly, small beetles, slugs, snails and thrips. Some soaps are said to be effective against certain fungi.

#### Method

- i A spray of half a tablespoon (10ml) of liquid soap in 1 litre of water has been found effective for aphids and whitefly.
- ii Stronger soap solutions are needed for stronger insects. The following mixtures have been found effective: 1 tablespoon (20ml) of soap powder to 1 litre of water and 7g of bar soap to 1 litre of water, with or without paraffin added.
- iii Any soapy water left over from washing the clothes or dishes can be diluted into a suitable spray.
- iv Soap is also added in small quantities to other preparations as it helps the spray to stay on the leaves and to spread out into a thin coating.

#### Frequency

Apply as few times as necessary to control the insects to low levels. As always, spray selectively (on the pest only) and avoid killing off the predators.

#### Warning

The amount of soap needed will depend upon the soap type and the users should experiment for themselves to find the correct mix. Strong soaps such as modern detergent for washing clothes can scorch plant leaves because of the caustic soda contained in them. Consistent use of these soap powders will also damage the fertility of the soil. Wherever possible use soft soap made from potash as it adds potassium to the soil. Even soft soaps can harm plant leaves, however, if applied too frequently or if the mixture is too strong. Young and fleshy leaves (not smooth and waxy) are the most vulnerable.

## A26: SOIL

#### Materials

Ordinary topsoil but not pure sand.

#### Targets

Maize stalk borer and larvae of fruit fly and cankerworm moths.

#### Method

- i To control maize stalk borer, pick up just enough dry soil from the ground next to the plant to block the central funnel of the plant. The soil firmly blocks the funnel when it rains and dries into a hard block during dry spells. In both cases the borers are smothered. Also, eggs are laid on the upper leaves of the maize plants. Borers can be prevented from crawling down these leaves and penetrating the stem by filling the 'v' made where the main upper leaves meet the stem with soil.
- ii Disturb the soil under fruit trees to enable poultry and birds to pick up the larvae of fruit flies and cankerworm moths.

#### Frequency

Where high infestations of stalk borer are the rule, use the method as a preventative by applying when the maize is very small and repeat later on any plants showing signs of leaf damage. When borer infestations are not common, apply as early as possible but to damaged plants only.

#### Warning

Do not use pure sand for controlling stalk borer as the

caterpillars can force their way through the voids between the sand particles and will not be smothered. Sand has been proven completely ineffective.

## A27 : SUGAR

*Targets*  
Nematodes.

*Method*  
Dissolve 2kg of sugar in a bucket of water and pour out over the soil to dehydrate nematodes. The treatment will kill all nematodes within 24 hours. The sugar should be flushed out of the soil with plenty of extra water after 24 hours. This treatment is not economically practical for large areas.

Soaking seed potatoes in sugar water for 24 hours prior to planting has been used to remove nematodes.

*Frequency*  
This treatment should be carried out as infrequently as possible, and applied only once on any soil infested through past poor treatment.

*Warning*  
Many nematodes are beneficial to the soil and keep each other in balance. Also, other useful soil organisms will be killed off by this treatment.

## A28: SULPHUR

*Targets*  
Spider mites, scale, fungi: especially botrytis, powdery mildew, and rust on beans and asparagus.

*Method*  
Dust flowers of sulphur on infected parts of plants. Lime sulphur is sold commercially and consists of 1 part sulphur to 3 parts lime dissolved in water. It is used primarily to control scale in orchards. Sulphur is sometimes added to Bordeaux mixture when used for orchards.

*Frequency*  
Use rarely and only as a last resort.

*Warning*  
Apply on cool, cloudy days. Do not use on tender leaves as the sulphur will burn the plants. Grapes, gooseberries, cucurbits, squashes and others are particularly sensitive. Sulphur kills a lot of benefi-

cial insects and micro-organisms and increases soil acidity.

## A29 : SUN

*Materials*  
Sunshine and polythene sheeting.

*Targets*  
Maize stalk borer, weeds and many plant diseases.

*Method*  
i The larvae of the stalk borer hibernates during the winter in the stem of the maize plant. It can be killed off by chopping the stover into small lengths and allowing the stalks to thoroughly dry in the sun while spread out as a surface mulch in the field.

ii Plant diseases (and weed seeds) can be killed off by using the heat of the sun while benefiting useful soil micro organisms, and improving soil tilth, nutrition, aeration and water movement. Ensure that the soil surface is smooth and moist to 60cm (70% of field capacity is recommended). Polythene is then stretched firmly over the soil to exclude air spaces. The edges are dug into the soil to prevent wind from moving it. Leave in place for 4-6 weeks. This will ensure that temperatures exceeding 37°C will be reached for extended periods. The treatment is best carried out during the dry season. Black polythene sheeting 1 to 1.5mm thick can be used but clear sheeting is preferable. The sheeting will last two months even if it has no ultraviolet stabiliser in it.

*Frequency*  
Apply treatment i once a year only but inspect the field to ensure that all the stalks have been thoroughly chopped up. Apply treatment ii during the winter on beds which have shown evidence of disease build up.

*Warning*  
When eradicating maize stalk borer, do not leave any standing maize stems as the borer will survive in the base of the rooted stem near to the ground. Cut through the stalk at ground level, thus leaving the root system intact to improve the soil and future infiltration.

## A30 : VINEGAR

*Materials*  
Vinegar, soap, manure and water.

*Targets*  
Aphid.

*Method*  
Spray with a mixture of 3 tablespoons of vinegar, 3 tablespoons of liquid soap, 2 tablespoons of liquid manure and 10 litres of water to kill off aphids and boost the plant at the same time.

*Frequency*  
Apply as often as necessary, usually 2 or 3 times at 4-day intervals.

*Warning*  
No dangers known as the amount of acidic vinegar recommended is too small to damage either the soil or the crop.

## A31: WATER

*Targets*  
Aphids, caterpillars, eggs and larvae, mites, termites, thrips and whitefly; fungal and bacterial diseases particularly damping off; any crawling pest in the nursery.

*Method*  
i Wash insects off with a strong jet of water by applying the thumb over the end of the hosepipe or by a suitable size nozzle on a

- knapsack spray. This treatment will also deter termites by washing away their mud tunnels and increasing the soil moisture.
- ii Wash insects and fungal diseases off the leaves, branches and trunks of any plants with cold water applied with a stiff brush.
  - iii Spray on hot water (40-50°C) over the plant and leaves to control pests. Pour boiling water over the soil surface or blow steam through the soil to control diseases, particularly damping off in nurseries.
  - iv Dip fruit and seeds briefly in hot water to remove bacterial and fungal diseases — see Part IV A5 for further details.
  - v Stand the legs of nursery tables in water. If the legs of the tables are made of wood, first paint them with used engine oil to prevent rot. Add a little oil to the water in the tin to reduce evaporation. Five litre oil tins cut in half make excellent containers for the water.

*Frequency*  
Several applications of i and ii will be necessary within a few days of each other.

*Warning*  
Control the force of the waterjet in i so that the insects are washed off but no harm occurs to the plants. New shoots are particularly vulnerable to strong jets.

For spraying over plants do not have water at higher temperatures or the leaves may be burnt.



## B Plant materials

### GUIDELINES FOR THE USE OF BOTANICAL SPRAYS

Not all is known concerning the use of vegetation to control pests. Much has been forgotten.

Do not assume that the vegetation named will control only the insects named. Try it on others. It may have wider application than originally supposed.

The effectiveness of the various plant remedies vary from place to place, season to season, within season and according to the insect type and maturity and age of the plant. The users should carry out their own experiments to find the plants and remedies most appropriate to their field conditions.

All preparations from plant materials should be used immediately and not exposed to sunlight before use. The best time to apply those with insecticidal properties is in the late evening. They will be more effective against the pest at this time while being less of a danger to beneficial insects such as bees.

Plant leaves can be ground down quickly by rubbing them with wet sand in a bucket.

Preparation time can be greatly reduced by boiling instead of soaking. Boil until water becomes strongly coloured with the juice: 10 minutes for most plants, 30 minutes for tobacco.

There are many plants with insecticidal and repellent properties waiting to be discovered. These can be found simply by observing which plants are not attacked by any particular insect, even though nearby ones of a different type are infested. The untouched plant should be tested for insecticidal or repellency properties.

In the following sections, the first plant name given is the botanical one (in italics for contrast) and then the name in English; names in other languages appear in an appendix at the back of the book.

### B1: USEFUL COMMON CULTIVATED PLANTS

Plants grown in humus-rich soil and fed on compost are known to be the most effective in pest and disease control.

#### B1.1: *ALLIUM CEPA*; ONION

*Description*  
Perennial bulbous herb.

*Action*  
Mildly fungicidal, insecticidal, repellent.

*Targets*  
Aphids, cabbage butterfly, mites, scale, thrips, tomato flies, ticks and whitefly, damping off, late blight and tomato leaf spot, mice and moles.

*Parts*  
Bulb and leaves.

*Application*  
Recipes vary from 10–100g onion peel or leaves per litre of water; leave for 4–7 days in a covered container before spraying. Onions can be planted to repel cabbage butterfly, mice, moles and other pests.

*Other uses*  
Food flavouring, antibiotic, antiseptic.

*Warning*  
The juice irritates the eyes.

#### B1.2: *ALLIUM SATIVUM*; GARLIC

*Description*  
Biennial or annual bulbous herb.

*Action*  
Anti-feedant, bactericidal, fungicidal, insecticidal, nematocidal, repellent.

*Targets*  
Ants, aphids, armyworm, beetles, birds, caterpillars, diamond-back moth, false codling moth, grubs, house flies, mice, mites, moles, mosquitoes, nematodes, peach borer, termites, ticks and animals; fungi and bacteria.

*Parts*  
Bulbs.

*Application*  
Different strengths are required for different pests. A common spray: crush 1 garlic bulb, add to 1 litre of water, mix in a little soap and use immediately. The bulbs can be dried, crushed and used as a dust. The dust can be made into a spray recommended for scab, mildew, bean rust and tomato blight. Garlic planted around fruit trees and other plants will repel aphids, fruit tree borers like peach borer, mice, moles and termites. A mixture made by covering 3 crushed cloves placed in a glass jar (not a tin) with liquid paraffin, soaking for 2 days, filtering and adding 10 litres of soapy water, is said to give a spray which will kill most insects. Garlic bulbs are often planted as a repellent.

*Other uses*  
Food flavouring, antibiotic, worm remedy.

*Warning*  
Garlic is a broad spectrum insecticide which will kill beneficial insects as well as pests. The taste remains on the sprayed or dusted plants for up to 1 month afterwards. Do not use on legumes.

#### B1.3: *CAMELLIA SINENSIS*; TEA

*Description*  
Shrub cultivated commercially for tea leaves.

*Action*  
Anti-feedant, insecticidal.

*Targets*  
Woolly aphid, squash bug, snails and termites.

*Parts*  
Leaves and fruit.

*Application*  
The used leaves can be spread around plants to repel snails; the liquid (tea) when cool can be sprayed on plants; the fruit can be soaked in water and used to repel termites.

*Warning*  
None known.

#### B1.4: *CAPSICUM FRUTESCENS* AND *ANNUUM*; CHILLI, SWEET PEPPERS

*Description*  
Shrubs grown for hot spicy or sweet pods to flavour foods.

*Action*  
Repellent and insecticidal.

*Targets*  
General insects, fungi, bacteria and viruses.

*Parts*  
Ripe pods and seeds.

*Application*  
Grind 2 handfuls of chillies, soak in 1 litre of water for 1 day, shake well for a few minutes, filter, add 5 litres of water and a little soap. Chilli powder can be applied around the base of plants to repel ants, cutworm, slugs, snails and a wide range of soil pests; the juice from sweet peppers will control mosaic virus and inhibit the spread of other viruses. Chillies are often planted as a repellent.

*Other uses*  
Food flavouring, medicine, vegetable (sweet peppers).

*Warning*  
Leaves can be burned if the mixture is too strong.

#### B1.5: *CARICA PAPAYA*; PAWPAW

*Description*  
A 2–10 m tall spindly, short-lived, soft-wooded tree cultivated for its large oval fruit.

*Action*  
Fungicidal, nematocidal and insecticidal.

*Targets*  
Aphids, bugs, caterpillars, cutworm, root-knot nematodes, termites; coffee rust, powdery mildew and rice brown leaf spot.

*Parts*  
Fruit, fresh leaves and roots.

*Application*  
Add 1kg of finely shredded leaves to 1 litre of water, shake vigorously, filter, add 4 litres of water, 2

teaspoons of paraffin and a little soap (20g or ml) spray or water into the soil for cutworm. Extract the juice from immature fruit to control termites.

*Other uses*

Edible fruit and young leaves; medicinal.

*Warning*

None known.

### B1.6: CHRYSANTHEMUM CINERARIIFOLIUM; PYRETHRUM

*Description*

Perennial, daisy-like flowering plant.

*Action*

Broad spectrum insecticide.

*Targets*

Insects in general.

*Parts*

Flowers.

*Application*

Pick flowers on a hot day, dry in shade, grind into powder, dust over insect pests; pour 1 litre of boiling water over 50g pyrethrum flowers (or 20g powder), soak for several hours, add a little soap, filter and spray.

*Other uses*

Medicinal and ornamental.

*Warning*

Apply late in the evening and spray selectively to protect bees etc. Frequency of use is often restricted by organic farming organisations.

### B1.7: CROTALARIA JUNCEA; SUNNHEMP

*Description*

Erect annual shrubby herb.

*Action*

Insecticidal, repellent and trap crop.

*Targets*

Insects in general, pests in stored grain, nematodes and fungi.

*Parts*

Whole plant.

*Application*

Rotate or interplant as a trap crop for nematodes and other pests; crush plant parts in water to make a spray (no details available).

*Other uses*

Fodder, soil improver, weed control (esp. striga), fibre, paper, sacks.

*Warning*

Mildly toxic to cattle under certain conditions; seed should not be stored in room where human beings are working.

### B1.8: ELEUSINE CORACANE; FINGER MILLET

*Description*

Small grain crop.

*Action*

Insecticidal, repellent.

*Targets*

Insects in general especially caterpillars, cutworm, fruit fly, snails.

*Parts*

Residue.

*Application*

Use residues as mulch for soil pests; grow as a trap crop for armyworm; soak residue in water for spraying fruit fly and other insects.

*Other uses*

Food crop, erosion control.

*Warning*

None known.

### B1.9: EUCALYPTUS SPP; GUM TREE

*Description*

Fast-growing, evergreen tree.

*Action*

Repellent, mildly insecticidal.

*Targets*

Insects in general.

*Parts*

Young leaves.

*Application*

Dry and grind into a powder for dusting; crush fresh leaves in water until water is green, add a little soap, spray. The powder mixed with water has been used as a spray to repel stalk borer moths at times when the moths are active.

*Other uses*

Timber, oils.

*Warning*

Heavy water user—do not plant near water or in wet areas. Do not plant too large an area in one location.

### B1.10: GLYCINE MAX; SOYABEAN

*Description*

Widely cultivated field grain crop.

*Action*

Insecticidal.

*Targets*

Ants, woolly aphids and codling moth.

*Parts*

Stems.

*Application*

Spray prepared by soaking stems in water.

*Other uses*

Beans, oil, soil improver.

*Warning*

None known.

### B1.11: IMPOMOEA BATATAS; SWEET POTATO

*Description*

Perennial with trailing ground stems and edible root tuber.

*Action*

Fungicidal, small inert insects.

*Targets*

Aphids; rice brown leaf spot, rice blast fungus; potential to control other fungi.

*Parts*

Leaves.

*Application*

Crush and soak in water, spray; heavily starched water after cooking potatoes should be tried on small immobile insects such as aphid.

*Other uses*

Food.

*Warning*

None known.

### B1.12: LYCOPERSICON ESCULENTUM; TOMATO

*Description*

Perennial soft herb, usually grown as an annual for the fruit; strong scent when leaves crushed.

*Action*

Insecticidal, repellent, attractant, anti-feedant, prevents egg laying, bactericidal, fungicidal.

*Targets*

Aphids, ants, asparagus beetle, cabbage worm, caterpillars, cockroaches, diamond-back moth, flies, grasshoppers, grubs, larvae, mites, nematodes, tomato hornworm, whitefly; fungi in general and bacterial wilt.

*Parts*

Any part of the plant including roots and fruit.

*Application*

Simmer 1kg of chopped leaves in 2 litres of water; shred 2 handfuls of leaves/stems/fruit in 2 litres of water so that the green juice is extracted, leave for 5 hours, filter, add a little soap; spray every 2 days when the butterflies of the cabbage worm are flying. Fresh plant parts are best but should be used immediately. Dried parts can be crushed into a powder and mixed with water to give a spray or applied as a dust but it is not as effective as the fresh material. Tomatoes planted around other plants will protect those plants from asparagus beetles; and the whole plant hung up in orchards or in houses is said to protect the fruit trees from many insects and the houses from cockroaches.

*Other uses*

Edible fruits, oil from seeds.



**Warning**

The leaves are poisonous to humans.

**B1.13: MANIHOT ESCULENTA; CASSAVA (SEE WARNING)****Description**

Short-lived shrub with starch-rich tuber.

**Action**

Nematicidal.

**Targets**

Nematodes; small inert insects.

**Parts**

Roots.

**Application**

Obtain juice by crushing roots, dilute 1:1 with water, spray immediately using 4 litres diluted extract per square metre, said to be very effective. Wait 20 days before sowing; use cassava peelings as a mulch against nematodes; try starchy extract after boiling and cooling cassava on aphids etc.

**Other uses**

Roots of bitter varieties are edible when the prussic acid has been removed by boiling; fodder, paper making, textiles, cosmetics, adhesives, flour.

**Warning**

Roots of bitter varieties contain prussic acid which must be removed before the root is edible.

**B1.14: MATRICARIA EXIMIA; FEVERFEW****Description**

Perennial, shrubby herb with small daisy-like flowers, natural (wild) and cultivated.

**Action**

Insecticidal (pyrethrin), nematicidal.

**Targets**

Caterpillars, flies and soft-bodied insects.

**Parts**

Flowers (especially), leaves.

**Application**

Cover 100g of leaves and flowers with 1 litre of boiling water, soak for 12–24 hours, filter, dilute

with 1 litre of water, add a little soap, spray.

**Other uses**

Food flavouring, medicinal, perfume.

**Warning**

Spot spray to avoid killing beneficial insects; use immediately.

**B1.15: MELIA AZEDARACH; SYRINGA (PERSIAN LILAC)****Description**

Fast-growing ornamental exotic deciduous tree now naturalised.

**Action**

Anti-feedant, contact poison, fungicidal, growth inhibitor, nematicide, insecticidal, repellent.

**Targets**

Wide range of insects including: ants, aphid, armyworm, beetles, bollworms, bugs, cabbage root fly, caterpillars, hoppers, grubs, moles, nematodes, stemborers, weevils, red spider mite, termites and fungi.

**Parts**

Bark, branches, leaves, fruit, oil.

**Application**

Soak 150g of fresh (50g dried) leaves in 1 litre cold water for 24 hours, filter, spray; for termites mulch with leaves and berries or dig them into planting holes; dry and grind plant parts for dusting powder. A handful of leaves and berries boiled in 5 litres of water is said to make a general fungicide and insecticide. The mixture will repel moles if made even stronger.

**Other uses**

Medicinal, fodder, timber, fuel, ornamental.

**Warning**

Effectiveness of the plant has not been tested sufficiently locally; fruit is poisonous.

**B1.16: NERIU INDICUM; OLEANDER (SEE WARNING)****Description**

An ornamental evergreen shrub of gardens.

**Action**

Anti-feedant, fungicidal, nematicidal, insecticidal, rodenticidal.

**Targets**

Ants, flies, beetles, diamond-back moth, rodents, weevils, rice brown leaf spot.

**Parts**

Whole plant.

**Application**

Cut and soak leaves, bark and flowers in water for 30 minutes, spray; dry and grind plant parts into a dusting powder.

**Other uses**

Medicinal, perfumery, incense, nectar, ornamental.

**Warning**

The whole plant is poisonous.

**B1.17: NICOTIANA TABACUM; TOBACCO (SEE WARNING)****Description**

Erect annual herb; important cash crop.

**Action**

Contact and stomach poison, fungicidal, insecticidal, molluscicidal, repellent, viricidal.

**Targets**

Broad spectrum insecticide including: aphids, bilharzia snails, fleas, flea beetles, lawn caterpillar, leaf curl, leaf miner, scale, slugs, snails, termites, thrips, ticks, whitefly and many others; rusts, viruses.

**Parts**

Whole plant.

**Application**

Soak 1kg of bruised leaves in 15 litres of water for 24 hours, add a little soap, filter, use as a general spray; dry and grind very finely into a powder for dusting onto aphids, spider mites, slugs, snails, thrips, and various caterpillars; dust on trees and crops to repel insects and to control leaf curl virus; mulch with scrap leaves to kill slugs and snails, pupae, larvae or any adults hiding in the soil or in other mulch material and to repel aphids, flea beetles and thrips.

**Other uses**

Cash crop, medicinal.

**Warning**

Nicotene is a strong poison, avoid contact with the liquid forms; wear protective clothing if spraying otherwise use a watering can; liquids damage plants if used in higher concentrations than recommended. Do not spray or dust on leaves of the Solanaceae family — tomatoes, potatoes, brinjals and peppers. Do not use tobacco in any form on roses or they will turn black. Liquid sprays made from tobacco are not acceptable to The Zimbabwe Organic Producers' Association (ZOPA) and use of powdered material is restricted by some organic farming organisations.

**B1.18: OCIMUM BASILICUM; SWEET BASIL****Description**

Cultivated, aromatic herb.

**Action**

Mildly fungicidal, insecticidal, repellent.

**Targets**

Wide range of insects, fungi in general.

**Parts**

Whole plant.

**Application**

Soak crushed leaves in water for 24 hours, filter, spray; dry out of sun and grind as a dusting powder.

**Other uses**

Food flavouring, medicinal, repels mosquitoes and ticks.

**Warning**

The effectiveness seems to vary greatly with variety and other factors.

**B1.19: RHEUM SPP; RHUBARB (SEE WARNING)****Description:**

Perennial, large-leaved herb.

**Action**

Fungicidal, insecticidal.

**Targets**

Ants, aphids, whitefly, small caterpillars, maggots and other soft-bodied insects; club root and mildew.

**Parts**

Whole plant.

**Application**

Soak 100g fresh leaves in 1 litre of water for 24 hours, add a little soap, spray; up to 5 large leaves in 0.5 litres of water have been used to give stronger sprays; stems dug into the soil next to cabbages prevents club root.

**Other uses**

Medicinal, stems edible.

**Warning**

Leaves are extremely poisonous.

**B1.20: TAGETES ERECTA; AFRICAN MARIGOLD****Description**

Annual, erect, cultivated herb.

**Action**

Bactericidal, contact poison, fungicidal, insecticidal, nematocidal, repellent.

**Targets**

Wide range of insects including ants, beetles, nematodes and fungi; late blight.

**Parts**

Whole plant.

**Application**

Crush 100–200g of leaves, roots, flowers, pour on 1 litre boiling water, soak for 24 hours, add 1 litre of cold water, spray on plants or into soil; grow in rotation for nematodes; plant as an intercrop to repel beetles.

**Other uses**

Ornamental, erosion control, medicinal, food and cloth dye.

**Warning**

None known.

**B2: USEFUL WILD PLANTS AND TREES**

Among the following useful plants, the trees can be planted around homes, at the edges of gardens and fields, and on the contour ridges. Not only will they provide a ready supply of materials for pest and disease control but act as windbreaks and encourage predators to do most of the pest-control work.

**B2.1: AGAVE AMERICANA; AMERICAN ALOE AND OTHER AGAVE SPP.****Description**

Stout, succulent plant growing to 7 metres.

**Action**

Repellent and insecticidal.

**Targets**

Pests in stored grain and termites; buck; worth trying on other field pests.

**Parts**

Any part of the plant.

**Application**

Crush plant parts in water as field spray, 1 part plant to 5 parts water or less; dry and grind materials for dusting; cut up leaves, boil, cool and spray over plants to repel buck.

**Other uses**

Planted as a fence, edible buds and flowers, fibre, alcohol and medicine.

**Warning**

None known.

**B2.2: AGERATUM CONYZOIDES; GOAT WEED****Description**

Erect, perennial or annual herb growing 1 metre high.

**Action**

Anti-feedant, bactericidal, insecticidal, nematocidal, and repellent.

**Targets**

Bacteria, cotton stainer, flies, vinegar fly, maize weevil; beetles in stored flour, groundnuts etc. and nematodes.

**Parts**

Any part of the plant.

**Application**

Dig the plant into the soil or use as mulch for nematodes; crush in water as a spray; dry and grind as a dust; can be used very dilute; effective for 2 weeks.

**Other uses**

Medicinal (antibiotic), weed control, ground cover in plantations.

**Warning**

None known.

**B2.3: ANNONA MURICATA, RETICULATA AND SQUAMOSA; SUGAR APPLE (SEE WARNING)****Description**

Widely distributed small trees with edible fruit.

**Action**

Insecticidal.

**Targets**

General insects, nematodes, head lice.

**Parts**

Roots, leaves, unripe fruit, seeds and bark (A. reticulata).

**Application**

The above parts can be soaked in water for a few days and the liquid used as a spray. The seeds can be dried and crushed and dusted directly on to the insects or diluted with 20 parts of water as a spray. The power of the extract can be increased by soaking crushed seeds in a few drops of paraffin or a little petrol for a few hours. The seeds of Annona squamosa have been crushed in water as a spray for aphids and other insects.

**Other uses**

Edible fruit, medicine.

**Warning**

The powder made from the seeds of Annona squamosa is said to cause great pain if it gets into the eyes.

**B2.4: ANNONA SENEGALENSIS; CUSTARD APPLE****Description**

Widely distributed shrub.

**Action**

Insecticidal.

**Targets**

General insects including aphids, diamond-back moth, grasshoppers and leaf hoppers.

**Parts**

Bark and seeds.

**Application**

Soak bark in water for several days and spray; crush dried seeds to produce a dusting powder; a stronger spray can be made by soaking the crushed seeds in a few drops of paraffin for a few hours before adding the water.

**Other uses**

Edible fruit.

**Warning**

None known.

**B2.5: ARGEMONE MEXICANA; MEXICAN POPPY****Description**

Tall, annual weed with red, flowering head.

**Action**

Bactericidal, fungicidal, insecticidal, nematocidal, repellent.

**Targets**

Ants, beetles, bugs, caterpillars, grubs, nematodes, termites, weevils; and fungi.

**Parts**

All plant parts.

**Application**

Plant parts can be soaked in water to make a spray or dried and ground for dusting.

**Other uses**

Medicinal, oil from seeds, soap.

**Warning**

None known.



**B2.6: BALANITES AEGYPTIACA;**  
DESERT DATE

*Description*  
Evergreen shrubby tree.

*Action*  
Insecticidal, molluscicidal.

*Targets*  
Melon worm, black carpet beetle, diamond-back moth, slugs, snails, bilharzia.

*Parts*  
Bark, fruits, roots, seeds.

*Application*  
Crush the plant parts in water to form a soapy liquid, spray; the dried seeds can be crushed to give a powder for dusting on to slugs and snails. The fruits are mentioned particularly as a very effective control of bilharzia.

*Other uses*  
Fibre, medicinal, soap.

*Warning*  
None known.

**B2.7: BERSAMA ABYSSINICA; BITTER BARK**

*Description*  
Perennial, evergreen tree of the Eastern Districts.

*Action*  
Insecticidal.

*Targets*  
Insects in general.

*Parts*  
Roots and leaves.

*Application*  
Crush leaves and roots, cover with water, soak for 24 hours, filter, dilute with water 10 times, add a little soap, spray; grind dried roots and leaves into a dusting powder.

*Other uses*  
Medicinal.

*Warning*  
Spot spray — will kill off beneficial insects.

**B2.8: BIDENS PILOSA; BLACKJACK**

*Description*  
Annual weed with clusters of spiky black seeds.

*Action*  
Anti-feedant, insecticidal, repellent.

*Targets*  
Aphids, ants, beetles, cabbage root fly, caterpillars, crickets, mites, termites, whitefly.

*Parts*  
All the plant but particularly the mature seeds.

*Application*  
Cover a cupful of mature seeds with water, boil for 10 minutes (or soak for 24 hours), cool, add 1 litre of water and a teaspoonful of soap, spray immediately. Seeds can be spread around bushes to deter termites. The whole plant can be crushed/hand-rubbed in water until the juice has come out, and used as a spray.

*Other uses*  
Edible young leaves.

*Warning*  
High concentrations may affect the flowers of some plants.

**B2.9: CATUNAREGAM SPINOSA**

*Description*  
Perennial, shrubby tree.

*Action*  
Anti-feedant, insecticidal, repellent.

*Targets*  
Aphids, diamond-back moth, grasshoppers, hairy caterpillar, scale, stored grain pests, termites.

*Parts*  
Roots, fruit.

*Application*  
Pound the fruit and bulbs together with water into a concentrated pulp, dilute and spray.

*Other uses*  
Edible fruit.

*Warning*  
None known.

**B2.10: CHENOPODIUM**  
AMBROSIOIDES; WORMSEED

*Description*  
Weakly perennial, aromatic herb up to 1 metre high.

*Action*  
Anti-feedant, insecticidal, nematocidal, repellent.

*Targets*  
Ants, beetles, larvae, mosquitoes, termites, stored grain pests.

*Parts*  
All plant parts.

*Application*  
Plant around buildings to repel termites and storage pests; mix with water as a spray.

*Warning*  
None known.

**B2.11: DATURA STRAMONIUM;**  
THORN APPLE, JIMSON WEED (SEE WARNING)

*Description*  
Tall, annual weed with angular leaves.

*Action*  
Anti-feedant, fungicidal, insecticidal, nematocidal.

*Targets*  
Insects in general, cutworm, nematodes and some fungal diseases.

*Parts*  
Whole plant.

*Application*  
Dry whole plant out of the sun, grind and use as a powder; crush a handful of leaves in 1 litre of water as a spray, add a little soap.

*Other uses*  
Medicinal.

*Warning*  
Plant is poisonous so do not put near the mouth.

**B2.12: EUPHORBIA HIRTA; ASTHMA WEED**

*Description*  
Short, annual weed with milky sap.

*Action*  
Fungicidal, insecticidal, viricidal.

*Targets*  
Some funguses; insects and viruses in general.

*Parts*  
Leaves.

*Application*  
Crush leaves in water and spray.

*Other uses*  
Medicinal.

*Warning*  
None known.

**B2.13: EUPHORBIA TIRUCALLI;**  
RUBBER HEDGE INCLUDING  
EUPHORBIA OR CANDELABRA (SEE WARNING)

*Description*  
Perennial, succulent shrubs or trees.

*Action*  
Fungicidal, insecticidal, repellent.

*Targets*  
Aphids, mosquitoes, red spider mite, termites, fungi and insects in general.

*Parts*  
Whole plant.

*Application*  
To repel termites apply finely chopped leaves to planting holes or place the branches around trees. As a general spray mix 10 drops of the milky sap thoroughly into 1 litre of water, the water should be slightly milky, spray.

*Other uses*  
Provides a protective hedge around gardens, fire break, medicinal.

**Warning**

The milky sap is poisonous, keep away from skin and eyes, do not spray on windy days.

**B2.14: JATROPHA CURCAS; BARBADOS NUT (SEE WARNING)****Description**

Shrub or tree, cultivated and wild.

**Action**

Insecticidal, molluscicidal.

**Targets**

Leaf-eating larvae, snails.

**Parts**

All parts.

**Application**

Soak plant parts in water.

**Other uses**

Medicine, soap, live fence.

**Warning**

Seeds are poisonous.

**B2.15: LANTANA CAMARA; LANTANA****Description**

Invasive, perennial shrub.

**Action**

Contact poison.

**Target**

Wide range of insects.

**Parts**

Leaves, flowers, branches.

**Application**

Crush 1 handful of leaves in 1 litre of water, add a little soap, spray; dust with the ashes; dry and grind into a dusting powder. Burn the branches and dust the ash over beetles and leaf miner.

**Other uses**

Medicinal.

**Warning**

Classified as a noxious weed in Zimbabwe and poisonous to cattle.

**B2.16: LEONOTIS NEPETIFOLIA; LION'S EAR****Description**

Hairy, annual, shrubby plant with square stems.

**Action**

Insecticidal, stomach poison.

**Targets**

Ants, bean leaf beetle, cotton stainer, melon worm, termites.

**Parts**

Stem, branches, seeds.

**Application**

Dry plant and grind to produce a powder.

**Other uses**

Leaves put into wounds to expel maggots.

**Warning**

None known.

**B2.17: LIPPIA JAVANICA; LIPPIA****Description**

Small, perennial shrub.

**Action**

Insecticidal, repellent.

**Targets**

Ants, aphid, caterpillars, termites, whitefly and others.

**Parts**

Leaves.

**Application**

Crush leaves and branches in water, stand for 2-3 days, add a little soap, spray; burn dried leaves and branches in the field to repel insects.

**Other uses**

Leaves in hot water for stomach problems.

**Warning**

None known.

**B2.18: LOBELIA COLUMNARIS; GOD'S TOBACCO****Description**

Shrubby herb 2-3 metres high.

**Action**

Insecticidal.

**Targets**

Aphids, caterpillars, stemborers, bugs and others.

**Parts**

Leaves.

**Application**

Crush 1 kg of leaves and soak in 10 litres of water for 1 day, add a little soap and lime, spray.

**Other uses**

None known.

**Warning**

None known

**B2.19: LONCHOCARPUS CAPASSA; RAIN TREE — L. BUSSEI, ERIOCALYX AND NELSII KNOWN AS LANCE-PODS****Description**

Perennial, deciduous trees.

**Action**

Insecticidal (rotenone).

**Targets**

Wide range of insects.

**Parts**

Roots.

**Application**

Dry and grind roots for dusting; crush 500g of roots, cover with 1 litre of boiling water, soak for 2 days, filter, add 5 litres of water, spray (do not use soap).

**Other uses**

Fodder, bee tree, timber, soil improver.

**Warning**

Spot spray to avoid spraying beneficial insects. Use of rotenone products is restricted by some organic farming organisations.

**B2.20: MUNDULEA SERICEA; CORK BUSH****Description**

Small tree or shrub with corky bark.

**Action**

Insecticidal (rotenone).

**Targets**

Wide range of insects; red spider mite specified.

**Parts**

Roots, bark, seeds.

**Application**

Boil water, remove from fire, immerse equal volume of finely crushed bark immediately, stand for 2-3 days, filter, dilute with 5 parts water, spray (do not add soap); grind dried plant parts for dusting powder.

**Other uses**

Soil improver.

**Warning**

Spot spray — avoid killing beneficial insects. Use of rotenone products is restricted by some organic farming organisations.

**B2.21: NICANDRA PHYSALODES; APPLE OF RU****Description**

A common weed of cultivated or disturbed ground.

**Action**

Anti-feedant, insecticidal, repellent.

**Targets**

Flies, whitefly, Mexican bean beetle, tomato hornworm.

**Parts**

Whole plant.

**Application**

A companion plant; place bruised leaves in area to be protected; crush plant in water and spray.

**Other uses**

None known.

**Warning**

None known.



**B2.22: NICOTIANA RUSTICA; WILD TOBACCO** (SEE WARNING).**Description**

Erect, annual herb; cultivated in small quantities.

**Action**

Contact and stomach poison, fungicidal, insecticidal, molluscicidal, repellent, viricidal.

**Targets**

Wide spectrum insecticide, rusts, leaf curl virus, ticks, slugs, snails, fleas.

**Parts**

Whole plant.

**Application**

Soak 1kg of bruised leaves in 15 litres of water for 24 hours, add a little soap, filter, spray; dry and grind into a powder for dusting to kill wet-bodied pests like slugs or snails; dust on trees and crops to repel insects and to cure leaf curl virus; mulch with scrap leaves.

**Other uses**

Dust repels ticks and fleas; medicinal.

**Warning**

Nicotine is a poison, avoid contact with the liquid forms; wear protective clothing if spraying otherwise use a watering can; liquids damage plants if used in higher concentrations than recommended.

**B2.23: OCIMUM CANUM AND URTICAFOLIUM; BASIL; HOARY AND WILD****Description**

Aromatic, shrubby herb.

**Action**

Mildly fungicidal, insecticidal, repellent.

**Targets**

Aphid, flies, hairy caterpillar, maggots, mosquitoes, pests on seedbeds, potato tuber moth, spider mites.

**Parts**

Whole plant.

**Application**

Soak crushed leaves in water for 24 hours, filter, spray. This infusion is used locally to kill pests in seedbeds.

**Other uses**

Medicinal, food flavouring, mosquito and fly repellent.

**Warning**

None known.

**B2.24: RICINUS COMMUNIS; CASTOR OIL PLANT** (SEE WARNING)**Description**

Fast-growing shrub or tree.

**Action**

Contact poison, fungicidal, insecticidal, nematocidal, repellent.

**Targets**

Aphids, leaf cutting ants, caterpillars, cutworm, mites, stink/shield bugs, termites, nematodes in general; fleas, lice, moles; fungi: anthracnose, brown patch, damping off, root rot.

**Parts**

Leaves and seeds mainly.

**Application**

As a general spray soak green seeds and leaves in water for 24 hours, filter and spray; dry green seeds and leaves and grind for dusting powder; for cutworm put 4 cups of crushed shelled seeds in 2 litres of water, boil for 10 minutes, add 2 teaspoons of paraffin and some soap, dilute to 10 litres and water immediately into the soil; put green seeds into mole holes or rat runs as effective repellent; dig seeds, leaves or oil cake into the soil to kill fungal diseases; mulch with branches and leaves to repel termites; varieties with red stems said to be more effective than plants with green stems.

**Other uses**

Vegetable oil, medicinal, soap, industrial uses.

**Warning**

Seed is poisonous to humans and fowl.

**B2.25: TAGETES MINUTA; MEXICAN MARIGOLD****Description**

Strong-smelling weed of cropland and disturbed ground.

**Action**

Fungicidal, insecticidal, nematocidal, repellent.

**Targets**

Ants, aphids, blowflies, caterpillars, flies, fleas, maggots, mosquitoes, termites, nematodes; prevention of coffee berry disease, late blight, mildew and others.

**Parts**

Whole plant.

**Application**

Soak crushed parts of 1 mature plant in 2 litres of water for 24 hours, filter and spray; a handful of wood ash can be added to improve the insect range; if plants are in flower and the soaking prolonged for 5-10 days with daily stirring, the spray is said to be more effective; dig fresh *Tagetes* plants into the soil around plants to be protected from ant attack; apply weekly for prevention of fungal diseases; grow in rotation for nematode control; dry and grind for dusting powder against fleas etc. Leaves rubbed on the skin will repel mosquitoes. Ants can be repelled either by watering the liquid from the crushed plants well into the soil around the plants to be protected or by digging the freshly crushed leaves of *Tagetes* into the soil before planting.

**Other uses**

Medicinal, soil improver.

**Warning**

The aroma makes bees aggressive so do not approach bees when smelling of this plant.

**B2.26: TEPHROSIA VOGELII; TEPHROSIA****Description**

Perennial shrub with mauve flowers and many pods.

**Action**

Anti-feedant, insecticidal (rotenone), repellent.

**Targets**

A wide range of insects, including leaf miner, shield bugs and cattle ticks.

**Parts**

Leaves.

**Application**

Crush 50 fresh leaflets in 1 litre of water, stand for 24 hours, do not use soap; it is said to be the most effective against aphids if the water is warm when spray applied.

**Other uses**

Medicinal, soil improver.

**Warning**

Spot spray to avoid killing beneficial insects. Use of rotenone products is restricted by some organic farming organisations.

**B3: SOME EASY REMEDIES FROM PLANT MIXTURES**

These have general insecticidal properties and, like all plant mixtures, should be used immediately.

**B3.1: CHILLIS AND GARLIC**

Stir 1 thoroughly crushed or finely grated garlic bulb with 1 teaspoon of powdered chilli peppers into 2 litres of hot water, cool, filter, add a little soft soap and stir well.

Spray for caterpillars in fruit trees.

**B3.2: CHILLIS, GARLIC AND ONIONS**

Chop or grind 1 garlic bulb, 1 onion, 1 tablespoon of hot pepper and mix with 1 litre water, stand for 1 hour, add 1 tablespoon of liquid soap.

Spray for insects in general.

**B3.3: CHILLIS, MEXICAN MARIGOLD AND ONIONS**

Chop up 4 chillis, 4 onions and a handful of Mexican marigold leaves. Cover with soapy water, soak for 24 hours, strain and add 2 litres of water.

This spray is recommended particularly for red spider mite.

**B3.4: CHILLIS, GARLIC, MEXICAN MARIGOLD AND ONIONS**

Take 3 large cloves of garlic, 2 handfuls of marigold leaves, 2 large onions, 2 large peppers, add water and bring to the boil, cool, dilute with 4 times the quantity of water, stir constantly.

Spray for insects in general.

**B3.5: GARLIC AND TEPHROSIA VOGELII**

Pound the fresh leaves of Tephrosia and dried leaves of garlic into a paste with a small amount of water, mix 50g of the paste with 1 litre of water and filter. Apply weekly from 1 week after emergence or transplanting until 2 weeks before harvesting.

Water the ground around plant for cabbage worm at the rate of 1 litre of liquid per square metre.

**B3.6: MEXICAN MARIGOLD (TAGETES MINUTA) AND SYRINGA**

Soak the leaves of Mexican marigold with syringa leaves and berries in water for a few days; dilute the liquid to a pale green or brown; spray against ants, cabbage root fly and mites.

**B4: More Plants with Potential**

When searching for new remedies, select from amongst a group of plants being attacked one that is not attacked by a particular pest or pests, or one that has little damage. Blend or grind up some of the leaves and cover with water, soak for a few hours, strain, add 4 parts of water, spray on the infested plants. Try different concentrations on small test areas if it looks promising.

Any aromatic plant is likely to have potential in pest control.

The information under each identified plant is given in the order Description/target insect/parts used/other uses.

**B4.1: ALBIZIA ANTHELMINTICA**

Shrub/termites, fleas, mites on chickens/roots and leaves/soil improver.

**B4.2: ALBIZIA LEBBEK; WOMAN'S TONGUE**

Tree/flies, larvae, mosquitoes/leaves/soil improver.

**B4.3: ALEURITES FORDII; TUNG TREE**

Annual tree/weevils, stinkbugs, woolly aphid/fruits/food flavouring, tools, building material, dye, ink, oil.

**B4.4: AMARANTHUS SPINOSUS; PIGWEED**

Annual weed/fungi/leaves/edible leaves and grain,

**B4.5: ANDROPOGON SPP; BLUE, SNOWFLAKE AND STAB GRASS**

Perennial grass/insects/leaves/forage, thatching, erosion control.

**B4.6: CALOTROPIS PROCERA; SWALLOW WORT**

Annual shrubby weed/red pumpkin beetle, cabbage white butterfly, termites/whole plant/medicine, fibre, weaving, resin products,

**B4.7: CHENOPODIUM ALBUM; FAT HEN**

Annual herb 0.7m/insects/whole plant/edible leaves and meristem tissue.

**B4.8: CLAUSENA ANISATA**

Perennial shrub/armyworm/leaves/medicinal.

**B4.9: COMBRETUM IMBERBE; LEADWOOD**

Small tree/insects/ash or boiled leaves plus sugar/timber, fuel, medicine.

**B4.10: DICHROSTACHYS CINEREA; CHINESE LANTERN**

Spreading shrub/insects, nematodes/roots, leaves/soil improver, rope, wood, tannin.

**B4.11: DIGITARIA TERNATA; FINGER GRASS AND OTHER SPP.**

Tufted annual grass/termites/planted/fodder.

**B4.12: HETEROPOGON CONTORTUS; SPEAR GRASS**

Perennial tufted grass/nematodes/planted, leaves watered into the soil/forage, land reclamation.

**B4.13: KHAYA NYASICA; RED MAHOGANY**

Large tree/insects, lice, rats, termites/soaked seeds/hardwood.

**B4.14: MYROTHAMNUS FLABELLIFOLIUS; RESURRECTION BUSH**

Small shrub/insects/leaves and branches/medicinal.

**B4.15: SECURINEGA VIROSA; SNOW BERRY**

Deciduous shrub/insects, fungi/bark, leaves/edible fruit.

**B4.16: SOLANUM PANDURIFORME; SODOM APPLE**

Shrub/ants, cutworm, cockroaches, termites/fruit juice and slices /medicinal.

**B4.17: STRYCHNOS SPP; MONKEY OR WILD ORANGE**

Small tree/insects/fruits, seeds/edible ripe fruit.

**B4.18: TITHONIA ROTUNDIFLORA; RED SUNFLOWER**

Common weed/insects/leaves/ornamental.

**B5: SUMMARY OF PLANTS WITH INSECTICIDAL PROPERTIES**

The following plants have been identified as having the ability to kill a wide range of insects. They are listed by their common English names followed (in brackets) by the reference number by which they can be looked up in Sections B 1-4.

African Marigold (1.20);	Gum tree (1.9);	Snow berry (4.15);
American aloe (2.1);	Lantana (2.15);	Soyabean (1.10);
Apple of Peru (2.21);	Leadwood (4.9);	Stab grass (4.5);
Asthma weed (2.12);	Lion's ear (2.16);	Sugar apple (2.3);
Barbados nut (2.14);	Lippia (2.17);	Sunn hemp (1.7);
Bitter bark (2.7);	Mexican marigold (2.25);	Sweet basil (1.18);
Blackjack (2.8);	Mexican poppy (2.5);	Sweet potato (1.11);
Cassava (1.13);	Monkey orange (4.17);	Syringa (1.15);
Castor oil (2.24);	Oleander (1.16);	Tea (1.3);
Catunaregam (2.9);	Onion (1.1);	Tephrosia (2.26);
Chilli (1.4);	Pawpaw (1.5);	Thorn apple (2.11);
Chinese lantern (4.10);	Plant mixtures (3.1 to 3.6);	Tobacco (1.17);
Cork bush (2.20);	Pyrethrum (1.6);	Tomato (1.12);
Custard apple (2.4);	Rain tree (2.19);	Wild basil (2.23);
Fat hen (4.7);	Red mahogany (4.13);	Wild tobacco (2.22);
Feverfew (1.14);	Red sunflower (4.18);	
Finger millet (1.8);	Resurrection bush (4.14);	
Garlic (1.2);	Rhubarb (1.19);	
Goat weed (2.2);	Rubber hedge (2.13);	



## B6: SUMMARY OF PLANTS WITH REPELLANT PROPERTIES

The following plants have been identified in the text as having the ability to repel insects. Some of them act also as general insecticides. The plants are listed by their common English names followed (in brackets) by the reference number by which they can be looked up in Sections B 1-4.

African marigold (1.20);	Goat weed (2.2);	Sweet basil (1.18);
American aloe (2.1);	Gum tree (1.9);	Syringa (1.15);
Apple of Peru (2.21);	Lippia (2.17);	Tephrosia (2.26);
Blackjack (2.8);	Mexican marigold (2.25);	Tobacco (1.17);
Castor oil (2.24);	Mexican poppy (2.5);	Tomato (1.12);
Cannaregan (2.9);	Onion (1.1);	Wild basil (2.23);
Chilli (1.4);	Plant mixtures (3.1 to 3.6);	Wild tobacco (2.22);
Finger millet (1.8);	Rubber hedge (2.13);	Wormseed (2.10);
Garlic (1.2);	Sunn hemp (1.7);	

## B7: LATE ENTRIES

### B7.1: CUCUMIS ANGURIA; WILD GHERKIN

#### Description

An annual vine, a member of the pumpkin family, usually growing on heavily manured soils around cattle pens.

#### Action

Insecticidal.

#### Targets

Aphids.

#### Parts

Ripe fruit (yellow) which are small balls with spikes.

#### Application

The ripe fruit is crushed, mixed with water and allowed to stand for 2 days in a covered container. When the solution has thus fermented, it is sprayed on the aphids.

#### Other uses

Edible leaves.

#### Warning

None known

### B7.2: SPIROSTACHYS AFRICANA; AFRICAN SANDALWOOD (SEE WARNING)

#### Description

A medium-size indigenous tree with a rounded crown.

#### Action

Repellant, insecticidal.

#### Targets

Aphids, cutworms, grasshoppers, weevils.

#### Parts

Branches.

#### Application

The ashes are applied in the planting holes to control cutworm or dusted over the crop leaves for aphids and grasshoppers. The ash is mixed with cow manure and plastered on the walls and floor of granaries to repel weevils.

#### Other uses

Furniture, termite and borer resistant. Fish and game poison.

#### Warning

The milky latex is very poisonous and should not be in contact with the skin or eyes; protect the eyes from the sawdust.

## C Baits, Barriers, Scarecrows and Traps

### C1: ARMYWORM AND OTHER MARCHING CATERpillARS

Dig or plough a trench about 60cm wide and 45cm deep along the side of the garden or field. The sides of the trench should be vertical. The caterpillars fall into the trench and are trapped. They can be destroyed by filling the trench with straw or similar materials and setting fire to it.

In an emergency a single deep plough furrow can halt the swarm long enough for the caterpillars to be killed off.

### C2: BANANA WEEVIL

When the pest is first detected, pieces of banana stem 50cm long can be cut and split in two to form traps. Lay the pieces on the ground with the cut side facing down. The weevils bore their way into the chopped sections in order to eat the sap. After a few days the traps are collected, burnt and fresh traps prepared.

### C3: BEETLES, BUTTERFLIES, GRASSHOPPERS AND MOTHS

Flying or hopping insects can easily be trapped in hand nets made of fine mesh rather like small fishing nets. This is a suitable task for children. It can be made into a fun game with prizes for the one who collects the most. Each person should be given a jar, tin or plastic container to hold their catch.

### C4: BEETLES

Certain beetles are attracted by bright yellow containers. Part fill the container with soapy water and place near the crop. A little sugar or honey added to the water is said to increase the attraction. The beetles fall in and drown. Some green beetles are said to be attracted to white containers and blister beetles to blue and red containers — see C6.

### C5: BIRDS, BABOONS AND BUCK

Light nets, shade cloth and fine wire netting placed over the crop will protect planted seeds and the later

harvest from birds. Lines of string hung over the crop between tall poles will also act as a deterrent.

Pieces of white cloth, plastic or paper bags containing a few dried beans to make a rattle, tied every 3-5m along a white string so that they flap in the wind will keep away the more timid birds.

Scarecrows should be made to look like humans, snakes or birds of prey should be put in place *before* the fruit ripens or else the birds will take little notice of them. Scarecrows with flapping clothes are said to be the most effective. When the effectiveness of one type of scarecrow wears off, replace it with another one. Scarecrows which move in the wind are best, particularly if old smelly clothes are used and replaced with further unwashed clothing at regular intervals. The smelly-clothing scarecrow is effective against buck and baboons. See *Noise* under Section A20.

Although not related to traps etc, the damage caused to crops by birds can be avoided simply by encouraging the presence of birds of prey and by not destroying (through overgrazing, burning and tree felling) their natural food supplies.


Providing birds with bird baths is said to reduce the damage to moist green shoots and fruits to which the birds are attracted when they are thirsty.

### C6: BLISTER BEETLES

- i Take any blue container, fill with soapy water and place around susceptible crops like beans and cowpeas. The beetles fall in and drown.
- ii Dig holes, line them with blue plastic bags and place soapy water in the bottom.
- iii The beetle is said to be attracted by other bright colours too. Red has been named as a possible.

### C7: CATERpillARS, GRUBS AND SLUGS


Bind a 300mm wide piece of wire mosquito gauze around the trunk of the fruit tree. Nail the gauze tightly to the trunk at the top to prevent insects



getting under it (seal any gaps with clay and cement or clay with builders' lime). Support the bottom end so that it stands out from the trunk (long nails can be used). Crawling grubs can enter the funnel-shaped trap at the wide bottom end and are trapped by the tightly closed top end.

### C8: CUTWORM, CABBAGE ROOT FLY AND CRICKETS



- i Mix together equal parts of hardwood sawdust or bran, a good portion of molasses and enough water to make the mixture sticky. Distribute a handful of this preparation over the endangered crop at evening twilight. The molasses attracts the cutworms, sticks to their bodies and hardens towards the morning. The cutworm cannot enter the soil and can be hand picked or eaten by birds.
- ii Mix 100g bran, 10g sugar, 10cc pyrethrum powder, 200cc water thoroughly together and put near endangered plants. This bait is recommended for the control of cutworm, crickets, dusty surface beetle and false wireworm. Apply to crops just after emergence for cutworm and crickets and in February for dusty surface beetle. Baits can also be made using tobacco dust and rotenone from the Cork bush and Rain tree.
- iii Tin cans placed over the seedlings as collars will prevent cutworm and cabbage root fly from reaching the plant. Press the can firmly into the soil so that the cutworm cannot crawl under it. If the cutworm is already inside the tin it should be dug out by hand.



- iv Press two thin sticks into the soil on each side of the plant. The sticks should touch the stem of the plant so the cutworm will not squeeze between the sticks and the plant stem. It is said the cutworm cannot cut through the stem unless its body can encircle it. If the sticks used are hard and aromatic like the stems of Mexican marigold, the insect will be repelled.
- v Put scraps of old lettuce or cabbage leaves on the soil surface when the bed is weed free. Lift the leaves early in the morning and pick up cutworm hiding underneath.

### C9: FLIES AND FRUITFLIES

- i Half fill a narrow-necked bottle with water, fermenting fruit juice or water smelling of rotting meat or fish, smear a little honey or sweetened water around the neck to attract the flies. The top of a plastic bottle can be cut off and inverted to give a good fly trap as shown in the sketch.



- An ordinary jar with holes in the lid gives another very effective trap.
- ii A bait of stale beer and fermented sugar placed in a tin in a box will attract flies. Throw a cloth or net over the box to trap the flies and remove or destroy them.

### C10: LEAF MINER

Yellow painted boards of 30cm by 30cm, coated with oil or any other sticky substance, erected vertically at a height of 60cm above the ground have been found to trap the flies.

### C11: LOCUSTS

See *Noise* under Section A.

### C12: MOTHS

Fruit tree moths can be trapped by putting 25mm of brown sugar and syrup or molasses in the bottom of a jar and hanging it with string in the fruit tree. Cover the top of the jar with 6mm mesh to exclude bees. See also C17 for light traps.

### C13: RATS, MICE, MOLES, RABBITS AND HARES

- i Where there is a shortage of surface water, rats and mice can be trapped in drums dug into the ground and part-filled with water. The effectiveness of the above trap can be increased by throwing a few peanuts into the water and by smearing a generous ring of peanut butter around the inside of the drum at a depth of 3cm. When rats and mice try to eat

- ii this they overbalance and fall into the water. Where water is short, a steep-sided pit can be dug into the soil about 1m deep. A stick is then laid so that one end reaches to the centre of the pit. A suitable bait is put on the end. The stick is balanced so that the rodent's weight tips it and the rodent falls into the hole.
- iii See *Noise A20* for moles.
- iv A tarred or creosoted rope around the plot is said to give protection from rabbits and hares.
- v Another means of trapping moles is to go out in the early morning when they can be seen clearing out their burrows. Dig a spade or hoe into the ground behind the mole and dig it out.
- vi Vetiver or lemon grass planted as barriers prevent moles from entering the area.
- vii Old seeds spread on the soil surface will act as a decoy to protect the sown seed against rats and mice digging them up. Alternatively, fill the gaps with seedlings 2-3 weeks old from the nursery.
- viii Scarecrows made to look like snakes or birds of prey are said to keep rats and mice away.

### C14: SLUGS, SNAILS AND SQUASH BUGS

- i Sink a suitable shallow container like a largish tin lid into the earth to ground level and fill with equal parts milk and water, or beer and/or maize meal; sawdust, sliced turnip or lettuce and cabbage leaves are all good baits to attract slugs and snails. Empty every few days. The left-over milk can be spread on plants susceptible to fungal diseases, snails can be fed to chickens, slugs burnt and the ash used to fertilise heavy feeding vegetables.
- ii Deepertins and bottles (with the tops knocked off) can be used if they are dug into the ground so that their tops are level with the soil surface.
- iii Place a dish upside down between plants and then keep the soil under it moist. Remove the slugs and snails every morning. A bait may or may not be present.
- iv Slugs and snails will congregate under boards or wet sacks laid on the soil surface. Half grapefruits with some of the pulp left inside



C16, C17

(open end downwards) are laid to make effective slug traps.

- v Place slightly tilted boards near the plants as traps for squash bugs which hide underneath.

### C15: WHITEFLY


Yellow-orange painted boards of 20cm by 35cm coated with oil or other adhesive substances can be used as traps and have been particularly effective in greenhouses.

### C16: WILD PIG

Bells hung up on string to jangle when the wind blows or on trip wires are said to be effective in frightening off wild pig. Strong-smelling human clothes, or urinating around the edge of the field are other options. Pig wire fencing is the most effective solution to these strong and persistent pests.

### C17: VARIOUS INSECTS

- 1 Light traps are effective in catching grasshoppers and a variety of night-flying insects, for example moths, mosquitoes, chafer beetles, American bollworm and weevils. One suitable trap is shown in the sketch below. An objection to light traps is that they catch many useful insects as well and should therefore be used only in an emergency.
- 2 A wide variety of crawling insects can be caught by placing baits such as crushed maize, fruit or vegetable matter into the bottom of tin cans and stuffing the cans with straw. Place upside down on the ground or on a stick and remove the insects hiding



- inside at regular intervals.
- 3 Yellow bars covered with sticky materials such as oils are said to trap aphids, leafhoppers, black flies, gnats and hornets. Yellow and blue ones have been used for thrips. Sticky red balls coated with attractive smells have been employed to trap fruit flies in orchards.

## PART IV

# REMEDIES FOR PROTECTING STORED SEED AND GRAIN

### Note to the reader

As a direct result of the increasing use of insecticides over the last few decades, traditional methods of protecting stored seed and grain are being forgotten to the extent that many farmers are now unaware that low-cost alternatives exist.

Many traditional methods have been compiled in this section to preserve for future generations the ancient art of protecting stored seed and grain.

## A Controlling pests in stored seed (Small amounts for replanting)

In all cases, store the seed for replanting in strong, airtight containers in a cool, dark, dry place. Weevils and rodents can penetrate all but the thickest plastic. Glass or tin containers are the best. Ensure that the seed is thoroughly dry and hard inside before it is stored otherwise it will rot.

### 1 ASHES, OTHER FINE POWDERY MATERIALS AND SAND

Place the seed in the container until it is about two-thirds full. Add the powdery material and shake so that all voids are filled with the powder/ash. Make sure the top of the seed is also covered then fix the lid firmly on.

Ashes from plants (wood, leaves and stalks), agricultural lime, crushed clay, laterite, coal and charcoal, chalk, cow dung and rock dust prevent insects from moving from seed to seed and smother adults and eggs.

Seeds soaked in water and wood ash for 24 hours before planting or seed stored in dry cowdung and wood ash are protected against fungal and bacterial diseases, and are less likely to be dug up by birds when planted out in the field.

Fine sharp-grained sand has also been found to protect stored grain by cutting the insect and causing it to dehydrate.

### 2 BUTTERMILK

Dipping seeds in buttermilk and allowing them to dry gives protection against fungal and viral diseases.

### 3 COLD

Exposing the seed to temperatures below 5 °C for several hours will kill off weevils and some other tropical insect pests. Put out seed containers on frosty nights.

### 4 HEAT

Fungi, bacteria and weevils can be killed off by heating the seed to no more than 50°C. Temperatures greater than this will destroy seed viability — see *Part IV B1.7* for further details.

### 5 HOT WATER AND AIR

Seed can be disinfected of the spores of fungi and bacteria by placing the seed (maximum 250g) into a cloth bag, soaking the bag in cold water for 30 minutes followed by 20 minutes in water at 50°C (just too hot to touch), then dipping the bag in cold water again for a few minutes before spreading the seed in the sun to dry rapidly.

Fungi on peaches and lemons has been controlled by dipping the freshly picked fruit in hot water. Dipping times are 7 minutes for 50°C, 3 minutes for 55°C and 2 minutes for 60°C. Treat within 24 hours of harvesting. Hot air can also be used.

### 6 NAPHTHALENE BALLS

Add 2 per cent per kg of stored seeds. The smell will taint grains to be eaten.

### 7 OIL

Oils of coconut, castor bean, cottonseed, groundnut, maize, mustard, safflower, soyabean, sunflowers and leaf mustard affect egg laying, embryo and larva development on the surface of stored seeds. The eggs and larvae die before the latter can bore into the seeds so the seeds have to be properly coated with oil.

Place a little oil in a largish container, fix the lid and shake until the seed is thoroughly coated with a thin oil film. Small, light seeds cannot be treated in this way as they stick in the oil. Seeds of peas, beans, small grains, maize and sorghum are suitable as well as many others.

The seeds have to be coated periodically because the toxic effect of the oil diminishes with time.

### 8 SALT

Small amounts of salt mixed with produce have been found effective.

### 9 SMOKE

Smoke is a preservative agent and insect repellent. You can hang seed pods, cobs and whole seeding plants up to dry in the roof above the fire in village kitchens.



## B Controlling pests in stored grain (Large quantities)

A useful pre-storage approach to minimise damage is to harvest early and then dry the produce quickly. Storage areas should be thoroughly cleaned and repaired before new grain is put into them; and well ventilated. Completely sealing surfaces with packed clay will give protection against rats and mice.

The following methods are to protect the produce in storage after it has been thoroughly dried.

### 1 Non-plant Materials

#### 1.1 Animal fat

Fat coating the grain at 1ml per kg. was found effective against bean weevils.

#### 1.2 Fine Powdery Materials

Mixing 1-3g fine clay, chalk, ground laterite, lime, rock dust, or alumina per kg of stored seeds has been found to give some protection.

Coal ash ground to powder has been very successful and so has powdered charcoal mixed with clay dust.

#### 1.3 Manure

Burn a mixture of dry maize cobs and cowdung and mix with the stored grain or beans at the rate of 1 bucket full (15 litres) of ashes to every 90kg (1 bag) of shelled maize. Burnt cattle manure is said to be effective for 9 months by itself.

Granary floors and walls plastered with cowdung are recommended for pest control. Ash of African sandalwood (see Part III, B7.2) is mixed with cowdung to increase effectiveness.

Goat dung, burned, crushed and mixed with the grain has been used successfully.

In the past, grain storage pits have been dug below ground in cattle pens putting grain into the hot manure, giving good protection to stored grain from weevils for up to one year. A stone was placed on top of the hole but this treatment is said to affect the taste of the grain.

Spraying the walls and floors of granaries with liquid goat dung has also been found to protect grain

from insect attack.

#### 1.4 Peels

The peel from citrus, lemon, grapefruit and limes have proven effective in repelling pests from stored produce.

#### 1.5 Sand

Mix 10kg fine sand with 100kg of stored grain or peas for some protection. The sand can be removed again by sieving and washing.

Ideally the sand should fill the voids in the grain, the grain placed in a metal drum and covered with a layer of 40mm of sand. This is said to protect the grain for about 6 months. Potatoes can be stored in dry sand or soil for months.

#### 1.6 Salt

Salt mixed with pigeon pea gave good protection in one recorded experiment.

#### 1.7 Solar energy

The grain to be disinfested is placed in a large envelope (1m by 1m square or larger) made from black, high density polythene sheeting with the edges folded over to prevent heat or grain loss from inside. The black envelope is then placed inside another envelope made of transparent low density plastic sheeting and the edges turned over as before. The envelopes are placed on sloping ground or a sloping board facing the sun. The edges of the envelope can be weighed down with bricks or wooden batons to prevent them lifting in the wind. The following times and temperatures should be carefully followed. A thermometer is essential.

- a temperature of 45°C maintained for 10 minutes will kill fungi, bacteria, moulds and weevils;
  - a sustained temperature of 50°C will destroy seed viability;
  - a sustained temperature of 60°C will destroy the food value of the grain.
- A temperature of 45°C for 10 minutes is thus

sufficient to protect grain in storage from bacteria, fungi, moulds and weevils but can go up to 60°C for no more than 10 minutes. This higher temperature and time is said to kill off all insects at all stages of development.

#### 1.8 Stirring

Weevil activity can be reduced by stirring, shaking or turning the grain every 2 or 3 weeks.

## 2 Plant Materials

The following plant materials have been reported in the literature as being used traditionally in Africa to protect stored seeds in large quantities.

Basically any strong-smelling plant material can be used to repel insects but some plants have insecticidal properties.

#### 2.1 General

Ashes from plants in general are said to reduce pest attack, with the effectiveness increasing when the ashes are mixed with powdered pyrethrum or syringa seeds. Ashes at 2-4 per cent

by weight of grain are said to give 4-6 months' protection if the moisture content of the grain is less than 11 per cent. Ashes from African sandalwood, casuarina spp, cork bush, derris, mango, pod mahogany, rice husks, scented pod acacia (nilotica) and tamarind being particularly favoured. Grass - burnt grass mixed with millet protected the millet from weevils.

**Oils** - vegetable oil has proven to give long-term protection to beans, cowpea and bambara nuts against weevils for 6 months when mixed with grains at the rate of 5-10 ml per kg of grain.

Oil from cashew nuts, castor bean, coconut, cottonseed groundnuts, maize, mustard, safflower, soyabean, sunflowers and leaf mustard have all proved effective. Oils are usually more appropriate to small or medium amounts of grain.

**Shelled** - Mixing finger millet and cowpea seeds together in storage is said to protect the cowpea.

**Unshelled** - Storing beans unshelled, maize on the cob and millet on the head reduces insect attack.

### 2.2 SPECIFIC PLANTS

Plant Names Botanical/English	Plant Parts	Treatment
<i>Atzelia quanzensis</i> Pod mahogany	Whole plant	Ashes are mixed with grain.
<i>Agave americana</i> American aloe	Whole plant	Parts dried, ground, and dust mixed with the grain.
<i>Andropogon grass spp.</i> Blue/Stab/Snowflake	Leaves	Mixed with grains as a repellent and insecticide.
<i>Annona reticulata</i> Sugar apple	Leaves and crushed seeds at 0.5% by weight	Mixed with sorghum millet and cowpeas
<i>Brassica juncea</i> Leaf mustard	Seeds	10ml of oil per 1 kg of seed mixed in.
<i>Capiscum frutescens</i> Chilli peppers	Ripe, dried pods with ashes, dung or fine clay. (all voids can be filled with clay/pepper/dung mixtures)	Whole pods mixed with grain or dusted as a powder on bean

Plant Names	Plant Parts	Treatment
<i>Catunaregam spinosa</i> Wormseed	Roots, fruit	Dry and grind into a powder and mix with stored food.
<i>Chenopodium ambrosioides</i> Wormseed	Whole plant	Mix 5kg of dried and ground material with every 100kg of seed.
<i>Chrysanth. cinerariifolium</i> Pyrethrum	Flower heads	Pick on hot days: Dry in the shade. Powder and mix with grain.
<i>Combretum imberbe</i> Leadwood	Wood ash	Mix 1 cup of ash to a 20 litre bucket of grain; sometimes with goat dung. Mix ash with cow manure and plaster walls and floors.
<i>Combretum micranthum</i>	Leafy branches	Made into woven mats to cover the inside or layered in the grain.
<i>Orotalaria ochroleuca</i> Sunhemp	Seeds	Mix seeds between all gaps in the stored grain.
<i>Datura stramonium</i> Thorn apple	Leaves and stems	Dry and mix with produce.
<i>Derris elliptica</i> Derris	All parts	Stored produce dusted or sprayed.
<i>Eleusine coracana</i> Finger millet	Leaves or chaff	Layered or mixed with stored produce.
<i>Eucalyptus</i> spp Gums	Leaves and bark	Fresh or powdered; 10-20 leaves per kg of seed recommended.
<i>Hyptis spicigera</i> Black sesame	Leaves	3g leaf powder per kg of stored seed; leafy branches in layers between stored grain.
<i>Lantana</i> spp Lantana	Leaves	Crushed and placed among pods of cowpea.
<i>Lippia javanica</i> Lippia	Leaves, branches	Put layer of 3-5cm leafy branches at bottom of grain Umsuzwane bins.
<i>Mangifera indica</i> Mango	Whole plant, leaves, flowers	Dry, grind into a powder, spread over the surface.
<i>Melissa azedarach</i> Syringa	Leaves, ripe seeds.	Dried, powdered, mixed with stored grain using 2% if powder from seeds & 4% from leaves.
<i>Murdulea senecioidea</i> Dork bush	Fresh or dried powdered leaves and bark.	Mixed with and covering sorghum, millet and cowpeas.
<i>Nicotiana tabacum</i> Tobacco	Leaves	Dried and chopped and mixed with grain.
<i>Cicium</i> spp Wild basil	Whole plant	Layer 3-5cm thick put in the bottom of grain bins.
<i>Sorghum bicolor</i> Sorghum	Chaff/husks	Layered or mixed with grains at 5%.

Plant Names	Plant Parts	Treatment
<i>Spirostachys africana</i> African sandalwood	Wood chips from the core	Mixed with grain gives long term protection. Mix ash with cow manure and plaster walls and floors.
<i>Swartzia madagascariensis</i> Snake bean	Pods	Powdered and mixed with grain.
<i>Tagetes Minuta</i> Mexican marigold	Whole plants	Add dried plant in layers, or mix in powdered plant or place 3-5cm layer of crushed plants in base of grain bins.
<i>Tephrosia vogelii</i> Tephrosia	Whole plant	Powdered or crushed and mixed with grain.
<i>Tulbaghia leucantha</i> Wild garlic	Whole plant	Put fresh crushed plants in storage bins.

## Recommended Reading

The following locally available books are recommended appropriate reading for those interested in improving their knowledge of or intending to practise holistic and sustainable agriculture.

*Holistic Resource Management*, Allan Savoury, Gilmour Publishing 1991  
Available from bookshops. Contains much information on ecological land management, with emphasis on grazing land, pertinent to Southern Africa.

*Introduction To Permaculture*, Bill Mollison, Tutorial Press 1992  
Available from the Permaculture Association Zimbabwe, P.O. Box 8515, Causeway, Harare.

*Production Without Destruction and Management Techniques For Land Use* 1994, Natural Farming Network or The Permaculture Association Zimbabwe. Both available at P.O. Box 8515 Causeway, Harare.

*Soil Conservation*, Henry Elwell, College Press Publishers, *New Directions in Education Series*, 1982  
Available from bookshops or from College Press Publishers P.O. Box 3041, 15 Douglas Road, Workington, Harare.



## PLANT NAMES IN REGIONAL LANGUAGES

LATIN	ENGLISH	SETSWANA	KISWAHILI	NDEBELE	SHONA	CHINYANGA
1.1 <i>Allium cepa</i>	Onion	Anaynyese/Eiye	Kitunguu maji		Hanyanisi	Anyenze
1.2 <i>Allium sativum</i>	Garlic		Kitunguu-saumu			Anyenze
1.3 <i>Cammelia sinensis</i>	Tea	Tee (liquid), Matihare (leaves)	(M)chai			
1.4 <i>Capsicum frutescens</i>	Chilli & sweet peppers	Pherefere	Mpilipili-kichaa & Mpili-hoho	Ibelebile	Mhiripiri	Mphillpill
1.5 <i>Carica papaya</i>	Pawpaw		Mpapal		Mupapaya, Mupopo	Chipapaya
1.6 <i>Chrysanthemum cinerariifolium</i>	Pyrethrum		Pareto			
1.7 <i>Crotalaria juncea</i>	Sunnhemp		Marejea	Umbandatshatsha	Mubotangowa	Mupalazumba
1.8 <i>Eleusine coracana</i>	Finger millet, Rapoko		Ulezi	Mazhovole	Zviyo	Mawele
1.9 <i>Eucalyptus</i> spp.	Gum tree		Mkaralusi		Mupuranga, Mugatiriyi	Bulugamu
1.10 <i>Glycine max</i>	Soya bean		Soya			Soya
1.11 <i>Ipomoea batatas</i>	Sweet potato	Dipotata	Kiazi-kitamu	Imbambayila, Isibula	Mumbamaira	Kachamba
1.12 <i>Lycopersicon esculentum</i>	Tomato	Tamati	(M)nyanya	Itimatisi	Mudomasi	Chinyamanti
1.13 <i>Manihot esculenta</i>	Cassava		Muhogo		Mufarinya	Tute
1.14 <i>Matricaria eximia</i>	Feverfew				Shangiku	
1.15 <i>Melia azedarach</i>	Syringa (Persian lilac)	Mosalaosi, Morulwana	Mposholala (Kipare)		Mutuna, Masyringa	Chikondi
1.16 <i>Nerium indicum</i>	Oleander					
1.17 <i>Nicotiana tabacum</i>	Tobacco	Motsoko	Tumbaku	Igwayi	Fodya	Fodya
1.18 <i>Ocimum basilicum</i>	Sweet basil		Mvumbasi, Vuo	Rukhovi, Amakha		
1.19 <i>Rheum</i> spp.	Rhubarb					
1.20 <i>Tageles erecta</i>	African marigold		Mbangimwitu	Imbanje yonxiwa	Mbanda	
2.1 <i>Agave</i> spp. incl. <i>A. americana</i>	Aloe		Mkonge, Mkatani	Isikutsha	Mukonje	Khonje
2.2 <i>Ageratum conyzoides</i>	Goat weed		Kimavi cha kuku/ Mtumbaku			Chiwanda
2.3 <i>Annona muricata</i>	Soursop		Mstafeli	Ububese	Muroro	
<i>Annona reticulata</i>	Custard apple		Mtomokomkubwa or Mtopetope			
<i>Annona squamosa</i>	Sugar apple		Mtomoko or Mtopetope			
<i>Annona chrysophylla</i>		Mokamanawe, Moko-kele, Mokokwenana, Mukukwenana				

LATIN	ENGLISH	SETSWANA	KISWAHILI	NDEBELE	SHONA	CHINYANGA
2.4 <i>Annona Senegalensis</i>	Custard apple		Mtopetope-mwitu	Ububese	Muroro	Mpovya
2.5 <i>Argemone mexicana/ subfusiformis</i>	Mexican poppy	Lopero		Ijatimani		Chinyaku
2.6 <i>Balanites aegyptiaca</i>	Desert date	Simboba			Muvambangoma	Mubambangoma
2.7 <i>Bersama abyssinica</i>	Bitter bark				Munyahava	Chenjakulo
2.8 <i>Bidens pilosa</i>	Blackjack	Mokolonyane	Kishonanguo	Ucucuza	Muuwu	Kanzota
2.9 <i>Catunaregam spinosa</i>				Isitalaqwa	Chirovadunguru	Chipembele
2.10 <i>Chenopodium - ambrosioides</i>	Wormwood	Motlhatla badimo			Zinhuhwenhuhwe	Kaluhe
2.11 <i>Datura stramonium Datura ferox</i>	Thorn apple, Jimson weed	Mokhure			Chowa	Chivumulo
2.12 <i>Euphorbia hirta</i>	Asthma weed		Mziwasiwa			Nyamatoka
2.13 <i>Euphorbia tirucalli</i>	Rubber hedge	Motsetse, Setihare-setela	Mchakazi, Mgoru, Mnyaa	Lugotsha	Rusungwe	Nkadze
<i>Euphorbia ingens</i>	Candelabra tree	Ngogo				Mlangati
<i>Euphorbia cooperi</i>	Lesser candelabra tree	Modkosa				
2.14 <i>Jatropha curcas</i>	Barbados nut		Mono, Mkaburi		Mupfuta, Munjirimono	Chivumulo
2.15 <i>Lantana camara Lantana rugosa</i>	Lantana	Mabeleadinonyane	Mwingajini	Ubuhobe	Mbarapati, Mubanda	Mfika
2.16 <i>Leonotis nepetifolia</i>	Lion's ear		Mkuwa usiku	Ibhetshulebadala	Mudyatsonzo	Nyasongwe
2.17 <i>Lippia javanica</i>	Lippia	Mosukujwane, Motso-kojane, Mosukudu, Mosibyane	Mpambake	Umsuzwane	Zumbani	Chazi
2.18 <i>Labelia columnaris</i>	God's tobacco		Msambali			
2.19 <i>Lonchocarpus capassa</i> incl. <i>L. bussei</i> , <i>ericalyx</i> & <i>nelsii</i> known as lance-pods	Rain tree	Mopororo		Mucitamuzi	Muparadamusha	Chimpakasa
2.20 <i>Mundulea sericea</i>	Cork bush	Mosilatalou	Utupa-pori, Mkwaja	Umsindandlovu	Mupumhamhene	Mtototo
2.21 <i>Nicandra physalodes</i>	Apple of Peru		Mchekecheke		Mugumachembere	
2.22 <i>Nicotiana rustica Nicotiana glauca</i>	Wild tobacco Tree tobacco	Mosilawankwe, Kgomodimetseng	Tumbaku-mwitu	Igwayi	Fodya	Fodya



LATIN	ENGLISH	SETSWANA	KISWAHILI	NDEBELE	SHONA	CHINYANGA
2.23 <i>Ocimum canum</i> & <i>urticifolium</i>	Basil, hoary & wild	Mokure, Muono	Mvumbasi, Vuo, Mtulle	Amakha	Rukovhi	Kapafumba
2.24 <i>Ricinus communis</i>	Castor oil plant	Mokure, Muono	Mbarika, Mbono or Mnyonyo	Umhlafulutho	Mupfuta	Mono
2.25 <i>Tagetes minuta</i>	Mexican marigold	Mokhane	Mbangimtu	Imbanje yonixwa	Nyakambanje	
2.26 <i>Tephrosia vogelii</i>	Tephrosia		Utupa	Muphumambene	Nyakanyimo, Kachaya, Korekore, Mutika	Mtezeza
3.1-1.6 See previous entries						
4.1 <i>Albizia anthelmintica</i>		Monoga	Mkenge		Musalati	Chigalogolo
4.2 <i>Albizia lebbek</i>	Woman's tongue		Mkenge			
4.3 <i>Aleurites fordii</i>	Tung tree		Mkaa			
4.4 <i>Amaranthus spinosus</i>	Pigweed		Mchicha mwiba	Imbuya	Mowadanga	Maule
4.5 <i>Andropogon</i> spp.	Blue, Stab, Snowflake grass		Nyasi			
4.6 <i>Calotropis procera</i>	Swallow wort		Mpamba mwitu			
4.7 <i>Chenopodium album</i>	Fat hen	Motsebe			Mubvunzandadya	Kaluche
4.8 <i>Clausena anisata</i>			Mjavikali, Mkunguni (Shambaa)		Samanobere, Muvengahonye	Msambansanya
4.9 <i>Combretum imberbe</i>	Leadwood	Motswere	Mlama, Msana	Umtshili	Mutsviri, Muchenarota	Musimbili
4.10 <i>Dichrostachys cinerea</i>	Chinese lantern	Moselesele	Mfunganyamba or Mtunduru	Ugagu	Mupangara	Lumpangala
4.11 <i>Digitaria temata</i>	Finger grass	Moseka, Namele		Umthando	Tsinde	Luba
4.12 <i>Heteropogon contortus</i>	Spear grass	Seloka		Inzala	Tsine	Nsoche
4.13 <i>Khaya nyasica</i> , <i>Khaya anthotheca</i>	Red mahogany		Mkangazi		Mubawa	Mlulu
4.14 <i>Myrothamnus flabelliformis</i>	Resurrection bush	Manna-yoo-nkang			Mufahore, Mufandichimuka	Njeza
4.15 <i>Securinea virosa</i> , <i>Fluggea virosa</i>	Snow berry	Mulingambu, Dijotsa, Dinonyane, Monwana	Mkwamba	Umklankomo	Muchagawuwe	Chipeyela
4.16 <i>Solanum panaduriforme</i>	Sodom apple	Ditholatholwana, Tholwa	Mtula, Ndulele			Katuntula
4.17 <i>Strychnos</i> spp. <i>Strychnos cocculoides</i> <i>Strychnos pungens</i>	Monkey orange	Mogorogwana Mogwagwa, Mutu	Mkwakwa			Mzimbill
4.18 <i>Tilthonia rotundiflora</i>	Red sunflower					